

2006

## Reactions of 1,4-bis(tetrazole)benzenes: Formation of Long Chain Alkyl Halides

Fintan Kelleher

*Technological University Dublin, [fintan.kelleher@tudublin.ie](mailto:fintan.kelleher@tudublin.ie)*

Andrew Bond

*University of Southern Denmark, Odense*

Adrienne Fleming

*Technological University Dublin*

*See next page for additional authors*

Follow this and additional works at: <https://arrow.tudublin.ie/ittsciart>

### Recommended Citation

Kelleher, F., Bond, A. Fleming, A., McGinley, J. and Prajapati, V. Reactions of 1,4-bis(tetrazole)benzenes: formation of long chain alkyl halides. Tetrahedron, deposited on Mar. 2008.

This Article is brought to you for free and open access by the School of Science and Computing at ARROW@TU Dublin. It has been accepted for inclusion in Articles by an authorized administrator of ARROW@TU Dublin. For more information, please contact [arrow.admin@tudublin.ie](mailto:arrow.admin@tudublin.ie), [aisling.coyne@tudublin.ie](mailto:aisling.coyne@tudublin.ie).



This work is licensed under a [Creative Commons Attribution-NonCommercial-Share Alike 4.0 License](https://creativecommons.org/licenses/by-nc-sa/4.0/)  
Funder: Irish Government; Danish Natural Sciences Research Council and the Carlsberg Foundation

---

**Authors**

Fintan Kelleher, Andrew Bond, Adrienne Fleming, John McGinley, and Vipa Prajapati

## Reactions of 1,4-bis(tetrazole)benzenes: formation of long chain alkyl halides.

Andrew D. Bond,<sup>a</sup> Adrienne Fleming,<sup>b</sup> Fintan Kelleher,<sup>b</sup> John McGinley,<sup>c,\*</sup> and Vipa Prajapati.<sup>b</sup>

<sup>a</sup> *Department of Chemistry, University of Southern Denmark, DK-5230 Odense M, Denmark,*

<sup>b</sup> *Department of Science and Advanced Smart Materials Research Center, Institute of Technology, Tallaght, Dublin 24, Ireland,*

<sup>c</sup> *Department of Chemistry, National University of Ireland Maynooth, Maynooth, Co. Kildare, Ireland.*

Corresponding Author : Dr. John McGinley

Address : Department of Chemistry,  
National University of Ireland Maynooth,  
Maynooth,  
Co. Kildare,  
Ireland.


Phone Number : (+353 1) 708 4615

Fax Number : (+353 1) 708 3815

e-mail address : john.mcginley@nuim.ie

## Reactions of 1,4-bis(tetrazole)benzenes: formation of long chain alkyl halides.

Andrew D. Bond, Adrienne Fleming, Fintan Kelleher, John McGinley and Vipa Prajapati.



Reactions of 1,4-bis[2-(tributylstannyl)tetrazol-5-yl]benzene with  $\alpha,\omega$ -dibromoalkanes led to the formation of several alkyl halide derivatives, substituted variously at N1 or N2 on the tetrazole ring. The X-ray crystal structures of a number of derivatives are discussed.

## Reactions of 1,4-bis(tetrazole)benzenes: formation of long chain alkyl halides.

Andrew D. Bond,<sup>a</sup> Adrienne Fleming,<sup>b</sup> Fintan Kelleher,<sup>b</sup> John McGinley,<sup>c,\*</sup> and Vipa Prajapati.<sup>b</sup>

<sup>a</sup> *Department of Chemistry, University of Southern Denmark, DK-5230 Odense M, Denmark,*

<sup>b</sup> *Department of Science and Advanced Smart Materials Research Center, Institute of Technology, Tallaght, Dublin 24, Ireland,*

<sup>c</sup> *Department of Chemistry, National University of Ireland Maynooth, Maynooth, Co. Kildare, Ireland.*

### Abstract

The reactions of 1,4-bis[2-(tributylstannyl)tetrazol-5-yl]benzene with  $\alpha,\omega$ -dibromoalkanes were carried out in order to synthesise pendant alkyl halide derivatives of the parent bis-tetrazole. This led to the formation of several alkyl halide derivatives, substituted variously at N1 or N2 on the tetrazole ring. The crystal structures of 1,4-bis[(2-(4-bromobutyl)tetrazol-5-yl)]benzene (2-*N*, 2-*N'*), 1,4-bis[(2-(4-bromobutyl)tetrazol-5-yl)]benzene (1-*N*, 2-*N'*) and 1,4-bis[(2-(8-bromooctyl)tetrazol-5-yl)]benzene (2-*N*, 2-*N'*) are reported. Further discussion involves the structure of 1,4-bis(2-(6-bromohexyl)-2*H*-tetrazol-5-yl)benzene (2-*N*, 2-*N'*) previously reported.

### Keywords

Tetrazole; organotin; X-ray; alkyl halide; NMR spectroscopy.

## Introduction

The synthesis of tetrazoles from a cycloaddition reaction between a nitrile and an azide is well documented<sup>1-6</sup> since tetrazoles have roles in coordination chemistry as ligands, in medicinal chemistry as metabolically stable surrogates for carboxylic acids and in materials science applications, including photography and explosives. The interest in the ability of tetrazoles to mimic the carboxylic acid group has motivated the incorporation of tetrazoles into biologically active molecules.<sup>2</sup> This potential use has led to the incorporation in therapeutic applications, including their incorporation in pharmacologically active compounds with anti-hypertensive, anti-allergic and antibiotic activities.<sup>3</sup>

In recent years, particular attention has been directed towards the use of polydentate aromatic nitrogen heterocycles, specifically ligands with five-membered rings (azoles). Among these, imidazoles and triazoles have been extensively used for their ability to construct open framework networks with a wide variety of topologies. Tetrazoles exhibit a strong networking ability usually acting as mono- or bidentate ligands in most of the reported complexes.<sup>7-9</sup> A possible application for these materials as molecular hosts is in generating supramolecular arrays which embody additional functional groups capable of metal complexation. This would result in a metallotetrazole framework with potential as new catalysts, anti-bacterial or therapeutic agents. Our interest in macrocycles containing tetrazoles surrounds their use as precursors for the formation of new functionalised poly-tetrazoles as, for example, sensors or in molecular recognition.

Both Molloy *et al.*<sup>5</sup> and Butler and Fleming<sup>10</sup> have synthesised bis(bromoalkyltetrazolyl)benzenes from either tributylstannyl-substituted bis-tetrazoles or *N*-unsubstituted tetrazoles with dihaloalkanes with the 2-*N*, 2-*N'*-isomer again being the predominant product; Molloy *et al.* studied both the 1,2- and 1,3-substituted bis-tetrazole while Butler and Fleming looked only at the 1,3-unsubstituted bis-tetrazoles. In fact, Butler has succeeded in using these bis-(bromoalkyltetrazolyl)-benzenes to generate the tetra-tetrazole macrocycle (Figure 1) which include a cavity of variable dimensions tailored by both the length and flexibility of the alkyl chain and also the substitution position on the benzene ring.<sup>10-12</sup> Furthermore, the X-ray crystal structure of one such macrocycle has been reported.<sup>13</sup>

<< Figure 1 >>

We have previously reported our initial findings regarding the addition of pendant short-chain alkyl halide arms of some bis-tetrazoles which yielded not only bis-tetrazole derivatives with pendant alkyl halide arms but also, and rather surprisingly, bis-tetrazole derivatives with pendant vinyl arms.<sup>14</sup> The X-ray crystal structure of 1,4-bis(tetrazole)benzene with bromohexyl pendant arms has been reported, but with limited structural and spectroscopic discussion.<sup>15</sup> In this paper, we report the reactions of 1,4-bis(tetrazole)benzenes with various long chain  $\alpha,\omega$ -dibromoalkanes and discuss their spectroscopic results. The crystal structures of three derivatives are also presented herein.

## Results and Discussion

While alkylation of 5-substituted mono-tetrazole derivatives is known to lead to mixtures of 1-*N*- and 2-*N*-substituted products,<sup>1</sup> the regioselectivity being dependent on the reaction conditions and the nature of the *C*- and *N*-substituents, the alkylation of bis-tetrazole derivatives can lead to several diverse products. For example, the reaction of 1,2-bis[2-(tributylstannyl)tetrazol-5-yl]benzene (1,2-(Bu<sub>3</sub>SnN<sub>4</sub>C)<sub>2</sub>C<sub>6</sub>H<sub>4</sub>) with 1,2-dibromoethane, reported by Molloy *et al.*,<sup>5</sup> has been shown to form either a cyclophane or bis(bromoethyltetrazolyl)benzenes, depending on the ratio of the dibromoethane employed in the reaction. When using a 10-fold excess, the cyclophane was obtained; a larger excess (25:1) resulted in the formation of the bis(bromoethyltetrazolyl)benzenes, either the 2-*N*, 2-*N'*- or the 1-*N*, 2-*N'*- isomer, with the 2-*N*, 2-*N'*-isomer predominating in a ratio of *ca.* 3:1. We have recently reported the reactions of either 1,*n*-bis(tetrazol-5-yl)benzene (1,*n*-(HN<sub>4</sub>C)<sub>2</sub>C<sub>6</sub>H<sub>4</sub>) or 1,*n*-bis[2-(tributylstannyl)tetrazol-5-yl]benzene (1,*n*-(Bu<sub>3</sub>SnN<sub>4</sub>C)<sub>2</sub>C<sub>6</sub>H<sub>4</sub>) (*n* = 2, 3, 4) with 1,2-dibromoethane which yielded compounds containing pendant bromoethyl or vinyl groups with substitution occurring at either 1-*N*,2-*N'* or 2-*N*,2-*N'* respectively.<sup>14</sup>

Our strategy was to use both of these approaches, that is, the use of both 1,*n*-bis(tetrazol-5-yl)benzene and 1,*n*-bis[2-(tributylstannyl)tetrazol-5-yl]benzene, to obtain sufficient quantities of the 2-*N*, 2-*N'*-isomer of various bis(bromoalkyltetrazolyl)benzenes with a view towards generating derivatised tetra-tetrazole macrocycles. One approach involved the reactions of the 1,4-bis(tetrazole)benzene with long chain  $\alpha,\omega$ -dibromoalkanes. When the reactions of 1,4-bis(tetrazol-5-yl)benzene (1,4-(HN<sub>4</sub>C)<sub>2</sub>C<sub>6</sub>H<sub>4</sub>) with either 1,4-dibromobutane, 1,6-dibromohexane or 1,8-dibromooctane were heated as neat suspensions, no products were obtained and starting materials were recovered in all cases. Furthermore, when the reactions were carried out in the presence of triethylamine in either methanol or toluene with heating to reflux for 24 hours, we found that the

recovered material, after work-up, contained mainly starting bis-tetrazoles, with approx. 30% of products obtained. We were able to grow crystals of 1,4-bis[(2-bromohexyl)tetrazol-5-yl]benzene, prepared by this method, which we have already reported.<sup>8</sup> However, when the reactions of 1,4-bis[2-(tributylstannyl)tetrazol-5-yl]benzene (1,4-(Bu<sub>3</sub>SnN<sub>4</sub>C)<sub>2</sub>C<sub>6</sub>H<sub>4</sub>) with either 1,4-dibromobutane, 1,6-dibromohexane or 1,8-dibromooctane were heated as neat suspensions, two products were obtained, in relatively high yields, in all the reactions, as well as some recovered starting material (Scheme 1), suggesting that the organotin route was the better method for the synthesis of this particular type of material. It should be pointed out that neither the cyclophane product nor any products containing additions on one ring only were obtained.

<< Scheme 1 >>

Column chromatography, using hexane/ethyl acetate mixtures as eluent, separated the products from the reactants. <sup>1</sup>H and <sup>13</sup>C NMR spectra were obtained for all samples and revealed that both the 2-*N*, 2-*N'*- and the 1-*N*, 2-*N'*- isomers of the bis(bromoalkyltetrazolyl)benzene had formed in the reaction, but the expected cyclophane was not present. The isomeric 2-*N*, 2-*N'*- and 1-*N*, 2-*N'*- derivatives are readily distinguishable by their <sup>1</sup>H and <sup>13</sup>C NMR spectra,<sup>5,10,14</sup> with the <sup>13</sup>C NMR chemical shift of the tetrazole carbon atom appearing at *ca.* 154.0 and 164.0 ppm in 1,5- and 2,5-disubstituted tetrazoles, respectively. The symmetrical 2-*N*, 2-*N'*-substituted compounds thus gave rise to a single resonance at 164.0 ppm while both signals were apparent in the 1-*N*, 2-*N'*-substituted compounds. For example, 1,4-bis[(2-bromobutyl)tetrazol-5-yl]benzene (2-*N*, 2-*N'*) (**2a**) has a single peak at 164.6 ppm while 1,4-bis[(2-bromobutyl)tetrazol-5-yl]benzene (1-*N*, 2-*N'*) (**2b**) has two peaks at 153.8 and 164.0 ppm, respectively. The main difference in the <sup>1</sup>H NMR spectra of the isomers was the doubling of signals in the case of the 1-*N*, 2-*N'*-derivatives.

Crystals of compounds **2a**, **2b** and **4a**, suitable for an X-Ray diffraction study, were obtained from chloroform and the structures confirmed the presence of the pendant bromoalkyl groups at the 2-*N*, 2-*N'*-positions for **2a** and **4a** and at the 1-*N*, 2-*N'*-positions for **2b** (Figures 2, 3, and 4). Table 1 shows important bond lengths and angles for the three structures. We have previously published the structure of **3a**,<sup>15</sup> allowing us to include this also in our comparison of the 2-*N*, 2-*N'* compounds.

<< Figure 2 >>

<< Figure 3 >>



<< Figure 4 >>

<< Table 1 >>

The structures of **2a**, **3a** and **4a** exhibit regularity. In each case, the molecular unit is centrosymmetric in the solid state and is sited on a crystallographic inversion centre. The tetrazole rings are essentially coplanar with the benzene ring to which they are attached, and the bromoalkyl groups adopt fully extended conformations, projecting to either side of this plane. The long axes of the alkyl chains form angles of *ca* 145° to the molecular plane (measured by the N4-C5-C7 angle in **2a**, and equivalent angles in **3a** and **4a**). Adjacent molecules adopt slipped  $\pi$ -stacking arrangements, with distances of *ca* 3.4 Å between the least squares planes of the molecules, and centroid-centroid separations of *ca* 4.6 Å (the lattice parameter *a*). The stacks are arranged into 2-D layers so that molecules in adjacent stacks are close to co-planar, introducing relatively short contacts in the range 2.7-2.8 Å between one H atom of the first methylene group in the bromoalkyl chain and the 1-N atom of the neighbouring tetrazole rings. The alignment of neighbouring stacks into a 2-D arrangement is consistent with effective lateral packing of the alkyl chains: in each case, the H atoms of the methylene groups in one chain project into the gaps between methylene groups in the neighbouring chains. The 2-D layer arrangements are effectively identical in each of **2a**, **3a** and **4a**. Adjacent 2-D layers meet so as to bring the terminal CH<sub>2</sub>—Br bonds into an offset co-linear alignment, forming “Type I” Br $\cdots$ Br interactions, as classified previously by Pedireddi *et al.*<sup>16</sup> The intermolecular Br $\cdots$ Br distances lie in the range 3.4802(4)–3.5351(8) Å, considerably shorter than twice the bromine van der Waals radius (3.90 Å) and within the range of those values previously reported, 3.415–3.691 Å.<sup>17-20</sup>

In the crystal structure of **2b**, the two tetrazole rings in each molecule are essentially co-planar (r.m.s. deviation of 10 fitted atoms = 0.007 Å), but the plane of the central benzene ring forms a dihedral angle of 9.7(3)° with this plane. The bromobutyl chains form greater angles to the molecular plane (N3-C9-C11 and N6-C13-C15 both *ca* 95.5°), and the terminal CH<sub>2</sub>—Br bond vectors lie close to perpendicular to each other (Figure 3). The arrangement of the central portions of the molecules in the crystal structure is closely comparable to that in **2a**, **3a** and **4a**. The least-squares planes of the  $\pi$ -stacked molecules are separated by *ca* 3.5 Å, with centroid-centroid separations of *ca* 4.4 Å. Adjacent molecules in the stacks are related by centres of inversion. The

molecules in adjacent stacks are again close to co-planar, but with a slightly greater lateral offset along the long axes of the molecules in **2b** compared to **2a**. This introduces a short H···H contact (*ca* 2.29 Å) between adjacent benzene rings, and the twisting of these rings from the planes of the tetrazole rings may be attributed at least in part to alleviation of the steric constraints associated with this close contact. The terminal CH<sub>2</sub>—Br bonds again form “Type I” Br···Br interactions, with intermolecular Br···Br distances of 3.591(2) and 3.677(2) Å. It is noteworthy that while one of these Br···Br distances is closely comparable to those in **2a**, **3a** and **4a**, the second is significantly longer. This less-than-optimal intermolecular contact presumably contributes to the significantly lower melting point of **2b** (*ca* 110°C) compared to the isomeric **2a** (*ca* 140°C).

## Conclusions

The reactions of 1,4-(Bu<sub>3</sub>SnN<sub>4</sub>C)<sub>2</sub>C<sub>6</sub>H<sub>4</sub> with 1,4-dibromobutane, 1,6-dibromohexane and 1,8-dibromooctane yield compounds containing pendant bromoalkyl groups with substitution occurring at either *1-N,2-N'* or *2-N,2-N'* respectively. The crystal structures of three derivatives were obtained, including both the (*1-N, 2-N'*) and the (*2-N, 2-N'*) isomers of one derivative. No cyclophane products were observed from any of the reactions. Similar reactions involving 1,3-dibromopropane, 1,5-dibromopentane and 1,7-dibromoheptane are currently being investigated, as are macrocyclic ring closure reactions with 1,4-(Bu<sub>3</sub>SnN<sub>4</sub>C)<sub>2</sub>C<sub>6</sub>H<sub>4</sub> and metal complexation reactions with the resulting macrocyclic ring.

## Experimental

<sup>1</sup>H and <sup>13</sup>C NMR ( $\delta$  ppm; *J* Hz) spectra were recorded on a JOEL JNM-LA300 FT-NMR spectrometer using saturated CDCl<sub>3</sub> solutions with Me<sub>4</sub>Si reference, unless indicated otherwise, with resolutions of 0.18 Hz and 0.01 ppm, respectively. Infrared spectra (cm<sup>-1</sup>) were recorded as KBr discs or liquid films between KBr plates using a Nicolet Impact 410 FT-IR. Melting points were measured with a Stuart Scientific melting point apparatus (SMP1) without correction. Microanalysis was carried out at the Microanalytical Laboratory of University College, Dublin. Standard Schlenk techniques were used throughout.

## Syntheses

1,4-(Bu<sub>3</sub>SnN<sub>4</sub>C)<sub>2</sub>C<sub>6</sub>H<sub>4</sub> (**1**) was prepared as described previously.<sup>14</sup> All other reagents were commercially obtained and used without further purification. **CAUTION:** Owing to their potentially explosive nature, all preparations of and subsequent reactions with organotin azides and tetrazoles were conducted under an inert atmosphere behind a rigid safety screen.

The numbering scheme for the 1,4-bis-tetrazoles is shown in the figure below and all NMR assignments are based on these diagrams.

<< Figure 5 >>

### General Synthesis of Compounds

1,4-(Bu<sub>3</sub>SnN<sub>4</sub>C)<sub>2</sub>C<sub>6</sub>H<sub>4</sub> (**1**) (10 g, 126.2 mmol) and the appropriate  $\alpha,\omega$ -dibromoalkane (300 mmol) were heated to 120°C for 24 hours, under nitrogen. After cooling, the excess  $\alpha,\omega$ -dibromoalkane was removed under reduced pressure to afford the mixture of isomers **a** and **b**, as well as some starting bis-tetrazole. These were separated by column chromatography on silica gel (initially at the ratio of hexane : ethyl acetate 80 : 20, followed by the ratio 60 : 40). All compounds were recrystallised from chloroform.

#### 1,4-Bis(2-(4-bromobutyl)tetrazol-5-yl)benzene (2-N, 2-N') (**2a**)

White solid. Analysis: Found: C, 39.61; H, 4.15; N, 22.85. Calc. For C<sub>16</sub>H<sub>20</sub>Br<sub>2</sub>N<sub>8</sub>: C, 39.69; H, 4.16; N, 23.14; Yield: 64 %, 3.91 g, 8.1 mmol;  $\nu_{\max}$  (KBr) 2958, 1609, 1584, 1538, 1503, 1443, 1384, 1357, 1254, 1222, 1190, 1035, 1006, 875, 782, 753 cm<sup>-1</sup>; Mp 138-142 °C;  $\delta_{\text{H}}$ : 1.95 [m, 4 H, CH<sub>2</sub>], 2.75 [m, 4 H, CH<sub>2</sub>], 3.47 [t, 4 H, *J* = 6.6 Hz, CH<sub>2</sub>Br], 4.74 [t, 4 H, *J* = 6.6 Hz, NCH<sub>2</sub>], 8.28 [s, 4 H, H<sup>1</sup>-C<sub>6</sub>H<sub>4</sub>];  $\delta_{\text{C}}$ : 27.8 [CH<sub>2</sub>], 29.2 [CH<sub>2</sub>], 32.2 [CH<sub>2</sub>], 52.3 [CH<sub>2</sub>N], 127.3 [C<sup>1</sup>-C<sub>6</sub>H<sub>4</sub>], 129.1 [i-C<sub>6</sub>H<sub>4</sub>], 164.6 [CN<sub>4</sub>].

#### 1,4-Bis(2-(4-bromobutyl)tetrazol-5-yl)benzene (1-N, 2-N') (**2b**)

White solid. Analysis: Found: C, 39.92; H, 4.13; N, 23.04. Calc. For C<sub>16</sub>H<sub>20</sub>Br<sub>2</sub>N<sub>8</sub>: C, 39.69; H, 4.16; N, 23.14; Yield: 21 %, 1.28 g, 2.6 mmol; Mp 108-112 °C;  $\nu_{\max}$  (KBr) 2929, 2853, 1628, 1558, 1538, 1469, 1438, 1370, 1307, 1260, 1225, 1195, 1112, 1037, 1005, 847, 734, 645 cm<sup>-1</sup>;  $\delta_{\text{H}}$ : 1.95

[m, 4 H, CH<sub>2</sub>], 2.15 [m, 2 H, CH<sub>2</sub>], 2.28 [m, 2 H, CH<sub>2</sub>], 3.40 [t, 2 H, *J* = 6.6 Hz, CH<sub>2</sub>Br], 3.47 [t, 2 H, *J* = 6.6 Hz, CH<sub>2</sub>Br], 4.52 [t, 2 H, *J* = 6.6 Hz, N<sup>1</sup>CH<sub>2</sub>], 4.75 [t, 2 H, *J* = 6.6 Hz, N<sup>2</sup>CH<sub>2</sub>], 7.85 [d, 2 H, *J* = 6.6 Hz, H<sup>2</sup>-C<sub>6</sub>H<sub>4</sub>]; 8.38 [d, 2 H, *J* = 6.6 Hz, H<sup>1</sup>-C<sub>6</sub>H<sub>4</sub>]; δ<sub>C</sub> : 27.8 [CH<sub>2</sub>], 28.2 [CH<sub>2</sub>], 29.1 [CH<sub>2</sub>], 29.2 [CH<sub>2</sub>], 32.0 [CH<sub>2</sub>], 32.1 [CH<sub>2</sub>], 47.3 [CH<sub>2</sub>N], 52.4 [CH<sub>2</sub>N], 125.5 [i-C<sub>6</sub>H<sub>4</sub>], 127.7 [C<sup>1</sup>-C<sub>6</sub>H<sub>4</sub>], 129.3 [C<sup>2</sup>-C<sub>6</sub>H<sub>4</sub>], 130.3 [i'-C<sub>6</sub>H<sub>4</sub>], 153.8 [CN<sub>4</sub>], 164.0 [CN<sub>4</sub>].

#### 1,6-Bis(2-(6-bromohexyl)tetrazol-5-yl)benzene (2-N, 2-N') (**3a**)

White solid. Analysis: Found: C, 44.45; H, 5.20; N, 20.35. Calc. For C<sub>20</sub>H<sub>28</sub>Br<sub>2</sub>N<sub>8</sub>: C, 44.46; H, 5.22; N, 20.74; Yield: 65 %, 4.43 g, 8.2 mmol; Mp 104-106 °C; ν<sub>max</sub> (KBr) 2931, 2862, 1610, 1580, 1543, 1515, 1449, 1396, 1352, 1293, 1250, 1182, 1150, 1082, 1041, 897, 783, 756 cm<sup>-1</sup>; δ<sub>H</sub> : 1.40 [m, 4 H, CH<sub>2</sub>], 1.55 [m, 4 H, CH<sub>2</sub>], 1.88 [m, 4 H, CH<sub>2</sub>], 2.11 [m, 4 H, CH<sub>2</sub>], 3.41 [t, 4 H, *J* = 6.6 Hz, CH<sub>2</sub>Br], 4.69 [t, 4 H, *J* = 6.6 Hz, NCH<sub>2</sub>], 8.28 [s, 4 H, H<sup>1</sup>-C<sub>6</sub>H<sub>4</sub>]; δ<sub>C</sub> : 25.6 [CH<sub>2</sub>], 27.5 [CH<sub>2</sub>], 29.2 [CH<sub>2</sub>], 32.4 [CH<sub>2</sub>], 33.5 [CH<sub>2</sub>], 53.1 [CH<sub>2</sub>N], 127.3 [C<sup>1</sup>-C<sub>6</sub>H<sub>4</sub>], 129.1 [i-C<sub>6</sub>H<sub>4</sub>], 164.5 [CN<sub>4</sub>].

#### 1,6-Bis(2-(6-bromohexyl)tetrazol-5-yl)benzene (1-N, 2-N') (**3b**)

White solid. Analysis: Found: C, 44.40; H, 5.20; N, 20.45. Calc. For C<sub>20</sub>H<sub>28</sub>Br<sub>2</sub>N<sub>8</sub>: C, 44.46; H, 5.22; N, 20.74; Yield: 12 %, 0.82 g, 1.5 mmol; Mp 84-86 °C; ν<sub>max</sub> (KBr) 2930, 2865, 1625, 1550, 1538, 1470, 1436, 1379, 1316, 1262, 1224, 1195, 1110, 1036, 1015, 846, 734, 649 cm<sup>-1</sup>; δ<sub>H</sub> : 1.46 [m, 4 H, CH<sub>2</sub>], 1.84 [m, 4 H, CH<sub>2</sub>], 2.05 [m, 8 H, CH<sub>2</sub>], 3.36 [t, 2 H, *J* = 6.6 Hz, CH<sub>2</sub>Br], 3.41 [t, 2 H, *J* = 6.6 Hz, CH<sub>2</sub>Br], 4.51 [t, 2 H, *J* = 6.6 Hz, N<sup>1</sup>CH<sub>2</sub>], 4.70 [t, 2 H, *J* = 6.6 Hz, N<sup>2</sup>CH<sub>2</sub>], 7.84 [d, 2 H, *J* = 6.6 Hz, H<sup>2</sup>-C<sub>6</sub>H<sub>4</sub>]; 8.32 [d, 2 H, *J* = 6.6 Hz, H<sup>1</sup>-C<sub>6</sub>H<sub>4</sub>]; δ<sub>C</sub> : 25.3 [CH<sub>2</sub>], 25.4 [CH<sub>2</sub>], 27.1 [CH<sub>2</sub>], 27.2 [CH<sub>2</sub>], 28.9 [CH<sub>2</sub>], 29.3 [CH<sub>2</sub>], 32.1 [CH<sub>2</sub>], 32.2 [CH<sub>2</sub>], 33.3 [CH<sub>2</sub>], 33.4 [CH<sub>2</sub>], 47.9 [CH<sub>2</sub>N], 53.0 [CH<sub>2</sub>N], 125.5 [i-C<sub>6</sub>H<sub>4</sub>], 127.7 [C<sup>1</sup>-C<sub>6</sub>H<sub>4</sub>], 129.3 [C<sup>2</sup>-C<sub>6</sub>H<sub>4</sub>], 130.3 [i'-C<sub>6</sub>H<sub>4</sub>], 153.8 [CN<sub>4</sub>], 164.1 [CN<sub>4</sub>].

#### 1,8-Bis(2-(8-bromooctyl)tetrazol-5-yl)benzene (2-N, 2-N') (**4a**)

White solid. Analysis: Found: C, 47.95; H, 6.08; N, 18.48. Calc. For C<sub>24</sub>H<sub>36</sub>Br<sub>2</sub>N<sub>8</sub>: C, 48.33; H, 6.08; N, 18.79; Yield: 68 %, 5.12 g, 8.6 mmol; Mp 88-94 °C; ν<sub>max</sub> (KBr) 2929, 2852, 1629, 1557, 1468, 1430, 1354, 1357, 1266, 1226, 1198, 1048, 1006, 854, 744, 723, 647 cm<sup>-1</sup>; δ<sub>H</sub> : 1.39 [m, 16 H, CH<sub>2</sub>], 1.85 [m, 4 H, CH<sub>2</sub>], 2.08 [m, 4 H, CH<sub>2</sub>], 3.39 [t, 4 H, *J* = 6.6 Hz, CH<sub>2</sub>Br], 4.67 [t, 4 H, *J* =

6.6 Hz, NCH<sub>2</sub>], 8.28 [s, 4 H, H<sup>1</sup>-C<sub>6</sub>H<sub>4</sub>]; δ<sub>C</sub> : 26.3 [CH<sub>2</sub>], 27.9 [CH<sub>2</sub>], 28.5 [CH<sub>2</sub>], 28.7 [CH<sub>2</sub>], 29.3 [CH<sub>2</sub>], 32.7 [CH<sub>2</sub>], 33.9 [CH<sub>2</sub>], 53.3 [CH<sub>2</sub>N], 127.3 [C<sup>1</sup>-C<sub>6</sub>H<sub>4</sub>], 129.1 [i-C<sub>6</sub>H<sub>4</sub>], 164.5 [CN<sub>4</sub>].

#### 1,8-Bis(2-(8-bromooctyl)tetrazol-5-yl)]benzene (1-N, 2-N') (**4b**)

White solid. Analysis: Found: C, 48.16; H, 6.07; N, 18.58. Calc. For C<sub>24</sub>H<sub>36</sub>Br<sub>2</sub>N<sub>8</sub>: C, 48.33; H, 6.08; N, 18.79; Yield: 10 %, 0.75 g, 1.3 mmol; Mp 64-68 °C; ν<sub>max</sub> (KBr) 2929, 2853, 1626, 1550, 1538, 1469, 1434, 1378, 1303, 1271, 1226, 1197, 1112, 1044, 1003, 848, 736, 645 cm<sup>-1</sup>; δ<sub>H</sub> : 1.29 [m, 8 H, CH<sub>2</sub>], 1.37 [m, 8 H, CH<sub>2</sub>], 1.82 [m, 4 H, CH<sub>2</sub>], 1.95 [m, 2 H, CH<sub>2</sub>], 2.09 [m, 2 H, CH<sub>2</sub>], 3.38 [t, 2 H, *J* = 6.6 Hz, CH<sub>2</sub>Br], 3.40 [t, 2 H, *J* = 6.6 Hz, CH<sub>2</sub>Br], 4.46 [t, 2 H, *J* = 6.6 Hz, N<sup>1</sup>CH<sub>2</sub>], 4.69 [t, 2 H, *J* = 6.6 Hz, N<sup>2</sup>CH<sub>2</sub>], 7.82 [d, 2 H, *J* = 6.6 Hz, H<sup>2</sup>-C<sub>6</sub>H<sub>4</sub>]; 8.36 [d, 2 H, *J* = 6.6 Hz, H<sup>1</sup>-C<sub>6</sub>H<sub>4</sub>]; δ<sub>C</sub> : 26.2 [CH<sub>2</sub>], 26.3 [CH<sub>2</sub>], 27.9 [CH<sub>2</sub>], 28.0 [CH<sub>2</sub>], 28.4 [CH<sub>2</sub>], 28.5 [CH<sub>2</sub>], 28.7 [CH<sub>2</sub>], 29.3 [CH<sub>2</sub>], 29.7 [CH<sub>2</sub>], 32.6 [CH<sub>2</sub>], 32.7 [CH<sub>2</sub>], 33.8 [CH<sub>2</sub>], 48.2 [CH<sub>2</sub>N], 53.4 [CH<sub>2</sub>N], 125.7 [i-C<sub>6</sub>H<sub>4</sub>], 127.6 [C<sup>1</sup>-C<sub>6</sub>H<sub>4</sub>], 129.3 [C<sup>2</sup>-C<sub>6</sub>H<sub>4</sub>], 130.3 [i'-C<sub>6</sub>H<sub>4</sub>], 153.8 [CN<sub>4</sub>], 163.8 [CN<sub>4</sub>].

#### X-Ray Crystallography

Suitable crystals of **2a**, **2b** and **4a** for X-ray study were obtained by recrystallisation from chloroform solutions, respectively. Data were collected at 180(2) K on a Bruker Nonius X8 APEX II diffractometer,<sup>21</sup> and a multi-scan correction was applied.<sup>22</sup> The structures were refined against *F*<sup>2</sup> using all data.<sup>23</sup> Hydrogen atoms were placed at calculated positions and refined using a riding model.

**Compound 2a.** *Crystal data:* C<sub>16</sub>H<sub>20</sub>Br<sub>2</sub>N<sub>8</sub>, *M* = 484.22, triclinic, *a* = 4.5722(7), *b* = 5.8805(9), *c* = 18.119(3) Å, α = 96.943(6), β = 96.426(6), γ = 101.287(6)°, *U* = 469.61(13) Å<sup>3</sup>, space group *P*-1, *Z* = 1, μ(Mo-Kα) = 4.334 mm<sup>-1</sup>. 7346 data (1717 unique, *R*<sub>int</sub> = 0.0386) were measured in the range 3.59 < θ < 25.74°. *R*<sub>1</sub>(*I* > 2σ(*I*)) = 0.0321 and *wR*<sub>2</sub>(all data) = 0.0815. Goodness of fit on *F*<sup>2</sup> = 1.06. CCDC No. 604247.

**Compound 2b.** *Crystal data:* C<sub>16</sub>H<sub>20</sub>Br<sub>2</sub>N<sub>8</sub>, *M* = 484.22, triclinic, *a* = 6.7256(14), *b* = 8.1427(19), *c* = 17.186(4) Å, α = 96.972(8), β = 92.193(8), γ = 90.034(8)°, *U* = 933.5(4) Å<sup>3</sup>, space group *P*-1, *Z* = 2, μ(Mo-Kα) = 4.361 mm<sup>-1</sup>. 11127 data (3428 unique, *R*<sub>int</sub> = 0.0607) were measured in the range

$3.59 < \theta < 25.84^\circ$ .  $R_1(I > 2\sigma(I)) = 0.0906$  and  $wR_2(\text{all data}) = 0.2519$ . Goodness of fit on  $F^2 = 1.04$ . CCDC No. 604248.

**Compound 4a.** *Crystal data:*  $C_{24}H_{36}Br_2N_8$ ,  $M = 596.43$ , triclinic,  $a = 4.6665(5)$ ,  $b = 5.6593(5)$ ,  $c = 25.284(2)$  Å,  $\alpha = 92.812(3)$ ,  $\beta = 90.773(4)$ ,  $\gamma = 102.138(3)^\circ$ ,  $U = 651.82(10)$  Å<sup>3</sup>, space group  $P-1$ ,  $Z = 1$ ,  $\mu(\text{Mo-K}\alpha) = 3.138$  mm<sup>-1</sup>. 11318 data (2448 unique,  $R_{\text{int}} = 0.0477$ ) were measured in the range  $3.69 < \theta < 25.84^\circ$ .  $R_1(I > 2\sigma(I)) = 0.0338$  and  $wR_2(\text{all data}) = 0.0863$ . Goodness of fit on  $F^2 = 1.04$ . CCDC No. 604249.

## Supplementary Data

Crystallographic data for **2a**, **2b** and **4a** have been deposited with the Cambridge Crystallographic Data Centre. Copies of this information may be obtained free of charge from deposit@ccdc.cam.ac.uk or www: <http://www.ccdc.cam.ac.uk>

## Acknowledgements

We would like to thank the Postgraduate R&D Skills programme (Technological Sector Research, Strand III, Project Code CRS01/TA02) for financial assistance, and Drs. B. Creaven and B. A. Murray for useful discussions. A. D. B. is grateful to the Danish Natural Sciences Research Council and the Carlsberg Foundation for provision of the X-ray equipment.


## References

1. R. N. Butler in *Comprehensive Heterocyclic Chemistry*; A. R. Katritzky, C. W. Rees and E. F. V. Scriven, Eds.; Pergamon: Oxford, U.K., **1996**, Vol. 4.
2. R. J. Herr, *Bioorg. & Med. Chem.*, **2002**, *10*, 3379.
3. K. Sisido, K. Nabika, T. Isida, S. Kozima, *J. Organomet. Chem.*, **1971**, *33*, 337.
4. M. Hill, M. F. Mahon, J. McGinley, K. C. Molloy, *J. Chem. Soc., Dalton Trans.*, **1996**, 835.
5. P. A. Bethel, M. S. Hill, M. F. Mahon, K. C. Molloy, *J. Chem. Soc., Perkin Trans. 1*, **1999**, 3507.
6. Z. P. Demko, K. B. Sharpless, *J. Org. Chem.*, **2001**, *66*, 7945.
7. A. F. Stassen, H. Kooijman, A. L. Spek, J. G. Haasnoot, J. Reedijk, *J. Chem. Crystallogr.*, **2001**, *31*, 307.

8. A. F. Stassen, M. Grunert, A. M. Mills, A. L. Spek, J. G. Haasnoot, J. Reedijk, W. Linert, *Dalton Trans.*, **2003**, 3628.
9. T. Wu, B.-H. Yi, D. Li, *Inorg. Chem.*, **2005**, *44*, 4130.
10. R. N. Butler, A. F. M. Fleming, *J. Heterocycl. Chem.*, **1997**, *34*, 691.
11. R. N. Butler, K. F. Quinn, B. Welke, *J. Chem. Soc., Chem. Commun.*, **1992**, 1481.
12. R. N. Butler, E. P. Ní Bhrádaigh, *J. Chem. Res. (S)*, **1994**, 148.
13. R. N. Butler, J. McGinley, M. F. Mahon, K. C. Molloy, E. P. Ní Bhrádaigh, *Acta Cryst.*, **2001**, *E57*, 195.
14. A. Fleming, F. Kelleher, M. F. Mahon, J. McGinley, V. Prajapati, *Tetrahedron*, **2005**, *61*, 7002.
15. A. F. M. Fleming, F. Kelleher, M. F. Mahon, J. McGinley, K. C. Molloy, V. Prajapati, *Acta Cryst.*, **2004**, *E60*, 2388.
16. V. R. Pedireddi, D. S. Reddy, B. Satish Goud, D. C. Craig, A. D. Rae, G. Desiraju, *J. Chem. Soc., Perkin Trans. 2*, **1994**, 2353.
17. A. M. Christofi, P. J. Garratt, G. Hogarth, J. W. Steed, *J. Chem. Soc., Dalton Trans.*, **2000**, 2137.
18. N. Kuhn, A. Abu-Rayyan, K. Eichele, S. Schwarz, M. Steinmann, *Inorg. Chim. Acta*, **2004**, *357*, 1799.
19. F. Ruthe, W.-W. Du Mont, P. G. Jones, *Chem. Commun.*, **1997**, 1947.
20. R. Savinsky, H. Hopf, I. Dix, P. G. Jones, *Eur. J. Org. Chem.*, **2001**, 4595.
21. Bruker Nonius. APEX2. Version 1.0-22. Bruker Nonius BV, Delft, The Netherlands, 2004.
22. G. M. Sheldrick, SADABS. Version 2.10. Bruker AXS Inc., Madison, Wisconsin, USA, 2003.
23. G. M. Sheldrick, SHELXTL. Version 6.10. Bruker AXS Inc., Madison, Wisconsin, USA, 2000.

## Schemes

Scheme 1.



$n = 2, 3, 4$

	<b>a</b>	<b>b</b>
<b>2 (n = 2)</b>	2a (64 %)*	2b (21 %)
<b>3 (n = 3)</b>	3a (65 %)	3b (12 %)
<b>4 (n = 4)</b>	4a (68 %)	4b (10 %)

\* % Yield in parenthesis



## Captions

### *Figures*

Figure 1: Schematic of tetratetrazole macrocycle with X = substituted benzene and Y = alkyl chain.

Figure 2: Molecular structure of **2a**, showing the labelling scheme for the crystallographically independent atoms. The non-labelled half of the molecule is related to the labelled half by a centre of inversion. Ellipsoids are represented at 50% probability for non-H atoms.

Figure 3: Molecular structure of **2b**, showing the labelling scheme used. Ellipsoids are represented at 50% probability for non-H atoms.

Figure 4: Molecular structure of **4a**, showing the labelling scheme for the crystallographically independent atoms. The non-labelled half of the molecule is related to the labelled half by a centre of inversion. Ellipsoids are represented at 50% probability for non-H atoms.

Figure 5: Labelling scheme used for central core in the 1,4-bis(tetrazole) derivatives.

## Figures

Figure 1




Figure 2




Figure 3




Figure 4






Figure 5



## Tables

Table 1. Selected bond lengths (Å) and angles (°) for **2a**, **3a** and **4a**.

<i>Bond lengths</i>	<b>2a</b>	<b>3a</b>	<b>4a</b>
C(1)-C(2)	1.389(4)	1.389(4)	1.378(3)
C(2)-C(3)	1.399(4)	1.402(3)	1.395(3)
C(3)-C(4)	1.477(4)	1.463(3)	1.469(3)
C(4)-N(1)	1.352(4)	1.356(3)	1.360(3)
C(4)-N(2)	1.334(4)	1.340(3)	1.323(3)
N(1)-N(3)	1.324(4)	1.321(3)	1.324(3)
N(3)-N(4)	1.325(4)	1.333(3)	1.319(3)
N(4)-N(2)	1.330(3)	1.328(3)	1.329(3)
N(4)-C(5)	1.472(4)	1.463(3)	1.458(3)
<i>Bond Angles</i>	<b>2a</b>	<b>3a</b>	<b>4a</b>
C(4)-N(1)-N(3)	106.1(3)	106.2(2)	105.7(2)
N(1)-N(3)-N(4)	105.9(2)	106.2(2)	106.35(19)
N(3)-N(4)-N(2)	114.2(2)	113.8(2)	113.7(2)
N(4)-N(2)-C(4)	101.4(3)	102.2(2)	102.1(2)
N(2)-C(4)-N(1)	112.4(3)	112.2(2)	112.2(2)