

2014-02-07

## Enzymatic Extraction of High-Value Ingredients From Food Waste

Amit Jaiswal

*Technological University Dublin, amit.jaiswal@tudublin.ie*

Nissreen Abu-Ghannam

*Technological University Dublin, nissreen.abughannam@tudublin.ie*

Follow this and additional works at: <https://arrow.tudublin.ie/schfsehcon>



Part of the [Food Biotechnology Commons](#), [Food Chemistry Commons](#), and the [Food Processing Commons](#)

---

### Recommended Citation

Jaiswal, A.K., & Abu-Ghannam, N. (2014). Enzymatic extraction of high-value ingredients from food waste. Presented at "Waste not Want not – Recovering Value from Food Waste". Teagasc Food Research Centre, Ashtown, Dublin 15, Ireland.

This Presentation is brought to you for free and open access by the School of Food Science and Environmental Health at ARROW@TU Dublin. It has been accepted for inclusion in Conference papers by an authorized administrator of ARROW@TU Dublin. For more information, please contact [arrow.admin@tudublin.ie](mailto:arrow.admin@tudublin.ie), [aisling.coyne@tudublin.ie](mailto:aisling.coyne@tudublin.ie), [vera.kilshaw@tudublin.ie](mailto:vera.kilshaw@tudublin.ie).

Funder: The authors gratefully acknowledge the financial support from the Food Institutional Research Measure (FIRM) strategic research initiative administered by the Irish Department of Agriculture, Fisheries and Food.

# Enzymatic Extraction of High-value Ingredients from Food Waste

**Dr. Amit Kumar Jaiswal**  
**Prof. Nissreen Abu-Ghannam**



**School of Food Science and Environmental Health,  
Dublin Institute of Technology, Dublin 1, Ireland**



Brewer's Spent Grain

# FOOD WASTE FACTS



- Food waste: concerns from an economical, environmental and food security perspectives.
- Over 1 million tonnes of food waste produced annually in Ireland.
- 89 million tonnes food waste produced annually throughout the Europe.
- Roughly, 1.3 billion tonnes food per annum is being wasted.

- Common utilization of food waste:
  - incorporation in animal feed
  - direct disposal in landfills




# REGULATIONS



- Government pressure on food industry to reduce the food waste.
- The management of food wastes is becoming extremely difficult due to legislative restrictions on landfill.
- In Europe, policies are created to promote the prevention of food waste in all the life stages:
  - Integrated product policy (EC, 2003)
  - Resource efficiency flagship initiative (EC, 2011) and
  - the bio-economy communication (EC, 2012)

# FOOD WASTE INTO ANIMAL FEED



- Some of the food waste is used as animal feed.
- Spent grain: up to 93% used as animal feed.

- Some of the food waste is not very ideal for animal feed.
  - Low in protein content
  - High in lignin content (olive waste)
  - High in potassium (potato) (only suitable for cattle feed)



# ADDING VALUE TO FOOD WASTE

- Offer a range of potentially useful alternatives for dealing with food waste other than disposal and/or landfilling.
- Basic valorisation strategies including composting, recycling and burning are able to recover/convert less than 50% of the waste into useful products.
- Novel strategies: physical, biological and enzymatic technologies
- These include (but are not restricted to):
  - Ultrasound-assisted extraction of useful components
  - Biological (e.g. fermentation) and
  - Enzymatic approaches for the isolation of valuable components

# WHAT HIGH ADDED-VALUE COMPONENTS??

- Lead to possibilities for the production of valuable products with a market value.
- These include:
  - Enzymes (Amylase, Protease etc.)
  - Nutraceuticals
  - Functional foods
  - Food preservatives (Polyphenols)
  - Organic acids (Lactic acids etc)
  - Food additives
  - Pharmaceutical products



# ONGOING RESEARCH IN OUR LAB

- Our goal is to obtain products of high added value from food waste.
- Most appropriate treatments and technologies from the environment friendly and cost-effective standpoints.
- In pursuit of waste recovery as a better alternative to waste disposal our research is focused on:
  - Biochemical characterization of waste and industrial by-products.
  - Enzymatic and microbial techniques for extraction of high added-value components.
  - Purification and characterization of high added-value components.
  - Functional characterization.
  - Cytotoxicity analysis



# $\beta$ - GLUCAN FROM BREWER'S SPENT GRAIN

- Brewer's spent grain: main waste product from beer production (85%)
- 20 kg per 100 litres beer produced
- 160,000 tonnes Brewer's spent grain produced in Ireland annually
- Rich in fibre, protein, carbohydrate and phenolic compounds
- High availability, low cost and proven safety for human consumption.
- Applications:
  - $\beta$ -glucans are the only dietary fiber recognized by the EFSA to be able to reduce a disease risk and also used to boost the immune system.
  - Food supplement and as an ingredient in product formulation.
  - Non-caloric thickening and stabilizing agents.

# CHITIN FROM MUSHROOM'S WASTE

- About 55,000 tonnes of mushrooms produced in Ireland annually.
- Waste ranges between 5 and 20% of production volume.
- Mushroom waste generally used as compost.
- Due to its physical and chemical properties, chitin is being used in different products and applications:
  - Emulsifiers, thickening, and gelling agent for stabilizing foods
  - Used as a dietary fibre in baked foods
  - Paper and textile industry
  - Cosmetics and biomedical industry

# GLYCOALKALOIDS FROM POTATO PEEL

- Potato production in 2010 was more than 450,000 tonnes in Ireland.
- Industrial processing of potatoes generates large quantities of peel that create disposal, sanitation, and environmental problems.
- The potato peels are rich in glycoalkaloids such as  $\alpha$ -solanine and  $\alpha$ -chaconine, carbohydrates, high in starch (8-28%) and about 1-4% protein.
- Has strong inhibitory effects on fungal and insect pests and can be exploited as natural pesticides.
- Medicinal properties

# PROBIOTIC BEVERAGES

- Development of probiotic beverages based on Bewer's spent grain as substrate
- Supported prolific growth of *L. brevis*, *L. plantarum* and *L. rhamnosus* up to 10.4 log cfu/ml
- Production of 2.95 g/l lactic acid
- Release of 268.6 mg gallic acid equivalent of phenolic compounds
- Rich in antioxidant properties
- Shelf stable in terms of bioactive components and probiotic counts for 15 days

# TRADITIONAL EXTRACTION APPROACH

- To date, most of the extraction procedures are solvent based.
- Solvent has negative impact on cost and environment.
- Possible solvent contamination of the final products leads to the health and safety issues.
- Focus moved to the safe technology which uses very small or no organic solvent.




# BIOLOGICAL EXTRACTION



- Eco-friendly extraction technologies
- Faster extraction
- Higher recovery
- Reduced solvent usage
- Lower energy consumption and

# ENZYME-ASSISTED EXTRACTION...



Enzyme-assisted extraction

- Highly specific and versatile
- Enzymes are not changed by the reaction
- Re-used again for the same reaction with other molecules
- Very little enzyme needed to help in many reactions
- Used only temporarily
- Relatively cheaper than organic solvents

# Thank You

**Dr. Amit Kumar Jaiswal**

Phone: +353 1402 4547

E-mail: [amit.jaiswal@dit.ie](mailto:amit.jaiswal@dit.ie)

**Prof. Nissreen Abu-Ghannam**

Phone: +353 1402 7570

E-mail: [nissreen.abughannam@dit.ie](mailto:nissreen.abughannam@dit.ie)