2020

A Comparison of Dietary Intake of Children/Adolescents with and without Myopia

Niamh Burke

Technological University Dublin, Niamh.Burke@TUDublin.ie

John Butler

Technological University Dublin, john.s.butler@tudublin.ie

Ian Flitcroft

Technological University Dublin, ian.flitcroft@dit.ie

See next page for additional authors

Follow this and additional works at: <https://arrow.tudublin.ie/scschphycon>

Part of the [Optometry Commons](#)

Recommended Citation

Burke, N., Flitcroft, I., & Butler, J. (2020). A comparison of dietary intake of children/adolescents with and without myopia. *Proceedings of the Nutrition Society, Nutrition Society Live 2020*, vol. 79, no. OCE3, E790, 14–15th July 2020. doi:10.1017/S0029665120007764

This Conference Paper is brought to you for free and open access by the School of Physics & Clinical & Optometric Science at ARROW@TU Dublin. It has been accepted for inclusion in Conference Papers by an authorized administrator of ARROW@TU Dublin. For more information, please contact arrow.admin@tudublin.ie, aisling.coyne@tudublin.ie, gerard.connolly@tudublin.ie.

Authors

Niamh Burke, John Butler, Ian Flitcroft, and James Loughman

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/346278793>

A comparison of dietary intake of children/adolescents with and without myopia

Article in *Proceedings of The Nutrition Society* · October 2020

DOI: 10.1017/S0029665120007764

CITATIONS

0

READS

24

4 authors:

Niamh Burke

Technological University Dublin - City Campus

5 PUBLICATIONS 3 CITATIONS

[SEE PROFILE](#)

Ian Flitcroft

Children's University Hospital, Temple Street

77 PUBLICATIONS 1,875 CITATIONS

[SEE PROFILE](#)

John S Butler

Technological University Dublin - City Campus

105 PUBLICATIONS 1,641 CITATIONS

[SEE PROFILE](#)

James Loughman

Technological University Dublin - City Campus

114 PUBLICATIONS 1,109 CITATIONS

[SEE PROFILE](#)

Some of the authors of this publication are also working on these related projects:

Environmental Sustainability and Health Institute [View project](#)

Mozambique Eye care project [View project](#)

A comparison of dietary intake of children/adolescents with and without myopia

N. Burke¹, I. Flitcroft², J.S. Butler³ and J. Loughman¹

¹Centre for Eye Research Ireland, School of Physics, Clinical and Optometric Sciences, College of Sciences and Health, Technological University Dublin, Dublin, Ireland,

²Ophthalmology Department, Temple Street Children's University Hospital, Dublin, Ireland and

³School of Mathematical Sciences, College of Sciences and Health, Technological University Dublin, Dublin, Ireland

Myopia is caused by an eye which has grown too long. The prevalence of myopia has increased drastically in recent years and half of the world's population is expected to be myopic by 2050.⁽¹⁾ Consequently, a significant increase in sight threatening diseases will follow.⁽¹⁾ Myopia most commonly develops in youth, and we now know that environmental factors affect its development. Nutritional status strongly influences growth and development, including that of the eye. It has been hypothesised, therefore, that nutrition may play a role in this excessive eye growth. Current findings are conflicted, with a clear gap in the evidence.^(2,3) Therefore, the aim of this study is to examine differences in dietary intake of myopic and non-myopic Irish children/adolescents, and identify any potential relationship between macro and/or micronutrient intake and myopia.

A hundred healthy children/adolescents aged 6–16 years were recruited into the study, seventy myopes and thirty non-myopes. Each participant was asked to complete two 24hr diet recalls online, using the 'INTAKE24' software. Spherical equivalent refraction was measured by non-cycloplegic auto refraction. Myopia was defined as ≤ -1.00 dioptres [D]. Energy mis-reporters were identified using Goldberg et al cut-offs. Descriptive analysis was preformed using SAS (Version 9.4). Nutrient intakes were reported as medians and interquartile range, as they were not normally distributed. Dietary data was log transformed, and the residual method was used to adjust for energy intake. Multivariate logistic regression was carried out to examine any potential relationships between nutrient intake and myopia.

Both the myopic and non-myopic groups had similar energy, protein, carbohydrate and total fat intake. However, the non-myopes had a significantly higher intake of vitamin D ($p < 0.01$), retinol ($p = 0.04$), cholesterol ($p = 0.01$) and vitamin B12 ($p = 0.02$). When multivariate logistic regression analysis was performed, vitamin D was the only nutrient to maintain a significant association with myopia, after adjustment for potential confounders [OR 0.18 (95% CI 0.03–0.89) ($p = 0.03$)].

Preliminary results show a statistically significant relationship between dietary vitamin D intake and myopia. Interestingly, previous studies have found an association between vitamin D status and myopia,⁽⁴⁾ but few have examined dietary intake alone, indicating a potential difference in diets of myopes and non-myopes. This study does have its limitations; the sample size is small, the use of a dietary intake tool is dependent on recall, and data on dietary supplement use was not available. However, given the corroboration of previous findings, further investigation is warranted, perhaps in a larger population with an extended focus on dietary patterns.

1. Holden BA, Fricke TR, Wilson DA *et al.* (2016) *Ophthalmology* **123**, 1036–1042.
2. Fedor M, Socha K, Urban B *et al.* (2017) *Biol Trace Elem* **176**, 1–9.
3. Lim LS, Gazzard G, Low YL, *et al.* (2010) *Ophthalmology* **117**:993–997.e4.
4. Yazar S, Hewitt AW, Black LJ, *et al.* (2014) *Invest Ophthalmol Vis Sci* **55**:4552–9.