

2006-01-01

Apoptosis is Initiated in Human Keratinocytes Exposed to Signalling Factors from Microbeam Irradiated Cells

Fiona Lyng

Technological University Dublin, Fiona.lyng@tudublin.ie

Paula Maguire

Technological University Dublin

A. Kilmurray

Technological University Dublin

Carmel Mothersill

Technological University Dublin

C. Shao

Mount Vernon Hospital

Follow this and additional works at: <https://arrow.tudublin.ie/radart>

See next page for additional authors

Recommended Citation

Lyng, F. et al. (2006) Apoptosis is Initiated in Human Keratinocytes Exposed to Signalling Factors from Microbeam Irradiated Cells. *International Journal of Radiation Biology*, Vol. 82, no.6, pp.393-399.
doi:10.1080/09553000600803904

This Article is brought to you for free and open access by the Radiation and Environmental Science Centre at ARROW@TU Dublin. It has been accepted for inclusion in Articles by an authorized administrator of ARROW@TU Dublin. For more information, please contact yvonne.desmond@tudublin.ie, arrow.admin@tudublin.ie, brian.widdis@tudublin.ie.



This work is licensed under a [Creative Commons Attribution-Noncommercial-Share Alike 3.0 License](https://creativecommons.org/licenses/by-nc-sa/3.0/)

Authors

Fiona Lyng, Paula Maguire, A. Kilmurray, Carmel Mothersill, C. Shao, M. Folkard, and K. Prise

1
2
3 **Apoptosis is initiated in human keratinocytes exposed to signalling factors from**
4 **microbeam irradiated cells**
5
6

7
8 Lyng FM^{a,1}, Maguire P^a, Kilmurray N^a, Mothersill C^{a,2}, Shao C^b, Folkard M^b and Prise
9
10 KM^b
11

12 ^aRadiation and Environmental Science Centre, Focas Institute, Dublin Institute of
13 Technology, Kevin St, Dublin 8, IRELAND
14
15

16 ^bCell and Molecular Biophysics Group, Gray Cancer Institute, Mount Vernon Hospital,
17 Northwood, UK
18
19
20
21

22
23
24 **Running Head:** Bystander factor induced apoptosis in microbeam irradiated cells
25
26

27 **Keywords:** Radiation, Bystander Effects, Cell Signalling, Apoptosis
28
29
30
31

32 ¹Person to whom all correspondence should be sent:
33

34 Dr Fiona M Lyng
35

36 Radiation and Environmental Science Centre
37

38 Focas Institute
39

40 Dublin Institute of Technology
41

42 Kevin St
43

44 Dublin 8
45

46 IRELAND
47

48 Email: fiona.lyng@dit.ie
49
50
51
52
53

54 ²Present address: Medical Physics & Applied Radiation Sciences, McMaster University,
55 Hamilton, Ontario, Canada
56
57
58
59
60

Abstract

Purpose: There is now no doubt that bystander signalling from irradiated cells occurs and causes a variety of responses in cells not targeted by the ionising track. However, the mechanisms underlying these processes are unknown and the relevance to radiotherapy and risk assessment remains controversial. Previous research by our laboratory has shown bystander effects in a human keratinocyte cell line, HPV-G cells, exposed to medium from γ irradiated HPV-G cells. The aim of this work was to investigate if similar mechanisms to those identified in medium transfer experiments occurred in these HPV-G cells when they are in the vicinity of microbeam irradiated cells. Demonstration of a commonality of mechanisms would support the idea that the process is not artifactual.

Materials and Methods: HPV-G cells were plated as two separate populations on mylar dishes. One population was directly irradiated using a charged particle microbeam (1 - 10 protons). The other population was not irradiated. Bystander factor induced apoptosis was investigated in both populations following treatment by monitoring the levels of reactive oxygen species and mitochondrial membrane potential using fluorescent probes. Expression of the anti-apoptotic protein, bcl-2, and cytochrome c were determined, as well as apoptosis levels.

Results: Microbeam irradiation induced increases in reactive oxygen species and decreases in mitochondrial membrane potential at 6 hours post exposure, increased expression of bcl-2 and cytochrome c release at 6.5 hours and increased apoptosis at 24 hours.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Conclusion: This study shows that similar bystander signalling pathways leading to apoptosis are induced following microbeam irradiation and following medium transfer. This demonstrates that the mechanisms involved are common across different radiation qualities and conditions and indicates that they may be relevant *in vivo*.

For Peer Review Only

Introduction

There has been considerable interest recently in non-targeted effects of radiation that cannot be attributed to direct DNA damage. Recent research has shown that low doses of ionising radiation can cause a “bystander effect”, where radiation damage occurs in cells not directly irradiated (see reviews; Mothersill and Seymour 2003, Lorimore *et al*, 2003, Little 2003, Morgan 2003).

Studies with very low doses of α particles have shown increases in the number of sister chromatid exchanges (SCE) (Nagasawa and Little 1992, Deshpande *et al*, 1997), chromosomal aberrations (Lorimore *et al*, 1998) and activation of stress inducible signalling pathways (Azzam *et al*, 1998, 2001, 2003) in considerably more cells than would have been traversed by an α particle.

Microbeams, which allow the targeting of individual cells or subcellular locations within a population with a charged particle beam, have become useful tools in the study of radiation induced bystander effects. Microbeam studies have shown that effects of single cell irradiation are not limited to the exposed cell but affect other cells in the vicinity. Micronucleus formation and apoptosis (Prise *et al*, 1998, Belyakov *et al*, 2001, 2003), mutation (Zhou *et al*, 2000, 2001) and oncogenic transformation (Sawant *et al*, 2001) have all been shown in cells distant from the target cell.

Medium transfer experiments have shown that bystander effects may be mediated by damage signals released into the culture medium by irradiated cells. Non-irradiated cells incubated with conditioned medium from irradiated cells have shown similar effects to the directly irradiated cells. Mothersill and Seymour (1997, 1998) reported that cell death was induced in unirradiated cells treated with medium from directly irradiated

1
2
3 epithelial cells. This was further shown to be associated with early apoptotic events such
4
5 as calcium fluxes, loss in mitochondrial membrane permeability and the induction of
6
7 reactive oxygen species (ROS) (Lyng *et al*, 2000, 2002). Lehnert *et al* (1997) and
8
9 Narayanan *et al* (1997) also used medium transfer experiments to show that extracellular
10
11 factors including ROS were released by α particle irradiated cells leading to increased
12
13 SCE in nonirradiated cells. Conditioned medium from α particle irradiated cells has also
14
15 been shown to stimulate proliferation in nonirradiated cells (Iyer *et al*, 2000). A recent
16
17 study by Suzuki *et al* (2004) demonstrated that cells irradiated with α particles released
18
19 medium borne factors which induced chromatin damage in bystander cells plated on the
20
21 other side of a medium filled double mylar dish.
22
23
24
25
26
27
28
29

30 To date the majority of reports on bystander effects have used either direct exposure to
31
32 high linear energy transfer (LET) helium ions delivered by low fluence sources
33
34 (Nagasawa and Little 1992), direct irradiation using microbeam approaches (Prise *et al*,
35
36 1998, Zhou *et al*, 2000, Sawant *et al*, 2001, Shao *et al*, 2003) or medium transfer after
37
38 low LET exposure (Mothersill and Seymour 1997).
39
40

41 Two main models have emerged on the mechanisms of bystander responses; cell - cell
42
43 communication through gap junctions and secretion of a cytotoxic factor into the
44
45 medium. The model appears to depend on the cell type used for the experiments and on
46
47 the way the experiments were performed. Some groups have demonstrated a requirement
48
49 for gap junctional intercellular communication (GJIC) to mediate a bystander response
50
51 (Azzam *et al*, 2001, Shao *et al*, 2003b) but other groups have shown GJIC not to be
52
53 involved (Mothersill and Seymour 1997, Lehnert *et al*, 1997).
54
55
56
57
58
59
60

1
2
3 A role for ROS in radiation induced bystander effects has been reported by many groups
4
5
6 (Lehnert *et al* 1997, Azzam *et al* 2002, Lyng *et al* 2006). Specifically, nitric oxide, an
7
8 important signalling molecule, has been shown to induce bystander effects (Matsumoto *et*
9
10 *al* 2000, Shao *et al* 2001, 2002, 2003a, 2004)

11
12
13
14
15 The aim of the present study was to investigate bystander responses following microbeam
16
17 irradiation in a human keratinocyte cell line (HPV-G cells) which has previously been
18
19 shown to undergo apoptosis when exposed to medium from γ irradiated cells (Lyng *et al*,
20
21 2000, 2002, Maguire *et al*, 2005). Mitochondrial membrane potential depolarisation,
22
23 cytochrome c release, bcl-2 expression, induction of reactive oxygen species (ROS) and
24
25 apoptosis levels were measured in HPV-G cells which were in the vicinity of microbeam
26
27 irradiated HPV-G cells.
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Materials and Methods

Cell Culture

Human keratinocytes, HPV-G cells, immortalised with the human papilloma virus (HPV) were originally obtained as a kind gift from Dr. J. Di Paolo, NIH Bethesda (Pirisi *et al*, 1988). HPV-G cells were cultured in Dulbecco's Modified Eagle Medium (DMEM) : F12 (1:1) medium (Sigma, Dorset, U.K.) containing, 10% fetal bovine serum (Gibco, Irvine, U.K.), 1% penicillin-streptomycin solution 1000 IU (Gibco, Irvine, U.K.), 2mM L- glutamine (Gibco, Irvine, U.K.) and 1µg/ml hydrocortisone (Sigma, Dorset, U.K.) Cells were maintained in an incubator at 37°C, with 95% humidity and 5% CO₂. Subculture was routinely performed when cells were 80-100% confluent, using a 1:1 solution of 0.25% trypsin and 1mM versene (Sigma, Dorset, U.K.) at 37° C.

Microbeam irradiation

For microbeam experiments, plateau phase cells were seeded into specially designed dishes (Folkard *et al*, 1997) consisting of a 34 mm diameter base composed of a 4 µm thick mylar membrane. Two areas of the dish diagonally opposed to each other had been pretreated with 1 µg/ml CellTak adhesive (Becton Dickinson, Oxford, U.K.). Each area was about 5 mm in diameter. Cells were seeded 16 h prior to irradiation to allow full attachment. Typically cells were seeded at a density to allow 600–800 cells in each area of the dish. One hour prior to irradiation cells were incubated with 1 µM Hoechst 33258 (Molecular Probes, Leiden, The Netherlands). At the time of irradiation the cell culture medium was replaced with fresh medium containing 20 mM 4-(2-hydroxyethyl)-1-piperazineethanesulfonic acid (HEPES) (Sigma, Dorset, U.K.) and irradiation was

1
2
3 performed at room temperature. All the cells in one area of the dish had exact numbers of
4
5 3.2 MeV protons delivered through the centre of the cell nucleus. Details of the automatic
6
7 cell finding and imaging system are described in Folkard *et al* (1997). The cells in the
8
9 unirradiated area of the dish were bystander cells. The irradiation procedure typically
10
11 took around 10 min after which fresh medium was added to the cells and incubation
12
13 continued at 37°C for 6 or 24 hours prior to scoring. Control cells were sham irradiated
14
15 by incubating with 1 µM Hoechst 33258 (Molecular Probes, Leiden, The Netherlands) as
16
17 before and scanning, finding and following the same number of cells as in the irradiated
18
19 dishes but without actual irradiation.
20
21
22
23
24
25
26

27 *Measurement of mitochondrial membrane potential*

28
29 Mitochondrial membrane potential was determined using rhodamine 123, a green
30
31 fluorescent dye that accumulates in active mitochondria with high membrane potential.
32
33 Cultures were washed twice with a buffer containing 130 mM NaCl, 5 mM KCl, 1 mM
34
35 Na₂HPO₄, 1 mM CaCl₂, 1 mM MgCl₂ and 25 mM HEPES (pH 7.4). Cells were loaded
36
37 with 5 µM Rhodamine 123 (Sigma, Dorset, U.K.) for 30 min in the buffer at 37 °C.
38
39 Subsequently, the cultures were washed three times with buffer. Fluorescence images
40
41 were recorded using a Zeiss Axioskope epifluorescence microscope with a BP 450-490
42
43 nm excitation filter and LP515 emission filter (Carl Zeiss Ltd, Welwyn Garden City,
44
45 U.K.) and a cooled charge coupled device (CCD) camera system (Photonic Science, UK).
46
47 The mean fluorescence intensity (or mean grey value) was determined using the software
48
49 package, NIH Image (National Institutes of Health, USA).
50
51
52
53
54
55
56
57
58
59
60

Measurement of reactive oxygen species

Induction of reactive oxygen species was measured using 2,7 - dichlorofluorescein diacetate (DCF-DA) (Sigma, Dorset, U.K.). Once inside a cell, the acetate group is cleaved by cellular esterases leaving dichlorofluorescein which emits green fluorescence when oxidized by the reactive oxygen species, hydrogen peroxide and nitric oxide. Cultures were washed twice with a buffer containing 130 mM NaCl, 5 mM KCl, 1 mM Na₂HPO₄, 1 mM CaCl₂, 1 mM MgCl₂ and 25 mM HEPES (pH 7.4). Cells were loaded with 5 µM 2,7 DCF-DA for 30 min in the buffer at 37 °C. Subsequently, the cultures were washed three times with buffer. The cells were then returned to the incubator for 30 mins and washed once more prior to scoring. Fluorescence images were recorded using a Zeiss Axioskope epifluorescence microscope with a BP 450-490 nm excitation filter and LP515 emission filter (Carl Zeiss Ltd, Welwyn Garden City, U.K.) and a cooled CCD camera system (Photonic Science, UK). The mean fluorescence intensity (or mean grey value) was determined using the software package, NIH Image (National Institutes of Health, USA).

Immunocytochemistry

Following the fluorescence measurements as described above (ie. approx 6.5 hours after microbeam irradiation), the cells were washed twice in phosphate buffered saline (PBS) to remove any debris and then fixed in 10% buffered formalin. The mylar on which the cells were attached was removed from the microbeam dish and adhered to a glass slide with the cells facing upwards. The slides were marked to indicate the irradiated and

1
2
3 bystander areas. The slides were immersed in PBS in plastic coplin jars and transported
4
5 back to Dublin, stored at 4°C and assayed within 48 hours. Immunocytochemical analysis
6
7 was performed using the Streptavidin Peroxidase method for cell culture using the
8
9 Vectastain ABC kits (Vector Laboratories, UK). The primary antibody, mouse
10
11 monoclonal, anti-Bcl-2 (Dako, Denmark) or anti-cytochrome c (Dako, Denmark) was
12
13 applied for one hour (1:50 and 1:250 dilution respectively). Biotinylated anti-mouse
14
15 reagent was then added to the cells for 30 minutes followed by Streptavidin Peroxidase for
16
17 a further 30 minutes, with a wash in PBS in between each step. The chromagen, 0.02%
18
19 DAB (Sigma, Dorset, U.K.) was then added for 10 minutes in darkness and washed off in
20
21 distilled water. Cells were then counterstained with Harris haematoxylin and mounted
22
23 with glycergel. A negative control, where no primary antibody was added was included
24
25 in each experimental run. Positive staining was determined by brown staining in the
26
27 cytoplasm. Numbers of cells positive for bcl-2 or cytochrome c were scored blind in both
28
29 direct and bystander areas on each of three replicate slides and expressed as the
30
31 percentage of the total cells counted (approx. 400 - 500 cells were counted in each area).
32
33
34
35
36
37
38
39
40

41 *Quantification of apoptosis*

42
43 Twenty four hours after microbeam irradiation, the cells were washed twice in PBS to
44
45 remove any debris and then fixed in 10% buffered formalin. The mylar was removed and
46
47 adhered to a glass slide as described above. The slides were transported back to Dublin
48
49 immersed in PBS in plastic coplin jars, stored at 4°C and assayed within 48 hours. The
50
51 cells were stained for 15 mins with 1 µg / ml Propidium Iodide (Sigma, Dorset, U.K.) to
52
53 visualise the nuclei. Slides were scored for the presence of apoptotic cells using a Zeiss
54
55
56
57
58
59
60

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Axioplan epifluorescence microscope equipped with BP546/12 nm excitation filter and LP590nm emission filter (Carl Zeiss Ltd, Welwyn Garden City, U.K.). Cells were defined as apoptotic if they displayed evidence of two or more of the following; cell volume shrinkage and pycnotic nucleus (chromatin condensation), nuclear fragmentation and formation of apoptotic bodies (Kerr and Harmon 1991). Numbers of apoptotic cells were scored blind in both direct and bystander areas on each of three replicate slides and expressed as the percentage of the total cells counted (approx. 500 cells were counted in each area).

Statistical analysis

Microscope and detector parameters were standardised to allow quantitative comparisons of the relative fluorescence intensity of the cells between groups. All measurements are presented as mean values \pm S.E. of 3 independent experiments with 3 replicate dishes per measurement. Significance of differences was determined by a student's unpaired t-test and the differences were considered significant if $p \leq 0.05$.

Results

A significant reduction in rhodamine 123 fluorescence, indicating mitochondrial membrane potential depolarisation, was observed in HPV-G cells 6 hours after direct microbeam irradiation (figure 1). Cells that were not directly irradiated but were in the same dish (bystander cells) also showed a similar significant reduction in fluorescence (figure 1).

A significant increase in DCF-DA fluorescence, indicating an increase in ROS, was observed in HPV-G cells 6 hours after direct microbeam irradiation (figure 2). Cells in another area of the same dish (bystander cells) also showed a significant increase in fluorescence (figure 2).

Both the directly irradiated cells and the bystander cells showed increased expression of bcl-2 6.5 hours after microbeam irradiation (table I). Similarly increased levels of cytochrome c were observed 6.5 hours after microbeam irradiation in both the directly irradiated cells and the bystander cells (table I).

HPV-G cells showed increased apoptosis 24 hours after direct microbeam irradiation (figure 3). Cells that were not directly irradiated but were in the same dish (bystander cells) also showed a similar increase in apoptosis levels (figure 3).

For all endpoints measured, there was no significant difference between the effect in the directly irradiated cells and the bystander cells. Similarly there was no significant difference between the effect following irradiation with 1 or 10 protons to the directly exposed cells.

Discussion

This study has shown a significant reduction in mitochondrial membrane potential and a significant increase in ROS in both directly irradiated and bystander HPV-G cells 6 hours after microbeam irradiation. Increased bcl-2 expression and cytochrome c release, after approx 6.5 hours, and increased apoptosis, after 24 hours, were also observed. No significant differences were observed between the effects of different doses.

These results are very similar to those previously reported for HPV-G cells exposed to medium from γ irradiated cells (Lyng *et al*, 2000, 2002, Maguire *et al*, 2005). This finding is important as it shows the mechanisms are similar for medium transfer and for microbeam irradiation and therefore more likely to be universal.

The findings from this study and from previous studies (Lyng *et al*, 2000, 2002, Maguire *et al*, 2005) have shown that early apoptotic events, such as mitochondrial membrane potential depolarisation, induction of ROS, expression of bcl-2 and release of cytochrome c are induced in normal human keratinocytes either exposed to medium from γ irradiated cells or in the vicinity of microbeam irradiated cells. The effects observed appear to be independent of the dose or number of protons delivered to the irradiated cells. A medium borne factor is likely to be involved in both cases. It is unlikely that gap junctional communication is involved in the medium transfer approach and in the present microbeam study the directly irradiated cells and the bystander cells were two distinct populations separated by on average 6 mm.

Most of the studies on bystander effects have used either direct exposure to low fluences of α particles, direct irradiation using microbeam approaches or medium transfer after

1
2
3 low LET exposure. There have been very few reports on the LET dependence of the
4 bystander effect. Hickman *et al* (1994) observed a bystander response, evidenced by
5 increased p53 expression, in rat lung epithelial cells exposed to low fluences of
6 α particles. No increase was seen in cells exposed to similar doses (less than 10 cGy) of
7 X-rays, indicating the existence of a relatively higher damage threshold for sparsely
8 ionizing radiation. Shao *et al* (2002, 2003b) reported an LET dependent induction of
9 micronuclei and cell proliferation in human neoplastic epithelial cells. High LET (100
10 keV/ μ m) carbon-ion irradiation was found to be more efficient at inducing the medium-
11 mediated bystander effect than low LET (13 keV/ μ m) carbon-ion irradiation. Further
12 studies by the same group compared the bystander responses in primary human
13 fibroblasts individually targeted by a high LET heavy particle microbeam of ^{40}Ar (1260
14 keV/ μ m) or ^{20}Ne (380 keV/ μ m) (Shao *et al*, 2003b). An increase in micronuclei was
15 observed independent of the LET and the number of particles delivered to the targeted
16 cells. Previous studies by our group has shown increased apoptosis in human
17 keratinocytes exposed to a medium borne factor from cells irradiated with low LET γ
18 rays (Lyng *et al*, 2000, 2002, Maguire *et al*, 2005). The protons used in the present study
19 (3.2 MeV with an LET of ~ 13 keV/ μ m) are essentially low LET and importantly, the
20 degree of bystander responses observed was similar to that obtained with γ -rays.
21
22 An important observation from these studies is that the level of effect is the same
23 regardless of whether cells were directly exposed or were neighboring non-targeted
24 bystander cells. This agrees with other studies showing that at least after low dose
25 exposure, the bystander response predominates the overall effect (Schettino *et al*, 2003,
26 Seymour and Mothersill 2000).
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

References

Azzam EI, de Toledo SM, Gooding T, Little JB. 1998. Intercellular communication is involved in the bystander regulation of gene expression in human cells exposed to very low fluences of alpha particles. *Radiation Research* 150: 497-504

Azzam EI, de Toledo SM, Little JB. 2001. Direct evidence for the participation of gap junction-mediated intercellular communication in the transmission of damage signals from alpha -particle irradiated to nonirradiated cells. *Proceedings of the National Academy of Sciences U.S.A.* 98: 473-478

Azzam EI, de Toledo SM, Spitz DR, Little JB. 2002. Oxidative metabolism modulates signal transduction and micronucleus formation in bystander cells from alpha-particle-irradiated normal human fibroblast cultures. *Cancer Research* 62: 5437-5442

Azzam EI, de Toledo SM, Little JB. 2003. Oxidative metabolism, gap junctions and the ionizing radiation-induced bystander effect. *Oncogene* 22: 7050-7057

Belyakov OV, Malcolmson AM, Folkard M, Prise KM and Michael BD. 2001. Direct evidence for a bystander effect of ionizing radiation in primary human fibroblasts. *British Journal of Cancer* 84: 674-679

1
2
3 Belyakov OV, Folkard M, Mothersill C, Prise KM, Michael BD. 2003. A proliferation-
4
5 dependent bystander effect in primary porcine and human urothelial explants in response
6
7 to targeted irradiation. *British Journal of Cancer* 88: 767-774
8
9

10
11
12 Deshpande A, Goodwin EH, Bailey SM, Marrone BL and Lehnert BE. 1997. Alpha-
13
14 particle-induced sister chromatid exchange in normal human lung fibroblasts: evidence
15
16 for an extranuclear target. *Radiation Research* 145: 260-267
17
18
19

20
21
22 Folkard M, Vojnovic B, Hollis KJ, Bowey AG, Watts SJ, Schettino G, Prise KM and
23
24 Michael BD. 1997. *International Journal of Radiation Biology* 72: 387-395
25
26
27

28
29 Hickman AW, Jaramillo RJ, Lechner JF, Johnson NF. 1994. Alpha-particle-induced p53
30
31 protein expression in a rat lung epithelial cell strain. *Cancer Research* 54: 5797-5800
32
33
34

35
36 Iyer R, Lehnert BE and Swensson R. 2000. Factors underlying the cell growth-related
37
38 bystander responses to alpha particles. *Cancer Research* 60: 1290-1298
39
40
41

42
43 Kerr JFR and Harmon BV. 1991. Definition and incidence of apoptosis. An historical
44
45 perspective. In: Tomei LD and Cope FO, editors. *Apoptosis. The Molecular Basis of*
46
47 *Cell Death*, New York: Cold Springs Harbour Laboratory, pp 5 - 9
48
49
50

1
2
3 Lehnert BE, Goodwin EH, Desppande A. 1997. Extracellular factor(s) following
4 exposure to alpha-particles can cause sister chromatid exchanges in normal human cells.
5
6

7
8 Cancer Research 57: 2164-2171
9

10
11
12 Little JB. 2003. Genomic instability and bystander effects: a historical perspective.
13

14
15 Oncogene 22: 6978-6987
16
17

18
19
20 Lorimore SA, Coates PJ, Wright EG. 2003. Radiation-induced genomic instability and
21 bystander effects: inter-related nontargeted effects of exposure to ionizing radiation.
22
23

24
25 Oncogene 22: 7058-7069
26
27

28
29 Lyng FM, Seymour CB, Mothersill C. 2000. Production of a signal by irradiated cells
30 which leads to a response in unirradiated cells characteristic of initiation of apoptosis.
31
32

33
34 British Journal of Cancer 83: 1223-1230.
35
36

37
38
39 Lyng FM, Seymour CB, Mothersill C. 2002. Initiation of apoptosis in cells exposed to
40 medium from the progeny of irradiated cells: a possible mechanism for bystander-
41
42

43
44 induced genomic instability? Radiation Research 157: 365-370
45
46

47
48
49 Lyng FM, Maguire P, McClean B, Seymour C, Mothersill C. 2006. The involvement of
50 calcium and MAP kinase signalling pathways in the production of radiation induced
51
52

53
54 bystander effects, Radiation Research (in press)
55
56
57
58
59
60

1
2
3 Maguire P, Mothersill C, Seymour C and Lyng FM. 2005. Medium from irradiated cells
4 induces dose dependent mitochondrial changes and bcl-2 responses in unirradiated
5 human keratinocytes. *Radiation Research* 163: 384-390
6
7
8
9

10
11
12 **Matsumoto H, Hayashi S, Hatashita M, Shioura H, Ohtsubo T, Kitai R, Ohnishi T,**
13 **Yukawa O, Furusawa Y, Kano E. 2000. Induction of radioresistance to accelerated**
14 **carbon-ion beams in recipient cells by nitric oxide excreted from irradiated donor cells of**
15 **human glioblastoma. *International Journal of Radiation Biology* 76: 1649-1657**
16
17
18
19
20
21

22
23
24 Morgan WF. 2003. Non-targeted and delayed effects of exposure to ionizing radiation: I.
25 Radiation-induced genomic instability and bystander effects in vitro. *Radiation Research*
26 159: 567-580
27
28
29
30
31

32
33
34 Mothersill C, Seymour C. 1997. Medium from irradiated human epithelial cells but not
35 human fibroblasts reduces the clonogenic survival of unirradiated cells. *International*
36 *Journal of Radiation Biology* 71 421-427
37
38
39
40

41
42
43 Mothersill C, Seymour CB. 1998. Cell-cell contact during gamma irradiation is not
44 required to induce a bystander effect in normal human keratinocytes: evidence for release
45 during irradiation of a signal controlling survival into the medium. *Radiation Research*
46 149 256-262
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 Mothersill C and Seymour CB. 2004. Radiation-induced bystander effects--implications
4 for cancer. *Nature Reviews Cancer* 4: 158-164
5
6
7

8
9
10 Nagasawa H and Little JB. 1992. Induction of sister chromatid exchanges by extremely
11 low doses of alpha particles. *Cancer Research* 52: 6394 – 6396
12
13

14
15
16
17 Narayanan PK, Goodwin EH, Lehnert BE. 1997. α particles initiate biological
18 production of superoxide anions and hydrogen peroxide in human cells. *Cancer Research*
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

57: 2963-3971

Pirisi L, Creek KE, Doniger J, DiPaolo J. 1988. Continuous cell lines with altered growth
and differentiation properties originate after transfection of human keratinocytes with
human papillomavirus type 16 DNA. *Carcinogenesis*. 9: 1573-1579

Prise KM, Belyakov OV, Folkard M, and Michael B. 1998. Studies of bystander effects
in human fibroblasts using a charged particle microbeam. *International Journal of
Radiation Biology* 74: 793-798

Sawant SG, Randers-Pehrson G, Geard CR, Brenner DJ, Hall EJ. 2001. The bystander
effect in radiation oncogenesis: I. Transformation in C3H 10T1/2 cells in vitro can be
initiated in the unirradiated neighbors of irradiated cells. *Radiation Research* 155: 397-
401

1
2
3 Schettino G, Folkard M, Prise KM, Vojnovic B, Held KD and Michael BD. 2003. Low-
4 dose studies of bystander cell killing with targeted soft X rays, *Radiation Research* 160:
5
6 505-511
7
8

9
10
11
12 Seymour CB, Mothersill C. 2000. Relative contribution of bystander and targeted cell
13 killing to the low-dose region of the radiation dose-response curve. *Radiation Research*
14
15 153: 508-511.
16
17
18

19
20
21
22 Shao C, Aoki M, Furusawa Y. 2001. Medium-mediated bystander effects on HSG cells
23 co-cultivated with cells irradiated by X-rays or a 290 MeV/u carbon beam. *Journal of*
24
25 *Radiation Research (Tokyo)* 42: 305-316
26
27
28

29
30
31 Shao C, Furusawa Y, Aoki M, Matsumoto H, Ando K. 2002. Nitric oxide-mediated
32 bystander effect induced by heavy-ions in human salivary gland tumour cells.
33
34 *International Journal of Radiation Biology* 78: 837-844
35
36
37

38
39
40
41 Shao C, Stewart V, Folkard M, Michael BD and Prise KM. 2003a. Nitric oxide-mediated
42 signaling in the bystander response of individually targeted glioma cells. *Cancer*
43
44 *Research* 63 8437-8442
45
46
47

48
49
50 Shao C, Furusawa Y, Kobayashi Y, Funayama T, Wada S. 2003b. Bystander effect
51 induced by high-LET particles in confluent human fibroblasts: a mechanistic study.
52
53 *FASEB Journal* 17: 1422-1427
54
55
56

1
2
3
4
5
6 Shao C, Aoki M, and Furusawa Y. 2004. Bystander effect in lymphoma cells vicinal to
7
8 irradiated neoplastic epithelial cells: Nitric oxide is involved. Journal of Radiation
9
10 Research 45: 97-103
11

12
13
14
15 Suzuki M, Zhou H, Geard CR, Hei TK. 2004 Effect of medium on chromatin damage in
16
17 bystander mammalian cells. Radiation Research 162: 264- 269
18

19
20
21
22 Zhou H, Randers-Pehrson G, Waldren CA, Vannais D, Hall EJ and Hei TK. 2000.
23
24 Induction of a bystander mutagenic effect of alpha particles in mammalian cells.
25
26 Proceedings of the National Academy of Sciences U.S.A. 97: 2099-2104
27

28
29
30
31 Zhou H, Suzuki M, Randers-Pehrson G, Vannais D, Chen G, Trosko JE, Waldren CA,
32
33 Hei TK. 2001. Radiation risk to low fluences of alpha particles may be greater than we
34
35 thought. Proceedings of the National Academy of Sciences U.S.A. 98:14410-14415
36
37

38 39 40 41 **Acknowledgements**

42
43 The authors acknowledge support from Science Foundation Ireland, the Royal Irish
44
45 Academy, the Royal Society, the Gray Cancer Institute and Cancer Research UK.
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Figure Legends


Figure 1 % Fluorescence from rhodamine 123 in both directly irradiated and bystander HPV-G cells 6 hours after microbeam irradiation. A decrease in fluorescence levels is indicative of a decrease in mitochondrial membrane potential. * $p < 0.001$

Figure 2 % Fluorescence from 2,7 dichlorofluorescein diacetate in both directly irradiated and bystander HPV-G cells 6 hours after microbeam irradiation. An increase in fluorescence levels is indicative of an increase in reactive oxygen species. * $p < 0.001$

Figure 3 % Apoptotic cells in both directly irradiated and bystander HPV-G cells 24 hours after microbeam irradiation. * $p < 0.05$, ** $p < 0.01$

Table I % HPV-G cells positive for bcl-2 and cytochrome c 6.5 hours following microbeam irradiation. * p<0.05, ** p<0.01, *** p<0.005

Dose	% bcl-2 positive cells		% cytochrome c positive cells	
	Direct	Bystander	Direct	Bystander
Control	2.17 ± 0.33	1.67 ± 0.60	1.83 ± 0.44	2.50 ± 0.73
1 proton	7.17 ± 0.73 **	7.00 ± 0.29 *	13.83 ± 0.60 ***	14.17 ± 0.88 **
10 protons	9.00 ± 0.59**	7.67 ± 1.01*	15.67 ± 0.73***	14.33 ± 0.19 **



1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49

