

2008-07-01

Green Tea Extract as a Natural Antioxidant to Extend the Shelf-life of Fresh-cut Lettuce

Catherine Barry-Ryan

Technological University Dublin, catherine.barryryan@tudublin.ie

Ana Belen Martin-Diana

Technological University Dublin, anabelen.martindiana@tudublin.ie

Daniel Rico

Technological University Dublin, daniel.rico@tudublin.ie

Follow this and additional works at: <https://arrow.tudublin.ie/schfsehart>

Recommended Citation

Barry-Ryan,C., Martin-Diana,A.,Rico,D. (2008) Green tea extract as a natural antioxidant to extend the shelf-life of fresh-cut lettuce. *Innovative Food Science and Emerging Technologies*, 9 (2008) 593–603.
doi:10.1016/j.ifset.2008.04.001

This Article is brought to you for free and open access by the School of Food Science and Environmental Health at ARROW@TU Dublin. It has been accepted for inclusion in Articles by an authorized administrator of ARROW@TU Dublin. For more information, please contact arrow.admin@tudublin.ie, aisling.coyne@tudublin.ie, vera.kilshaw@tudublin.ie.

Green tea extract as a natural antioxidant to extend the shelf-life of fresh-cut lettuce

Innovative Food Science and Emerging Technologies 9 (2008) 593–603

A. B. Martín-Diana, D. Rico, Catherine Barry-Ryan

School of Food Science and Environmental Health, Dublin Institute of Technology (DIT), Cathal Brugha Street, Dublin 1, Ireland

Keywords: Green tea, *Camellia sinensis* L., Fresh-cut lettuce, Polyphenols, Browning

ABSTRACT

Green tea extract (GT) was evaluated as a preservative treatment for fresh-cut lettuce. Different quality markers, e.g. respiration, browning, ascorbic acid and carotenoid content were evaluated. GT concentration (0.25, 0.5 and 1 g 100 mL⁻¹) and temperature (20 °C and 50 °C) were tested. Optimal GT treatment (0.25 g 100 mL⁻¹ at 20 °C) was compared with chlorine (120 ppm at 20 °C). High GT concentrations (0.5 g 100 mL⁻¹ and 1.0 g 100 mL⁻¹) maintained better prevent ascorbic acid and carotenoid loss than 0.25 g 100 mL⁻¹ GT and chlorine. GT increased browning of samples, probably due to the content of polyphenols of the treatment; the use of heat-shock reduced this negative effect. GT and heat-shock combined also showed negative effects, reducing the antioxidant content (ascorbic acid and carotenoids). No significant differences were observed between chlorine and optimal GT (0.25 g 100 mL⁻¹ at 20 °C) in browning appearance and sensory properties. GT better kept the antioxidant activity of the samples than chlorine.

Industrial relevance: An alternative treatment for minimally processed Iceberg lettuce is tested, based on its antioxidant capacity. Minimally processed industry is constantly looking for new treatments to avoid the use of chlorine which is a standard at the moment.

1. Introduction

Green tea (*Camellia sinensis* L.) is a popular beverage and it has been reported to exert beneficial effects on several life-style related diseases: chemopreventive, anticarcinogenic, antiatherogenic, antioxidant and antimicrobial activities (Si et al., 2006; McKay & Blumberg, 2002; Rietveld & Wiseman, 2003; Cooper, Morre & Morre, 2005). The ability to inhibit food-borne pathogens such as *Escherichia coli*, *Salmonella typhimurium*, *Listeria monocytogenes*, *Staphylococcus aureus*, and *Campylobacter jejuni* has been reported for different varieties of tea or tea extracts, including Oolong, Jasmine and Black (Si et al., 2006; Hamilton-Miller, 1995; Hamilton-Miller & Shah, 2005; Yeo, Ahn, Lee, Lee, Park & Kim, 1995).

However, results showing antimicrobial activity of tea extract components against bacterial pathogens have been inconsistent due to the variability of the methods used (e.g. different varieties of tea, different processing and extraction procedures) and the mechanisms of action described in the literature remain controversial (Si et al., 2006).

Green tea is an excellent source of polyphenols, which are natural antioxidants that can be used as alternatives to synthetic antioxidants as they are typically less harmful than synthetic ones and appears to have an equivalent effect upon the inhibition of oxidation (Cao, Sofic & Prior, 1996). These antioxidants, which inhibit the oxidation of organic molecules, are very important, not only for living systems and their defence against oxidative stress but also for food preservation

(Masuda et al., 2003). Phenolic antioxidants interrupt the propagation of the free radical autoxidation chain by contributing a hydrogen atom from a phenolic hydroxyl group, with the formation of a relatively stable free radical that does not initiate or propagate further oxidation processes (Kaur & Kapoor, 2001).

Fresh-cut vegetable industry (~ 10% p.a.) has undergone an important growth due to the increasing demand for fresh, healthy and convenient foods. A substantial portion of vitamins and minerals in the diet comes from fruit and vegetable consumption (Klein, 1987). There is mounting evidence to support the alleviation of many degenerative diseases including cardiovascular disease, cancer and ageing by the consumption of fruit and vegetables (Kaur & Kapoor, 2001). These beneficial health effects of fruit and vegetables have been attributed to the presence of antioxidants that act as receptors of free radicals. Ascorbic acid and β -carotene are the antioxidants present in the greatest quantities in fruit and vegetables and usually are destroyed after processing or decontamination procedures.

Consumers have also become more critical to the use of artificial additives in order to preserve food or enhance characteristics such as colour, flavour and nutritional value (Bruhn, 2000). Ohlsson (1994) suggests that minimal processing techniques have emerged to meet the challenge of replacing traditional methods of preservation while retaining nutritional and sensory quality. RTU vegetables typically involve peeling, slicing, dicing or shredding prior to packaging and storage (Barry-Ryan & O'Beirne, 1999). The International Fresh-cut Produce Association (IFPA) defines fresh-cut products as fruit or vegetables that have been trimmed and/or peeled and/or cut into




Fig. 1. Experimental design for the testing of Green tea extracts treatments on fresh-cut lettuce.

100% usable product that is bagged or pre-packaged to offer consumers high nutrition, convenience, and flavour while still maintaining its freshness (Lamikanra, 2002).

It is well-known that processing of vegetables promotes a faster physiological deterioration, biochemical changes and microbial degradation of the product even when only slight processing operations can be used (O'Beirne & Francis, 2003), which may result in degradation of the colour, texture and flavour (Kabir, 1994; Varoquaux & Wiley, 1994). While conventional food-processing methods extend the shelf-life of fruit and vegetables, the minimal processing to which fresh-cut fruit and vegetables are subjected renders products highly perishable, requiring chilled storage to ensure a reasonable shelf-life (Garcia & Barret, 2002).

The vast majority of fresh minimally processed produce manufacturers use chlorine in washing and decontamination procedures (Seymour, 1999). There is controversy about the formation of carcinogenic chlorinated compounds in water (chloramines and trihalo-methanes), calling into question the use of chlorine (Wei et al., 1995).

The current concern associated with chlorine outlines the need to find new alternatives for preservation of minimally processed vegetables. Traditional and novel technologies for maintaining quality while inhibiting undesired microbial growth are being investigated, driven by an increasing demand from the industry and consumers. These include use of temperature blanching/steamer (Abreu, Beirão-da-Costa, Gonçalves, Beirão-da-Costa & Moldão-Martins, 2003; Martín-Diana et al., 2007a; Rico et al., 2008a,b); short thermal shocks (Loaiza-Velarde, Mangrich, Campos-Vargas, & Saltveit, 2003; Martín-Diana et al., 2005a;

Koukounaras, Diamantidis & Sfakiotakis, 2008); organic acids (Rosen & Kader, 1989; Agar, Massantini, Hess-Pierce, & Kader, 1999; Luna-Guzmán & Barrett, 2000; Martín-Diana et al., 2005b; Martín-Diana et al., 2007b), irradiation (Allende & Artes, 2003a; Han et al., 2004), modified atmosphere packaging (Kato-Noguchi & Watada, 1997; Allende & Artes, 2003b), ozone (García, Mount & Davidson, 2003; Zhang, Zhaoxin, Zhifang & Xiang, 2005; Rico, Martín-Diana, Frías, Henehan & Barry-Ryan, 2006), electrolysed water (Izumi, 1999; Koseki, Yoshida, Kamitani, Isobe & Itoh, 2004; Park, Hung, Doyle, Ezeike & Kim, 2001; Rico et al., 2008a,b), electric pulses (Giner et al., 2000), edible coating (Molnar-Perl & Friedman, 1990; Guilbert, Gontard, & Gorris, 1996; Son, Moon, & Lee, 2001) among others.

During the last years a special renewed interest in the use of natural products as decontaminants has been noticed. Research and commercial applications have shown that natural antimicrobials could replace traditional sanitizing agents (Graham, 1997; Cherry, 1999; Martín-Diana, Rico et al., 2006). Green tea has been used to extend the shelf-life of dry-fermented sausage (Bozkurt, 2006), enrich candy jellies (Gramza-Michalowska & Regula, 2007) and cooked pork patties (Nissen, Byrne, Bertelsen & Skibsted, 2004). However, only a few studies can be found on the use of Green tea in fresh-cut fruits or vegetables.

Due to the lack of information regarding the use of Green tea in minimally processed vegetables, the objective of this research was to evaluate its ability to extend the shelf-life of fresh-cut lettuce. Quality markers, mainly indicators of respiration, browning, sensory and nutritional parameters were used to optimise the Green tea extract treatment conditions (concentration and temperature) and to compare its effect on fresh-cut lettuce quality with chlorine (standard treatment).

2. Materials and methods

2.1. Sampling and experimental design

The experiments were carried out between December 2004 and June 2005. All procedures were performed in a food-processing room at 20 °C. Each batch consisted of ~ 18–25 kg (~ 55–65 heads of lettuce of ~ 450 g each). For colour and texture analyses all pieces in the assigned bag (20 to 25 approx.) were analysed individually to minimise error due to product variability. For sensory analysis, samples from the whole storage bag were analysed. For destructive analysis (enzymes, ascorbic acid, carotenoids and pH) samples were taken from a pool of all the pieces in a storage bag.

The experimental design was divided in three parts (Fig. 1): two-step optimisation of Green tea extract (GT) treatment conditions to extend the quality and nutritional value of fresh-cut lettuce (Parts I and II) and comparison of GT and chlorine treatments (Part III).

Preliminary sensory studies using GT concentrations from 0 to 5 g 100 mL⁻¹ vs. 120 ppm chlorine as control were carried out in fresh-cut lettuce. Sensory results showed that it was necessary to use at least concentrations of 0.25 g 100 mL⁻¹ and a maximum of 1 g 100 mL⁻¹ to obtain samples considered as acceptable during the shelf-life (10 days).

The optimisation of GT concentration (Part I) was done using concentrations (0.25, 0.5 and 1.0 g 100 mL⁻¹). Part II consisted of a

comparative analysis using GT treatments at a concentration obtained from the optimisation step (Part I) at two temperatures: 20 and 50 °C. The temperatures were selected to analyse the combined effect of heat-shock and GT. The two temperatures were selected based on previous research (Martin-Diana et al., 2005a,b,c). The last part of the experiment (Part III) was carried out in order to compare the optimal GT conditions obtained in the two previous steps (Part I and II) with an industrial standard treatment.


2.2. Raw material

Iceberg lettuce (*Lactuca sativa* sp.) was grown in Ireland and purchased from a local supplier. The lettuce was brought to the laboratory within 24 h after harvest and stored in cardboard boxes, in bulk and unwrapped at 4 °C until processing.

2.3. Processing (washing and packaging procedure)

Lettuce was prepared according to safety statements and recommendations (Beuchat & Ryu, 1997; Beuchat, 1998). The two outer leaves were removed and the heads cored with a stainless steel knife. The lettuce was cut in halves and each half was further cut into four pieces.

Washing treatments consisted of immersion of the fresh-cut lettuce in the treatment solution. Dried and ground Green tea (*C. sinensis* L.) leaves



*Points designated on any curve by the same letter are not significantly different ($p > 0.05$)

Fig. 2. Changes in gas composition (carbon dioxide (A,C,E) and oxygen (B,D,F)) of package headspace of fresh-cut lettuce treated with GT and stored at 4 °C over 10 days. Part I: GT at different concentrations (0.25, 0.5 and 1 g 100 mL⁻¹ at 20 °C). Part II: 0.25 g 100 mL⁻¹ GT at 20 °C and 50 °C. Part III: 0.25 g 100 mL⁻¹ GT at 20 °C and 120 ppm chlorine at 20 °C.

were brewed for 5 min in distilled water at 90 °C at different concentrations (0.25 g 100 mL⁻¹, 0.5 g 100 mL⁻¹ and 1 g 100 mL⁻¹ with a content in total phenolics respectively of 350, 700 and 1260 mg of gallic acid per litre of infusion). The obtained extracts were filtrated two times through a Whatman No. 1 filter paper. Chlorinated water was prepared by adding sodium hypochlorite (≥ 120 ppm available chlorine) to distilled water to obtain a 120 ppm free chlorine solution (pH ~ 8). Free chlorine concentration in the washing solution was determined with the DPD method, using a DREL/2000 spectrometer with appropriate AccuVac® pillows (Hach Company, Loveland, Colorado, USA).

For all treatments the solutions were prepared using distilled water stored at selected temperatures according to the experimental design. The cut produce was placed in a basket and immersed with agitation in the solution (~ 100 g vegetable product per litre) for 1 min and subsequently dried for 5 min using an automatic salad spinner.

To minimise product heterogeneity, processed vegetables were pooled, mixed and subsequently packaged in bags (200 \times 320 mm) of oriented polypropylene (Amcor Flexibles Europe-Brighthouse, United Kingdom). Film permeabilities to oxygen and carbon dioxide were $\sim 12,000$ and $\sim 13,000$ (mL m⁻² day⁻¹ atm⁻¹ at 5 °C) respectively. Each package contained ~ 100 g of the product. The packages were chilled in a blast freezer set at 0 °C for 2 min before being sealed to avoid water condensation during storage (10 days at 4 °C).

2.4. Shelf-life analysis

The objective was to study the effect of GT on the quality of fresh-cut lettuce. Respiration, pH, browning appearance (browning-related enzymes, colour and potential browning) and nutritional changes (ascorbic acid and carotenoids) were evaluated as indicators of shelf-life. Shelf-life is defined as the length of time at which the vegetable can maintain the appearance, safety and antioxidant content that appeals to the consumer (Delaquis, Stewart, Toivonen & Moyls, 1999), therefore markers were monitored throughout the storage (10 days).

2.4.1. Headspace analysis

A Gaspacer analyser (Systech Instruments, UK) was used to measure levels of CO₂ and O₂ during storage. Gas extractions were performed with a hypodermic needle, inserted through an adhesive septum previously fixed to the bags, at a flow rate of 30 mL/min for 10 s. The accuracy is in percentage of readings: $\pm 0.5\%$ below 10% and $\pm 0.1\%$ above 10% for O₂; $\pm 0.1\%$ below 1%, $\pm 0.2\%$ between 1% and 10% and $\pm 2\%$ above 10% for CO₂. Three bags per treatment and batch were monitored throughout the storage.

2.4.2. pH. Ten grams sample of lettuce tissue was blended for 2 min in 20 mL of deionised water. The pH of the slurry was measured at room temperature using an Orion research pH-meter.

2.4.3. Browning

2.4.3.1. Colour measurement. Colour was quantified using a colorimeter (Minolta, Model CM-3600d, UK). A lettuce piece was placed directly on the colorimeter sensor (3.5 cm of diameter) and measured; 20–30 measurements were taken per treatment and day. The instrument was calibrated with a white tile standard (L* = 93.97, a* = -0.88 and b* = 1.21) and a green tile standard (L* = 56.23, a* = -21.85, b* = 8.31) under D₆₅ luminosity conditions. The a* parameter measures the degree of red (+a) or green (-a) colour and the b* parameter measures the degree of yellow (+b) or blue (-b) colour.

2.4.3.2. Potential browning. Potential browning was measured according to the methodology of Viña and Chaves (2006). Ten grams of tissue frozen and crushed was treated with ethanol for 60 min and then centrifuged at 6000 rpm at 10 °C for 10 min, retaining the supernatants. After a further amount of ethanol was added to bring the final volume to 25 mL. Absorbance at 320 nm was measured on aliquots of these extracts. The results were expressed as absorbance units (AU) g⁻¹ fresh tissue.

2.4.3.3. Browning-related enzymes (peroxidase (POD) E.C.1.11.1.7 and polyphenol oxidase (PPO) E.C.1.10.3.1.). Both enzymes were assayed in homogenates that were prepared as follows: 10 g of lettuce was placed in a polytron homogeniser (Polytron model PT 3000) in a 1:2 (w:v) ratio with 0.5 M phosphate buffer pH 6.5 containing 50 g L⁻¹ polyvinylpyrrolidone. Homogenisation was carried out two times at 4 °C, and 5500 rpm, for 1 min each time with a break of 3 min between homogenisations in order to avoid excessive heating of the sample. The homogenate was then centrifuged at 12,720 \times g for 30 min at 4 °C. It was filtered through one layer of crepe bandage. The resulting crude extract was used without further purification. All the extracts were stored at 4 °C in the dark and used immediately.

POD activity was assayed spectrophotometrically. The reaction mixture contained 0.2 mL of extract and 2.7 mL of 0.05 M phosphate buffer pH 6.5 containing 100 μ L of hydrogen peroxide (1 mL/L v/v) as an oxidant and 200 μ L of p-phenylenediamine as hydrogen donor. The oxidation of p-phenylenediamine was monitored at 485 nm and 25 °C. An enzyme activity unit was defined as an increment of 0.1 in absorbance per minute. PPO activity was assayed spectrophotometrically by a modified method based on Galeazzi, Sgarbieri and Constantinides (1981) and Tan and Harris (1995). The reaction mixture contained 0.1 mL crude extract and 2.9 mL substrate solution (0.020 mol/L catechol in 0.05 mol/L phosphate buffer, pH 6.5). The rate of catechol oxidation was followed at 400 nm for 2 min at 25 °C. An enzyme activity unit was defined as an increase of 0.1 in absorbance per minute. Three independent trials were carried out. All the enzymatic measurements were carried out in duplicate.

2.4.4. Sensory analysis. Analytical-descriptive tests were used to evaluate the sensory quality attributes of fresh-cut lettuce prepared by different washing treatments. The panel, which consisted of 10 judges aged 20–30 years with sensory evaluation experience, were trained in discriminate evaluation of fresh-cut tissue. Panelists were required to score changes in total browning, photosynthetic browning, vascular browning and fresh appearance. Before the start of the sensory experiments, panel members were familiarised with the product and scoring methods. This consisted of demonstration exercises involving examination of packs at different levels of deterioration and agreeing appropriate scores. When the panel members had become familiar with the test facilities and scoring regime, they were invited to score samples. This procedure was repeated several times until a level of consistency in scoring was obtained. The same packages were scored during the entire test for sensory analysis (10 days), due to the high variability of the product. During this stage, the samples were presented to the panel to evaluate and measure the reproducibility of the judge's answers and their capability in discriminating among samples. During the analyses, samples were presented in randomized order to minimise possible carry-over effects.

Fresh-cut lettuce was evaluated for appearance using a ten point numerical rating scale, where: (I) General Browning, where 1 = nothing, 5 = moderate and 10 = a lot (severe browning); (II) Photosynthetic Browning, where 1 = nothing, 5 = moderate and 10 = a lot (severe browning); (III) Vascular Browning, where 1 = nothing, 5 = moderate and 10 = a lot (severe browning); (IV) Fresh appearance, where 1 = poor/no fresh appearance, 5 = moderate and 10 = very good/fresh appearance.

The results of the sensory analysis were reported as means of three separate trials. Data analysis was collected using Compusense® Five software (Release 4.4, Ontario, Canada).

2.4.5. Nutritional markers

2.4.5.1. Ascorbic acid. Ascorbic acid determination was carried out according to the 2,6-dichlorolindophenol method recommended by AOAC (1995) for the analysis of Vitamin C in fruits and juices. grams of fresh-cut was homogenised with 50 mL of acetic-methaphosphoric acid at low speed (in order to avoid foaming) for




Fig. 3. Changes in pH of fresh-cut lettuce stored at 4 °C for 10 days. Part I: GT at different concentrations (0.25, 0.5 and 1 m 100 mL⁻¹ at 20 °C). Part II: 0.25 g 100 mL⁻¹ GT at 20 °C and 50 °C. Part III: 0.25 g 100 mL⁻¹ GTE at 20 °C and 120 ppm chlorine at 20 °C.

3 min at 4 °C in darkness. –The homogenate was filtered and aceticmethaphosphoric added up to 100 mL. Aliquots of 10 mL were titrated rapidly in triplicate, using aceticmethaphosphoric acid as a blank. The results were expressed as mg ascorbic acid in g of sample.

$$\text{mg ascorbic acid/g sample} = (X - B) * (F/E) * (V/Y)$$

where X is mL 2,6-dichlorolindophenol (DCPIP) used to obtain a permanent pink colour which persists more than 5 s, B is the DCPIP volume used for the control (blank), F* is the DCPIP factor (F* = 2 mg ascorbic acid (standard) mL⁻¹ DCPIP), E is g of sample (20 g), V is the final volume prepared (100 mL) and Y is mL titrated (10 mL).

2.4.5.2. Total carotenoids. Total carotenoids were extracted in the dark by homogenising lettuce tissue (5 g) with 30 mL of an acetone/ethanol (50:50 v:v) solution, containing 200 mg L⁻¹ of butylated hydroxytoluene (BHT). The homogenate was filtered under suction in a buchner funnel and washed with acetone/ethanol solvent until

colourless. Filtrate was adjusted to 100 mL volume with acetone/ethanol. An aliquot was placed in a 1 cm cuvette and its absorbance was measured at 470 nm. Total carotenoids (mg/g of sample) were calculated as described by Gross (1991).

2.5. Statistical analysis

Analysis of ANOVA (Multifactor and one-way) was used to find differences between treatments, storage and interaction of both factors for each one of the variables studied. Means were compared by significant difference (LSD) test, at a significance level p = 0.05 using the Statgraphics software (version 2.1; Statistical Graphics Co., Rockville, USA). Three independent trials were carried out.




Fig. 4. Changes in luminosity (CIE L*a*b* parameters) in fresh-cut lettuce stored at 4 °C for 10 days. Part I: GT at different concentrations (0.25, 0.5 and 1 g 100 mL⁻¹ at 20 °C). Part II: 0.25 g 100 mL⁻¹ GT at 20 °C and 50 °C. Part III: 0.25 g 100 mL⁻¹ GT at 20 °C and 120 ppm chlorine at 20 °C.

3. Results and discussion

3.1. Headspace

The package headspace was monitored over 10 days of storage. Carbon dioxide and oxygen content, starting at atmospheric values, significantly ($p < 0.05$) changed regardless of the treatment. GT concentration (Fig. 2A,B) affected the respiration rate of the samples. Significantly ($p < 0.05$) lower oxygen and higher carbon dioxide levels were observed in samples treated with GT concentrations (0.5 and 1.0 g L⁻¹) compared to those treated with the lowest concentration (0.25 g L⁻¹). Higher GT concentrations produced an increase in the




Fig. 5. Changes in a* (CIE L*a*b* parameters) in fresh-cut lettuce stored at 4 °C for 10 days. Part I: GT at different concentrations (0.25, 0.5 and 1 g 100 mL⁻¹ at 20 °C). Part II: 0.25 g 100 mL⁻¹ GT at 20 °C and 50 °C. Part III: 0.25 g 100 mL⁻¹ GT at 20 °C and 120 ppm chlorine at 20 °C.




Fig. 6. Changes in potential browning (AU g⁻¹) in fresh-cut lettuce stored at 4 °C for 10 days. Part I: GT at different concentrations (0.25, 0.5 and 1 g 100 mL⁻¹ at 20 °C). Part II: 0.25 g 100 mL⁻¹ GT at 20 °C and 50 °C. Part III: 0.25 g 100 mL⁻¹ GT at 20 °C and 120 ppm chlorine at 20 °C.

respiration rate, maybe related to a higher stress caused by the more concentrated treatments (Surjadinata & Cisneros-Zevallos, 2003). As indicated by Saltveit (2001) for fresh produce subjected to heat-shock for less than 2 min, treatment temperature (Fig. 2C,D) did not show any effect on the oxygen and carbon dioxide concentrations, results in agreement with previous work of the authors (Martin-Diana et al., 2005b). Headspace composition was significantly ($p < 0.05$) different in samples treated with the optimised GT treatment (0.25 g 100 mL⁻¹, 20 °C) and those treated with chlorine (Fig. 2E,F). These indicated that the GT treatment increased respiration of the produce in comparison with chlorine, probably indicating a higher stress caused by the GT treatment.

In all cases, the levels of oxygen were above 6%, avoiding hypoxic atmosphere conditions which favour the fermentative processes that cause the formation of acetaldehyde and the appearance of off-flavours compounds (Kays, 1991; Beaudry, 1999). In general, the highest respiration rates were observed in samples treated with the intermediate and high concentrations (0.5 and 1.0 g 100 mL⁻¹) GT, indicating a possible negative effect of increased stress with the use of an excessively high concentration of Green tea.

3.2. pH

Changes in pH were monitored in all the samples through the 10-day storage (Fig. 3, Part I, Part II and Part III). A significant increase in pH was observed for all the samples, with a peak at day 7 followed by a decrease at day 10. Different authors (King & Bolin, 1989; Jacxsens,




Fig. 7. Changes in polyphenol oxidase enzymatic activity (PPO, units g⁻¹) in fresh-cut lettuce stored at 4 °C for 10 days. Part I: GT at different concentrations (0.25, 05 and 1 g 100 mL⁻¹ at 20 °C). Part II: 0.25 g 100 mL⁻¹ GT at 20 °C and 50 °C. Part III: 0.25 g 100 mL⁻¹ GT at 20 °C and 120 ppm chlorine at 20 °C.




Fig. 8. Changes in peroxidase enzymatic activity (POD, units g⁻¹) in fresh-cut lettuce stored at 4 °C for 10 days. Part I: GT at different concentrations (0.25, 05 and 1 g 100 mL⁻¹ at 20 °C). Part II: 0.25 g 100 mL⁻¹ GT at 20 °C and 50 °C. Part III: 0.25 g 100 mL⁻¹ GT at 20 °C and 120 ppm chlorine at 20 °C.

Devlieghere, Ragaert, Vanneste & Debevere, 2003; Gomez-Lopez, Devlieghere, Bonduelle & Debevere, 2005) have explained this pH increase with the growth of gram-negative microorganisms, which play an important role in the spoilage of fresh-cut lettuce. Lower GT concentrations kept significantly (p < 0.05) lower pH values during the storage than higher concentrations (Fig. 3A). The lower pH values at day 10 could respond to an increase in lactic acid bacteria. The samples treated with GT and heat-shock maintained significantly lower produce pH, with no peak observed at day 7 (Fig. 3B). This could be due to the inhibitory effect of temperature on bacterial growth. Delaquis et al. (1999) described a reduction in bacterial counts in lettuce treated with warm water. Similar behaviour was observed in samples treated with chlorine compared with those treated with GT (Fig. 3C).

Table 1
Sensory evaluation of fresh-cut lettuce using a rating score (1–10) in fresh-cut lettuce stored at 4 °C for 10 days

Sensory analysis						
Part I	Storage (days)	Treatment	General Browning	Photosynthetic Browning	Vascular Browning	Fresh appearance
ANOVA treatments			GT 0.25/20 °C A	GT 0.25/20 °C A	GT 0.25/20 °C A	GT 0.25/20 °C A
			GT 0.5/20 °C B	GT 0.5/20 °C B	GT 0.5/20 °C B	GT 0.5/20 °C B
			GT 1.0/20 °C C	GT 1.0/20 °C C	GT 1.0/20 °C C	GT 1.0/20 °C C
1		GT 0.25/20 °C	1.90 ± 0.55a	0.90 ± 0.14a	0.95 ± 0.37a	7.30 ± 0.45f
		GT 0.5/20 °C	3.05 ± 0.70b	1.40 ± 0.29b	2.11 ± 0.69b	4.14 ± 0.47d
		GT 1.0/20 °C	4.20 ± 0.20c	5.75 ± 0.67d	4.00 ± 0.56d	2.36 ± 0.55b
3		GT 0.25/20 °C	3.65 ± 0.22b	2.30 ± 0.67b	3.14 ± 0.59c	4.80 ± 0.27d
		GT 0.5/20 °C	5.55 ± 2.08c	3.61 ± 0.90c	4.95 ± 0.84d	3.66 ± 0.52c
		GT 1.0/20 °C	8.00 ± 0.72d	7.58 ± 0.10e	7.21 ± 0.62e	1.38 ± 0.48a
7		GT 0.25/20 °C	4.46 ± 0.39c	3.38 ± 0.91c	3.32 ± 2.06c	5.50 ± 0.35e
		GT 0.5/20 °C	8.70 ± 0.98d	8.28 ± 0.70g	8.86 ± 0.92f	0.98 ± 0.13a
		GT 1.0/20 °C	8.80 ± 0.91d	7.72 ± 0.40f	8.55 ± 0.94f	2.30 ± 0.52b
10		GT 0.25/20 °C	7.66 ± 0.50d	4.90 ± 0.89d	4.70 ± 0.66d	5.80 ± 0.27e
		GT 0.5/20 °C	8.90 ± 0.82d	7.60 ± 0.66f	8.61 ± 0.99f	1.38 ± 0.21a
		GT 1.0/20 °C	8.50 ± 0.55d	7.60 ± 0.65f	6.71 ± 0.97f	1.17 ± 0.64a
Part II	Storage (days)	Treatment	General Browning	Photosynthetic Browning	Vascular Browning	Fresh appearance
ANOVA treatments			GT 0.25/20 °C B	GT 0.25/20 °C B	GT 0.25/20 °C A	GT 0.25/20 °C A
			GT 0.25/50 °C A	GT 0.25/50 °C A	GT 0.25/50 °C B	GT 0.25/50 °C B
1		GT 0.25/20 °C	1.90 ± 0.55ab	0.90 ± 0.14a	0.95 ± 0.37a	7.30 ± 0.45c
		GT 0.25/50 °C	1.00 ± 0.41a	0.69 ± 0.47a	0.87 ± 0.48a	8.86 ± 0.72e
3		GT 0.25/20 °C	3.65 ± 0.22c	2.30 ± 0.37c	3.14 ± 0.58d	4.80 ± 0.27a
		GT 0.25/50 °C	1.12 ± 0.25a	1.62 ± 0.48b	1.87 ± 0.25b	8.31 ± 0.19d
7		GT 0.22/20 °C	4.46 ± 0.38d	3.38 ± 0.61c	3.32 ± 0.96d	5.50 ± 0.35b
		GT 0.25/50 °C	2.30 ± 0.76 b	2.50 ± 0.51c	2.80 ± 0.27c	7.30 ± 0.67c
10		GT 0.25/20 °C	7.66 ± 0.50e	4.90 ± 0.49e	4.70 ± 0.25e	5.80 ± 0.27b
		GT 0.25/50 °C	3.62 ± 0.63c	3.12 ± 0.25d	3.25 ± 0.29d	6.50 ± 0.41b
Part III	Storage (days)	Treatment	General Browning	Photosynthetic Browning	Vascular Browning	Fresh appearance
ANOVA treatments			GT 0.25/20 °C A	GT 0.25/20 °C A	GT 0.25/20 °C A	GT 0.25/20 °C A
			Chlorine/20 °C A	Chlorine/20 °C A	Chlorine/20 °C B	Chlorine/20 °C A
1		GT 0.25/20 °C	1.95 ± 0.27a	0.95 ± 0.11a	0.85 ± 0.42a	7.35 ± 0.65d
		Chlorine/20 °C	2.02 ± 0.25a	0.87 ± 0.14a	0.94 ± 0.12a	7.25 ± 0.50d
3		GT 0.25/20 °C	3.50 ± 0.45b	3.20 ± 0.57b	4.04 ± 0.50c	5.40 ± 0.12c
		Chlorine/20 °C	3.65 ± 0.22b	2.30 ± 0.67b	3.14 ± 0.58b	4.80 ± 0.27b
7		GT 0.25/20 °C	4.40 ± 0.19c	3.85 ± 0.24c	5.30 ± 0.54d	5.55 ± 0.77c
		Chlorine/20 °C	4.46 ± 0.38c	3.38 ± 0.11c	3.32 ± 0.26b	5.50 ± 0.35c
10		GT 0.25/20 °C	7.50 ± 0.42d	3.95 ± 0.37c	5.55 ± 0.37d	5.10 ± 0.59c
		Chlorine/20 °C	7.66 ± 0.50d	4.40 ± 0.39d	4.70 ± 0.25c	5.80 ± 0.27c

Part I: GT at different concentrations (0.25, 0.5 and 1 g 100 mL⁻¹ at 20 °C). Part II: 0.25 g 100 mL⁻¹ GT at 20 °C and 50 °C. Part III: 0.25 g 100 mL⁻¹ GT at 20 °C and 120 ppm chlorine at 20 °C.

3.3. Browning

3.3.1. Colour measurement

Colour was monitored in all the samples during the entire storage (Figs. 4 and 5). Luminosity values (L^*) decreased during storage (Fig. 4A), which can be related to the appearance of browning. The concentration of the treatment (Part I) significantly affected luminosity and increasing GT concentration decreased luminosity values. This might indicate higher browning levels in the lettuce treated with higher GT concentrations. Increasing the temperature of the treatment from 20 °C to 50 °C (Part II) resulted in higher luminosity values of the samples (Part II, Fig. 4B), probably due to the inhibitory effect of heat-shock on enzymes involved in the enzymatic browning (Saltveit, 2001). Compared with a control treatment (chlorine), GT samples showed lower values in luminosity during the storage (Part III, Fig. 4C). The amount of polyphenols present in the Green tea might have favoured a quicker appearance of browning. Another colour parameter related to browning (Castaner, Gil, Ruiz & Artes, 1999) and to the breakdown of chlorophyll (Bolin & Huxsoll, 1991) is a^* value. A significant increase in a^* , indicating a shift from greenness to redness, was observed in all the samples during storage (Fig. 5). GT

concentration showed an effect from day 1 on this parameter, increasing a^* values with increasing concentration (Part I, Fig. 5A). As observed from luminosity results, heat-shock (50 °C) reduced the appearance of browning, which was reflected in lower a^* values during storage (Part II, Fig. 5B). No difference between GT and chlorine samples was observed (Part III, Fig. 5C).

3.3.2. Potential browning

Potential browning was monitored in all the samples during storage (Fig. 6). A significant increase of potential browning during storage was observed ($p < 0.05$). This increase in the concentration of phenolic compounds is caused by the wounding stress from the treatment (Hanotel, Fleuriot & Boisseau, 1995).

Lower potential browning was found in samples treated with the lowest GT concentration (GT 0.25 g L⁻¹) compared with the other two concentrations (0.5 g 100 mL⁻¹ and 1.0 g L⁻¹) (Part I, Fig. 6A). The combination of GT and heat-shock kept lower potential browning values than GT treatment at room temperature (Part II, Fig. 6B) those results are in agreement with findings from Murata, Tanaka, Minoura and Homma (2004) and Saltveit (2001). Samples treated with chlorine showed higher levels of potential browning than samples treated with

GT (Part III, Fig. 6C). GT is known for the high antioxidant capacity; however this resulted in a negative effect applied to a commodity affected by browning such as fresh-cut lettuce is, mainly due to the nature of this natural antioxidant. Green tea is rich in polyphenols, which act as substrate for browning-related enzymes.

3.3.3. Browning-related enzymes (peroxidase (POD) E.C.1.11.1.7 and polyphenol oxidase (PPO) E.C.1.10.3.1.)

Browning-related enzymes were measured as indicators of browning in all the samples after treatment and during the 10 days of storage (Fig. 7). A significant increase in PPO was observed during the storage. Samples treated with the highest concentrations of GT (0.05 and 1.0 g 100 mL⁻¹) showed lower levels of PPO activity (Part I, Fig. 7A) than samples treated with 0.25 g 100 mL⁻¹ GT. These results contrast with those obtained for the other browning markers (Figs. 4A, 5A, 6A and Table 1), showing higher browning incidence in the samples treated with increasing concentrations of GT. Samples treated with the GT at 50 °C (heat-shock) showed significantly ($p < 0.05$) lower PPO activity than those treated with GT at 20 °C (Part II, Fig. 7B). The inhibition of PPO activity due to heat-shock treatment has been previously reported (Saltveit, 2001). No significant percentage of PPO or POD activity is lost due to the direct effect of the treatment (Martin-Diana et al., 2005c). The reduction in the activity of PPO and POD might be due to an indirect effect caused by feedback inhibition by the lack of phenolic compounds (substrate) or due to a direct effect on a receptor (unknown) implicated

Table 2
Changes in ascorbic acid (mg g⁻¹) in fresh-cut lettuce stored at 4 °C for 10 days

Part I	Storage (days)	Treatment	Ascorbic acid (mg g ⁻¹)
ANOVA treatments	GT 0.25/20 °C A	GT 0.5/20 °C B	GT 1.0/20 °C B
	1	GT 0.25/20 °C GT 0.5/20 °C GT 1.0/20 °C	0.07 ± 0.001d 0.07 ± 0.006d 0.8 ± 0.001e
	3	GT 0.25/20 °C GT 0.5/20 °C GT 1.0/20 °C	0.06 ± 0.002c 0.06 ± 0.001c 0.07 ± 0.001d
	7	GT 0.25/20 °C GT 0.5/20 °C GT 1.0/20 °C	0.05 ± 0.003b 0.06 ± 0.001c 0.06 ± 0.001c
	10	GT 0.25/20 °C GT 0.5/20 °C GT 1.0/20 °C	0.04 ± 0.002a 0.05 ± 0.001b 0.05 ± 0.001b
Part II	Storage (days)	Treatment	Ascorbic acid (mg g ⁻¹)
ANOVA treatments		GT 0.25/20 °C B	GT 0.20/50 °C A
	1	GT 0.25/20 °C GT 0.25/50 °C	0.07 ± 0.001f 0.02 ± 0.001b
	3	GT 0.25/20 °C GT 0.25/50 °C	0.06 ± 0.002e 0.02 ± 0.000b
	7	GT 0.25/20 °C GT 0.25/50 °C	0.05 ± 0.003d 0.02 ± 0.001b
	10	GT 0.25/20 °C GT 0.25/50 °C	0.04 ± 0.002c 0.01 ± 0.001a
Part III	Storage (days)	Treatment	Ascorbic acid (mg g ⁻¹)
ANOVA treatments		GT 0.25/20 °C B	Chlorine/20 °C A
	1	GT 0.25/20 °C Chlorine/20 °C	0.07 ± 0.001e 0.05 ± 0.000c
	3	GT 0.25/20 °C Chlorine/20 °C	0.06 ± 0.002d 0.04 ± 0.000b
	7	GT 0.25/20 °C Chlorine/20 °C	0.05 ± 0.003c 0.03 ± 0.002a
	10	GT 0.25/20 °C Chlorine/20 °C	0.04 ± 0.002b 0.03 ± 0.000a

Part I: GT at different concentrations (0.25, 05 and 1 g 100 mL⁻¹ at 20 °C). Part II: 0.25 g 100 mL⁻¹ GT at 20 °C and 50 °C. Part III: 0.25 g 100 mL⁻¹ GT at 20 °C and 120 ppm chlorine at 20 °C.

Table 3
Changes in carotenoids (mg g⁻¹) in fresh-cut lettuce stored at 4 °C for 10 days

Part I	Storage (days)	Treatment	Carotenoids (mg g ⁻¹)
ANOVA treatments	GT 0.25/20 °C A	GT 0.5/20 °C B	GT 1.0/20 °C B
	1	GT 0.25/20 °C GT 0.5/20 °C GT 1.0/20 °C	0.41 ± 0.11a 0.99 ± 0.11c 1.01 ± 0.13c
	3	GT 0.25/20 °C GT 0.5/20 °C GT 1.0/20 °C	0.47 ± 0.04a 1.01 ± 0.17c 0.97 ± 0.14c
	7	GT 0.25/20 °C GT 0.5/20 °C GT 1.0/20 °C	0.41 ± 0.11a 0.70 ± 0.11b 0.80 ± 0.13b
	10	GT 0.25/20 °C GT 0.5/20 °C GT 1.0/20 °C	0.49 ± 0.19a 0.77 ± 0.07b 0.68 ± 0.06b
Part II	Storage (days)	Treatment	Carotenoids (mg g ⁻¹)
ANOVA treatments		GT 0.25/20 °C B	GT 0.25/50 °C A
	1	GT 0.25/20 °C GT 0.25/50 °C	0.41 ± 0.01c 0.09 ± 0.02b
	3	GT 0.25/20 °C GT 0.25/50 °C	0.47 ± 0.04c 0.07 ± 0.02b
	7	GT 0.25/20 °C GT 0.25/50 °C	0.41 ± 0.01c 0.02 ± 0.00a
	10	GT 0.25/20 °C GT 0.25/50 °C	0.49 ± 0.02c 0.01 ± 0.01a
Part II	Storage (days)	Treatment	Carotenoids (mg g ⁻¹)
ANOVA treatments		GT 0.25/20 °C B	Chlorine/20 °C A
	1	GT 0.25/20 °C Chlorine/25 °C	0.41 ± 0.01d 0.31 ± 0.01b
	3	GT 0.25/20 °C Chlorine/25 °C	0.47 ± 0.04d 0.35 ± 0.03c
	7	GT 0.25/20 °C Chlorine/25 °C	0.41 ± 0.01d 0.36 ± 0.02c
	10	GT 0.25/20 °C Chlorine/25 °C	0.49 ± 0.01d 0.24 ± 0.04a

Part I: GT at different concentrations (0.25, 05 and 1 g 100 mL⁻¹ at 20 °C). Part II: 0.25 g 100 mL⁻¹ GT at 20 °C and 50 °C. Part III: 0.25 g 100 mL⁻¹ GT at 20 °C and 120 ppm chlorine at 20 °C.

in the synthesis of PPO and POD. No differences between samples treated with chlorine and GT 0.25 g L⁻¹ (Part III, Fig. 6C) were found.

The GT controlled significantly ($p < 0.05$) the POD activity (Fig. 8, Part A). Higher levels of GT inhibited better the POD than lower GT levels. The temperature increase produced an inhibitory effect on this enzyme when the treatment was combined with GT 0.25 g 100 mL⁻¹ (Part II, Fig. 8B). Lower levels of POD activity were observed in samples treated with GT than in chlorinated samples (Part III, Fig. 8C). This inhibitory effect responds to the antioxidant capacity of the catechines, the main polyphenol compounds in GT, which can act as free radical-scavengers, quenching hydroxyl radicals (-OH) or superoxide anion radicals (O⁻) (Hanasaki, Ogawa & Fukui, 1994; Sichel, Corsaro, Scalia, Bilio & Bonomo, 1991; Aoshima, & Ayabe, 2007). In contrast with PPO, the polyphenols are not specific substrates of the POD enzyme.

3.4. Sensory analysis

Browning in photosynthetic and vascular tissues, general levels of browning and fresh appearance of the samples were evaluated by the sensory panel. The panel found differences in browning between samples treated with different GT concentrations. Increasing GT concentrations produced more visual browning on the samples and differences between treatments were observed throughout the storage (Table 1, Part I). Scores for fresh appearance also showed the unsuitability of high GT concentrations (0.5 and 1.0 g 100 mL⁻¹)

compared with 0.25 g 100 mL⁻¹ GT. Samples treated with GT and heat-shock showed less browning (Table 1, Part II). No differences between chlorine and GT-treated samples were observed (Table 1, Part III). Fresh appearance of samples treated with GT at 0.25 g 100 mL⁻¹ or chlorine received higher scores than samples treated with GT at higher concentrations, due mainly to the presence of browning.

3.5. Nutritional markers

3.5.1. Ascorbic acid

A significant ($p < 0.05$) decrease in ascorbic acid was observed during the storage. Increasing GT concentration showed a protective effect, avoiding ascorbic acid degradation (Table 2, Part I). This effect might be due to the antioxidant capacity of catechines; the aromatic rings with hydroxyl groups possess strong antioxidant properties (Peterson & Totlani, 2005). A significant ($p < 0.05$) decrease of the ascorbic acid content was observed after the treatment with heat-shock and GT (Table 2, Part II), probably due to the low stability of ascorbic acid at high temperatures (Nagy & Smooth, 1976; Dewanto, Wu, Adom & Liu, 2002). When small concentrations of GT (0.25 g 100 mL⁻¹) were compared with chlorine a larger protection effect was observed in the case of GT (Table 2, Part II).

3.5.2. Carotenoids

Similar behaviour as the ascorbic acid was observed for carotenoids (Table 3, Part I). Carotenoid concentration significantly ($p < 0.05$) dropped during the storage in all the treatments. The use of temperature (heat-shock) increased carotenoid degradation (Table 3, Part II). The use of GT (0.25 g 100 mL⁻¹) resulted in higher carotenoid levels, compared with the chlorine based treatment (Table 3, Part III).

4. Conclusion

The use of Green tea extracts, at the lowest concentration tested (0.25 g 100 mL⁻¹), is a feasible preservation method for minimally processed lettuce. Green tea treatment positively affected the preservation of ascorbic acid and carotenoid content of the samples. However, the use of high GT concentration (0.5 or 1.0 g 100 mL⁻¹) resulted in an unacceptable product when evaluated by the sensory panel. Browning appears to be an important factor involved in this negative effect.

Further research involving safety markers would be needed in order to study the efficacy of GT in controlling microorganism spoilage and pathogen growth. The use of GT in combination with other treatments, as for example heat-shock, is recommended by the authors.

Acknowledgements

This research was supported by a Technological Sector Research grant (2002–2006). The authors would like to thank Karen Roe for the assistance to the group in the laboratory part of the study. Thanks are also extended to Amcor Flexibles Europe (Bristol, UK) for kindly providing the packaging film.

References

- Abreu, M., Beirão-da-Costa, S., Gonçalves, E. M., Beirão-da-Costa, M. L., & Moldão-Martins, M. (2003). Use of mild heat pre-treatments for quality retention of fresh-cut 'Rocha' pear. *Postharvest Biology and Technology*, 30, 153–160.
- Agar, I. T., Massantini, R., Hess-Pierce, B., & Kader, A. A. (1999). Postharvest CO₂ and ethylene production and quality maintenance of fresh-cut kiwi slices. *Journal of Food Science*, 64(3), 433–440.
- Allende, A., & Artes, F. (2003). UV-C radiation as a novel technique for keeping quality of fresh processed 'Lollo Rosso' lettuce. *Food Research International*, 36, 739–746.
- Allende, A., & Artes, F. (2003). Combined ultraviolet-C and modified atmosphere packaging treatments for reducing microbial growth of fresh processed lettuce. *Food Science and Technology*, 36, 779–786.
- Official methods of analysis (pp. 690), 16th ed. (1995). Washington, DC: R. Association of Official Analytical Chemists.
- Aoshima, H., & Ayabe, W. S. (2007). Prevention of deterioration of polyphenol-rich beverages. *Food Chemistry*, 100, 350–355.
- Barry-Ryan, C., & O'Beirne, D. (1999). Ascorbic acid retention in shredded iceberg lettuce as affected by minimal processing. *Journal of Food Science*, 64(3), 498–500.
- Beaudry, R. M. (1999). Effect of O₂ and CO₂ partial pressure on selected phenomena affecting fruit and vegetable quality. *Postharvest Biology and Technology*, 15(3), 293–303.
- Beuchat, L. R. (1998). Surface decontamination of fruits and vegetables eaten raw: A review. Food Safety Unit, World Health Organisation, Brussels, Belgium. WHO/FSF/FOS/98.2.
- Beuchat, L. R., & Ryu, J. H. (1997). Produce handling and processing practices. *Emerging Infectious Diseases*, 3, 459–465.
- Bolin, H. R., & Huxoll, C. C. (1991). Effect of preparation procedures and storage parameters on quality retention of salad-cut lettuce. *Journal of Food Science*, 56, 416–418.
- Bozkurt, H. (2006). Utilization of natural antioxidants: Green tea extract and Thymbra spicata oil in Turkish dry-fermented sausage. *Meat Science*, 73, 442–450.
- Bruhn, C. (2000). Food labelling: Consumer needs. In J. Ralph (Ed.), *Food labelling*. Blanchfield Cambridge: Woodhead Publishing Limited.
- Cao, G., Sofic, E., & Prior, R. L. (1996). Antioxidant capacity of tea and common vegetables. *Journal of Agricultural and Food Chemistry*, 44, 3426–3431.
- Castaner, M., Gil, M. I., Ruiz, M. V., & Artes, F. (1999). Browning susceptibility of minimally processed Baby and Romaine lettuces. *European Food Research Technology*, 209, 52–56.
- Cherry, J. P. (1999). Improving the safety of fresh produce with antimicrobials. *Food Technology*, 53, 54–59.
- Cooper, R., Morre, J., & Morre, D. (2005). Medicinal benefits of green tea: Part II. Review of anticancer properties. *Journal of Alternative and Complementary Medicine*, 11(4), 639–652.
- Delaquis, P. J., Stewart, S., Toivonen, P. M. A., & Moyls, A. L. (1999). Effect of warm, chlorinated water on the microbial flora of shredded iceberg lettuce. *Food Research International*, 32, 7–14.
- Dewanto, V., Wu, X., Adom, K. K., & Liu, R. F. (2002). Thermal processing enhances the nutritional value of tomatoes by increasing total antioxidant activity. *Journal of Agricultural and Food Chemistry*, 50, 3010–3014.
- Galeazzi, M. A. M., Sgarbieri, V. C., & Constantinides, S. M. (1981). Isolation, purification and physicochemical characterization of polyphenoloxidases (PPO) from dwarf variety of banana (*Musa cavendishii*, L.). *Journal of Food Science*, 46, 150–155.
- García, E., & Barrett, D. M. (2002). Preservative treatments for fresh-cut fruits and vegetables. In O. Lamikanra (Ed.), *Fresh-cut fruits and vegetables*. Science, technology and market Boca Raton, Florida: CRC Press.
- García, A., Mount, J. R., & Davison, P. M. (2003). Ozone and chlorine treatment of minimally processed lettuce. *Food Microbiology and Safety*, 68(9), 2747–2751.
- Giner, J., Gimeno, V., Espachs, A., Elez, P., Barbosa-Cánovas, G. V., & Martín, O. (2000). Inhibition of tomato (*Lycopersicon esculentum* Mill.) pectin methylesterase pulsed electric fields. *Innovative Food Science & Emerging Technologies*, 1, 57–67.
- Gomez-Lopez, V. M., Devlieghere, F., Bonduelle, V., & Debevere, J. (2005). Intense light pulses decontamination of minimally processed vegetables and their shelf-life. *International Journal of Food Microbiology*, 103, 75–89.
- Graham, N. J. D. (1997). The role of ozone and potassium permanganate in drinking water treatment. 3rd international workshop on drinking water quality management and treatment technology, 5–6 March. Taipei, Taiwan.
- Gramza-Michalowska, A. G., & Regula, J. (2007). Use of tea extracts (*Camellia sinensis*) in jelly candies as polyphenols sources in human diet. *Asian Pacific Journal of Clinical Nutrition*, 16, 43–46.
- Gross, J. (1991). Carotenoids. Pigments in vegetables: Chlorophyllase and carotenoids (pp. 124–128). New York: Van Nostrand Reinhold.
- Guilbert, S., Gontard, N., & Gorris, L. (1996). Prolongation of the shelflife of perishable food products using biodegradable films and coating. *LWT Food Science and Technology*, 29, 10–17.
- Hamilton-Miller, J. M. (1995). Antimicrobial properties of tea (*Camellia sinensis* L.). *Antimicrobial Agents and Chemotherapy*, 39, 2375.
- Hamilton-Miller, J. M., & Shah, S. J. (2005). Medicinal benefits of green tea: Part I. Review of noncancer health benefits. *Journal of Alternative and Complementary Medicine*, 11(3), 521–528.
- Han, J., Gomes-Feitosa, C. L., Castell-Perez, E., Moreira, R. G., & Silva, P. F. (2004). Quality of packaged romaine lettuce hearts exposed to low-dose electron beam irradiation. *Food Science and Technology*, 37, 705–715.
- Hanasaki, Y., Ogawa, S., & Fukui, S. (1994). The correlation between active oxygens scavenging and antioxidative effects of flavonoids. *Free Radical Biology & Medicine*, 16, 845–850.
- Hanotel, L., Fleuriot, A., & Boisseau, P. (1995). Biochemical changes involved in browning of c-irradiated cut witloof chicory. *Postharvest Biology and Technology*, 5, 199–210.
- Izumi, H. (1999). Electrolyzed water as a disinfectant for fresh-cut vegetables. *Journal of Food Science*, 64, 536–539.
- Jacxsens, L., Devlieghere, F., Ragaert, P., Vanneste, E., & Debevere, J. (2003). Relation between microbiological quality, metabolite production and sensory quality of equilibrium modified atmosphere packaged fresh-cut produce. *International Journal of Food Microbiology*, 83, 263–280.
- Kabir, H. (1994). Fresh-cut vegetables. In A. L. Brods, & V. A. Herndon (Eds.), *Modified atmosphere food packaging* (pp. 155–160). Institute of Packaging Professionals.
- Kato-Noguchi, H., & Watada, A. E. (1997). Citric acid reduces the respiration of fresh-cut carrots. *Hortscience*, 32, 136.
- Kaur, C., & Kapoor, H. C. (2001). Antioxidants in fruits and vegetables: The millennium's health. *International Journal of Food Science and Technology*, 36, 703–725.
- Kays, S. J. (1991). *Postharvest physiology of perishable plant products*. New York: Van Nostrand Reinhold.
- King, A. D., & Bolin, H. R. (1989). Physiological and microbiological storage stability of minimally processed fruits and vegetables. *Food Technology*, 43, 132–135.

- Klein, B. P. (1987). Nutritional consequences of minimal processing on fruits and vegetables. *Journal of Food Quality*, 10, 179–193.
- Koseki, S., Yoshida, K., Kamitani, Y., Isobe, S., & Itoh, K. (2004). Effect of mild heat pre-treatment with alkaline electrolyzed water on the efficacy of acidic electrolyzed water against *Escherichia coli* O157:H7 and *Salmonella* on lettuce. *Food Microbiology*, 21(5), 559–566.
- Koukounaras, A., Diamantidis, G., & Sfakiotakis, E. (2008). The effect of heat treatment on quality retention of fresh-cut peach. *Postharvest Biology and Technology*, 48, 30–36.
- Lamikanra, O. (2002). Preface. In O. Lamikanra (Ed.), *Fresh-cut fruits and vegetables. Science, technology and market* CRC Press: Boca Raton, Florida.
- Loaiza-Velarde, J., Mangrich, M., Campos-Vargas, R., & Saltveit, M. (2003). Heat shock reduces browning of fresh-cut celery petioles. *Postharvest Biology and Technology*, 27, 305–311.
- Luna-Guzmán, I., & Barrett, D. M. (2000). Comparison of calcium chloride and calcium lactate effectiveness in maintaining shelf stability and quality of fresh-cut cantaloupes. *Postharvest Biology and Technology*, 19(1), 61–72.
- Martin-Diana, A. B., Rico, D., Barry-Ryan, C., Frias, J. M., Mulcahy, J., & Henehan, G. T. M. (2005). Comparison of calcium lactate with chlorine as a washing treatment for fresh-cut lettuce and carrots: Quality and nutritional parameters. *Journal of the Science of Food and Agriculture*, 85, 2260–2268.
- Martin-Diana, A. B., Rico, D., Barry-Ryan, C., Frias, J. M., Mulcahy, J., & Henehan, G. T. M. (2005). Comparison of calcium lactate with chlorine as a washing treatment for minimally processed lettuce and carrots: Quality and nutritional markers. *Journal of the Science of Food and Agriculture*, 85(13), 2260–2268.
- Martin-Diana, A. B., Rico, D., Frias, J. M., Barry-Ryan, C., Mulcahy, J., & Henehan, G. T. M. (2005). Effect of heat-shock on browning-related enzymes in minimally processed iceberg lettuce and crude extracts. *Bioscience, Biotechnology and Biochemistry*, 69(8), 2131–2139.
- Martin-Diana, A. B., Rico, D., Frias, J., Mulcahy, J., Henehan, G. T. M., & Barry-Ryan, C. (2006). Whey permeate as a bio-preservative for shelf life maintenance of fresh-cut vegetables. *Innovative Food Science & Emerging Technologies*, 7, 112–123.
- Martin-Diana, A. B., Rico, D., Barry-Ryan, C., Frias, J. M., Henehan, G. T. M., & Barat, J. M. (2007). Efficacy of steamer jet-injection as alternative to chlorine in fresh-cut lettuce. *Postharvest Biology Technology*, 45(1), 97–107.
- Martin-Diana, A. B., Rico, D., Henehan, G. T. M., Frias, J., Barat, J. M., & Barry-Ryan, C. (2007). Calcium for extending the shelf-life of fresh whole and minimally processed fruits and vegetables: A review. *Trends in Food Science & Technology*, 18(4), 210–218.
- McKay, D. L., & Blumberg, J. B. (2002). The role of tea in human health: An update. *Journal of the American College of Nutrition*, 21(1), 1–13.
- Masuda, T., Inaba, Y., Maekawa, T., Takeda, Y., Yamaguchi, H., Nakamoto, K., Kuninaga, H., Nishizato, S., & Nonaka, A. (2003). Simple detection method of powerful antiradical compounds in the raw extract of plants and its application for the identification of antiradical plant constituents. *Journal of Agricultural and Food Chemistry*, 51, 1831–1838.
- Molnar-Perl, I., & Friedman, M. (1990). Inhibition of browning by sulphur amino acids. 3. Apples and potatoes. *Journal of Agricultural and Food Chemistry*, 38(8), 1652–1656.
- Murata, M., Tanaka, E., Minoura, E., & Homma, S. (2004). Quality of cut lettuce treated by heat shock: Prevention of enzymatic browning, repression of phenylalanine ammonia-lyase activity, and improvement of sensory evaluation during storage. *Bioscience, Biotechnology and Biochemistry*, 68, 501–507.
- Nagy, S., & Smooth, J. M. (1976). Temperature and storage effects on percent retention and percent U.S. recommended dietary allowances of vitamin C in canned single-strength orange juice. *Journal of Agricultural and Food Chemistry*, 25(1), 135–138.
- Nissen, L. R., Byrne, D. V., Bertelsen, G., & Skibsted, L. H. (2004). The antioxidative activity of plant extracts in cooked pork patties as evaluated by descriptive sensory profiling and chemical analysis. *Meat Science*, 68, 485–495.
- O'Beirne, D., & Francis, G. A. (2003). Reducing the pathogen risk in MAP-prepared produce. In R. Ahvenainen (Ed.), *Novel food packaging techniques* (pp. 231–286). Cambridge, UK: Woodhead Publishing Limited/CRC Press LLC.
- Ohlsson, T. (1994). Minimal processing-preservation methods of the future — An overview. *Trends in Food Science and Technology*, 5, 341–344.
- Park, C. M., Hung, Y.-C., Doyle, M. P., Ezeike, G. O. I., & Kim, C. (2001). Pathogen reduction and quality of lettuce treated with electrolyzed oxidizing and acidified chlorinated water. *Journal of Food Science*, 66, 1368–1372.
- Peterson, D. G., & Tottani, V. M. (2005). Influence of flavonoids on the thermal generation of aroma compounds. In F. Shahidi (Ed.), *Phenolics in foods and natural health products* Washington, DC: American Chemical Society.
- Rico, D., Martin-Diana, A. B., Henehan, J., Frias, G. T. M., & Barry-Ryan, C. (2006). Effect of ozone and calcium lactate treatments on browning and textured properties of fresh-cut lettuce. *Journal of Food Science and Agriculture*, 86(13), 2179–2188.
- Rico, D., Martin-Diana, A. B., Barry-Ryan, C., Frias, J. M., Henehan, G. T. M., & Barat, J. M. (2008). Optimisation of steamer jet-injection to extend the shelf-life of fresh-cut lettuce. *Postharvest Biology Technology*, 48(3), 431–442.
- Rico, D., Martin-Diana, A. B., Barry-Ryan, C., Frias, J. M., Henehan, G. T. M., & Barat, J. M. (2008). Use of neutral electrolysed water (EW) for quality maintenance and shelf-life extension of minimally processed-lettuce. *Innovative Food Science & Emerging Technologies*, 9, 37–48.
- Rietveld, A., & Wiseman, S. (2003). Antioxidant effects of tea: Evidence from human clinical trials. *Journal of Nutrition*, 133, 3285–3292.
- Rosen, J. C., & Kader, A. A. (1989). Postharvest physiology and quality maintenance of sliced pear and strawberry fruits. *Journal of Food Science*, 54, 656–659.
- Saltveit, M. E. (2001). Wound induced changes in phenolic metabolism and tissue browning are altered by heat shock. *Postharvest Biology and Technology*, 21, 61–69.
- Seymour, I. J. (1999). Review of current industry practice on fruit and vegetable decontamination. Gloucestershire, UK: Campden & Chorleywood Food Research Association [review no. 14].
- Si, W., Gong, J., Tsao, R., Kalab, M., Yang, R., & Yin, Y. (2006). Bioassay-guided purification and identification of antimicrobial components in Chinese green tea extract. *Journal of Chromatography*, 1125, 204–210.
- Sichel, G., Corsaro, C., Scalia, M., Di Bilio, A. J., & Bonomo, R. O. (1991). In vitro scavenger activity of some flavonoids and melanin against O₂. *Free Radical Biology & Medicine*, 11, 1–8.
- Son, S. M., Moon, K. D., & Lee, C. Y. (2001). Inhibitory effects of various antibrowning agents on apple slices. *Food Chemistry*, 73, 23–30.
- Surjadinata, B. B., & Cisneros-Zevallos, L. (2003). Modeling wound-induced respiration of fresh-cut carrots (*Daucus carota* L.). *Journal of Food Science*, 68, 2735–2740.
- Tan, B. K., & Harris, N. D. (1995). Maillard products inhibit apple polyphenolase. *Food Chemistry*, 53, 267–273.
- Wei, C. I., Huang, T. S., Kim, J. M., Lin, W. F., Tamplin, M. L., & Bartz, J. A. (1995). Growth and survival of *Salmonella montevideo* on tomatoes and disinfection with chlorinated water. *Journal of Food Protection*, 58, 829–836.
- Varoquaux, P., & Wiley, R. (1994). Biological and biochemical changes in minimally processed refrigerated fruits and vegetables. In R. C. Wiley (Ed.), *Minimally processed refrigerated fruits and vegetables* (pp. 226–268). New York: Chapman and Hall.
- Vina, S. Z., & Chaves, A. S. (2006). Antioxidant responses in minimally processed celery during refrigerated storage. *Food Chemistry*, 94, 68–74.
- Yeo, S. G., Ahn, C. W., Lee, Y. W., Lee, T. G., Park, Y. H., & Kim, S. B. (1995). Antioxidative effect of tea extracts from green tea, oolong tea and black tea. *Journal of Korean Society of Food and Nutrition*, 24, 229–304.
- Zhang, L., Zhaoxin, L., Zhifang, Y., & Xiang, G. (2005). Preservation of freshcut celery by treatment of ozonated water. *Food Control*, 16, 279–283.