

2012-12

Effects of Salinity on the Toxicity of Ionic Silver and Ag-PVP Nanoparticles to *Tisbe Battagliai* and *Ceramium Tenuicorne*

Ailbhe Macken

Technological University Dublin


Hugh Byrne

Technological University Dublin, hugh.byrne@tudublin.ie

Kevin Thomas

Norwegian Institute for Water Research

Follow this and additional works at: <https://arrow.tudublin.ie/nanolart>

 Part of the [Environmental Health Commons](#), [Terrestrial and Aquatic Ecology Commons](#), and the [Toxicology Commons](#)

Recommended Citation

Macken, A., Byrne, H.J., Thomas, K.V., (2012) Effects of salinity on the toxicity of ionic silver and Ag-PVP nanoparticles to *Tisbe battagliai* and *Ceramium tenuicorne*, *Ecotoxicology and Environmental Safety*, 86, 101-110 (2012) doi:10.1016/j.ecoenv.2012.08.025

This Article is brought to you for free and open access by the NanoLab at ARROW@TU Dublin. It has been accepted for inclusion in Articles by an authorized administrator of ARROW@TU Dublin. For more information, please contact yvonne.desmond@tudublin.ie, arrow.admin@tudublin.ie, brian.widdis@tudublin.ie.



This work is licensed under a [Creative Commons Attribution-NonCommercial-Share Alike 3.0 License](#)

1 **Effects of salinity on the toxicity of ionic silver and Ag-PVP nanoparticles to *Tisbe***
2 ***battagliai* and *Ceramium tenuicorne***

3

4 **Ailbhe Macken^{ab}, Hugh J Byrne^a, Kevin V Thomas^b**

5

6 ^aFocas Research Institute, Dublin Institute of Technology, Dublin 8, Ireland.

7 hugh.byrne@dit.ie

8 ^bNorwegian Institute for Water Research (NIVA), Gaustadalléen 21, Oslo, NO-0349,

9 Norway. kevin.thomas@niva.no

10

11 *Corresponding author. Ailbhe Macken, Phone: +47 98215472, E-mail: ama@niva.no

12

13

14 **Abstract**

15 The toxic effects of polyvinylpyrrolidone (PVP) coated silver nanoparticles (Ag-NP_{PVP}) and
16 ionic Ag, to *Tisbe battagliai* (Tb) and *Ceramium tenuicorne* (Ct) were investigated and the
17 usefulness of standardised marine guidelines for ENP risk assessment were assessed. The
18 toxicity of Ag-NP_{PVP} [CtEC₅₀ = 26.6 µg/L, TbEC₅₀ = 7.9 µg/L] and Ag⁺ [CtEC₅₀ =
19 2312.2µg/L, Tb EC₅₀ = 90.9 µg/L] to both test species differed, with the silver ENPs being
20 more toxic. In contrast to Ag⁺ the toxicity of Ag-NP_{PVP} increased significantly with
21 increasing salinity, however, after thorough characterisation it was not possible to correlate
22 the behaviour of the particles with an increase in toxicity and salinity. The results suggest that
23 the observed toxicity is being elicited by the free ionic silver complexing in solution and also
24 from an unknown potential particle related effect.

25

26 **Keywords:** silver nanoparticles, marine ecotoxicity, salinity, *Tisbe battagliai*, *Ceramium*
27 *tenuicorne*

28

29

30

31 **1 Introduction**

32 Silver nanoparticles are one of the most widely used engineered nanomaterials (NMs)
33 employed in commercial applications and their wide use in consumer products has led to
34 them becoming a potentially important factor in environmental risk assessment (ERA) and
35 monitoring (Klaine et al., 2008; Meyer et al., 2010). Naturally occurring nanoparticles have
36 always existed in the environment without documented adverse effects (Murr et al., 2004;
37 Handy et al., 2008). Despite this, the risks associated with engineered nanoparticles (ENPs)
38 in the environment are becoming a significant concern (Moore, 2006; Ju-Nam and Lead,
39 2008; Tiede et al., 2009).

40 The ERA of NMs is a relatively new field and many questions about appropriate
41 assessment methods exist (Tiede et al., 2009; Bhatt and Tripathi, 2011). At present there is
42 inadequate information on the levels of nanomaterials in the environment (Paterson et al.,
43 2011). Most work to date has focused on laboratory based evaluation of risk, where few
44 interactions between the ENPs and natural environmental constituents occur. It is only in
45 recent years that scientists have started to incorporate realistic environmental exposures into
46 their laboratory work (Diegoli et al., 2008; Van Hoecke et al. 2011). Despite recent changes
47 to a more “environmentally relevant” approach in ENP ecotoxicity and fate assessment, it can
48 still be stated that, at present there exists no robust, reliable method for the quantification or
49 assessment of ENPs in the environment.

50 Most ENP ecotoxicological assessments have, up to now, focused on the potential
51 effects of ENPs in freshwater environments covering a wide array of species including fish
52 (Farkas et al., 2011; Farkas et al., 2010; Bilberg et al., 2010), daphnids (Naddy et al., 2007),
53 nematodes (Meyer et al., 2010), chironomids (Nair et al., 2011) and microalgae (Petit et al.,
54 2010). Due to the increased use of silver ENPs in consumer products there is a likelihood that
55 they will also end up in marine, estuarine and coastal environments. The possible adverse

56 effects elicited by these compounds must therefore be assessed employing suitable marine
57 test systems. A small number of studies employing marine species in the laboratory have
58 been performed (Galloway et al, 2010; Ringwood et al. 2010; Canesi et al., 2010), but there is
59 still a paucity of available data. Fabrega et al (2011) demonstrated that biofilm succession
60 was impeded on silver ENP treated biofilms, which in turn affected the abundance of major
61 bacterial groups that may cause further long term effect on the marine bacterial community.
62 In contrast Miao et al (2009) observed that silver ENPs formed non-toxic aggregates (>
63 0.22 μm) in seawater. Assessing the effects of ENP exposure to marine organisms poses new
64 and complex issues, hence, the importance of a realistic environmental assessment of both the
65 behaviour and effects needs to be emphasised.

66 The high ionic strength of seawater is likely to affect the behaviour of ENPs and may
67 prevent the introduced ENPs from becoming completely dispersed within the marine and
68 estuarine environment due to agglomeration (Gebauer and Treuel, 2011). There have been
69 some recent studies investigating the effects of salinity on ENPs revealing that only minute
70 changes in salinity may drastically affect the behaviour, effects and fate of ENPs (Matson,
71 2009). In the case of an estuarine environment, it is of even more importance that the
72 behaviour of ENPs and the effects of salinity be understood. Estuaries are constantly
73 changing and highly dynamic environments, in which organisms have to adapt to and tolerate
74 rapid changes in physico-chemical conditions over short periods of time. Estuaries and
75 coastal environments are also the most likely of the marine ecosystems to be impacted by
76 potential ENP contamination (Kaegi et al., 2008).

77 Silver ENPs are of particular interest in the marine environment, not only because of
78 their abundant use and potential entry into this ecosystem, but also because the speciation
79 behaviour of silver in saline waters increases bioavailability (Luoma et al., 1995). Upon
80 entering the marine environment, silver speciation becomes a significant factor (Cowan et al.,

81 1985). Therefore, in the marine and estuarine environment, silver is unlikely to remain in the
82 form in which it was introduced. Another potential effect on the behaviour of ENPs in saline
83 media, is the fact that their tendency to aggregate/agglomerate may increase with increased
84 salinity.

85 The primary production of micro and macro algal species in the marine environment
86 is of paramount importance. Effects on primary production within an ecosystem could lead to
87 devastating effects on ecosystem functioning. Sub-lethal effects on these organisms may also
88 lead to transfer and biomagnification of metal contaminants up the food chain (Quigg, 2008).
89 At present, research on the interactions and toxicity of ENPs to macrophytes is lacking for
90 both the terrestrial and aquatic ecosystems. Due to this lack of information on higher plants,
91 *Ceramium tenuicorne*, a marine rhodophyte, was selected for use in this study. *C. tenuicorne*
92 was selected as it represents a primary producer in both the marine and estuarine environment
93 (i.e. euryhaline species) and could be used to assay the ENPs under a wide range of salinities.
94 *Ceramium* species are found world-wide and are particularly abundant in temperate climates.
95 The marine clone employed in this study originated from the Oslofjord (20 – 25 ‰), an inlet
96 in the south-east of Norway, and has been maintained as a laboratory culture for over 30
97 years and can be adapted to a wide range of environmentally relevant salinities (ISO, 2010).

98 In order to augment the multi-trophic nature of this study, a marine harpacticoid
99 copepod *Tisbe battagliai*, was incorporated into the testing. *T. battagliai* represents a primary
100 consumer within the marine food web and is of environmental significance because, along
101 with nematodes, copepods are the most abundant multicellular organisms on earth (Humes,
102 1994). It is also a standard test species in regulatory testing and employed in ERA. To the
103 best of our knowledge, no data regarding the effects of ENPs to *T. battagliai* and *C.*
104 *tenuicorne* exist in the literature. Therefore, the information gained in this work is considered

105 to contribute to the knowledge gap in understanding the effects of ENPs to these marine
106 species.

107 Although there are many studies in the literature on the ecotoxicological assessment
108 of ENPs, there is no standardisation in the methodologies used between different laboratories.
109 Significant factors, such as sample preparation and ENP characterisation, vary significantly
110 between laboratories. This variation leads to considerable differences in results for similar
111 particles and incomparable inter-laboratory data, for both ecotoxicological tests and
112 characterisation (Roebben et al., 2011, Petersen and Henry, 2012). The applicability of
113 standard guidelines for the assessment of ENPs in the environment is of great interest at
114 present. Several major international standards organisations (e.g. OECD and ISO) are
115 currently investigating the applicability of the standard methods for use in ERA and
116 regulation of ENPs. Due to the potential for more rigorous ERA of ENPs in the future, where
117 possible the studies performed in this work were performed in line with standard regulatory
118 guidelines (ISO 14669, ISO 10710) and according to GLP (Good Laboratory Practice, NIVA
119 has a GLP certified ecotoxicological laboratory), any deviations are described in full.

120 The main objectives of this study were to: (1) investigate the toxicity of silver nitrate
121 (AgNO_3) and polyvinylpyrrolidone coated silver nanoparticles ($\text{Ag-NP}_{\text{PVP}}$) to two marine
122 species and to fully support all ecotoxicological studies with a thorough characterisation of
123 the ENPs in all environmental media; (2) to investigate species specific differences in
124 sensitivity; (3) evaluate the potential modifying effects of varying salinity on the toxicity of
125 AgNO_3 and $\text{Ag-NP}_{\text{PVP}}$ to *C. tenuicorne* and (4) to place the results in an environmentally
126 relevant context and to discuss the applicability of these standardised test guidelines for use
127 in the regulatory assessment of engineered NPs within the marine and estuarine environment.

128

129 **2 Materials and Methods**

130 **2.1. Nanoparticles and test chemicals**

131 The silver nanoparticles employed in this study were coated with
132 polyvinylpyrrolidone (PVP). These PVP (Ag-NP_{PVP}) capped ENPs were kindly prepared and
133 provided in a solution of MilliQ water by the University of Manchester, UK. Post analysis of
134 the stock solution by ICP-MS, indicated that the concentration was 348 mg/L, based on a
135 measurement of total silver.

136 Four different types of media were used in these studies: non-enriched natural
137 seawater (NSW) for the *T. battagliai* assays and an enriched algal media for *C. tenuicorne*
138 assays (Eklund, 2005, ISO, 2010), at three different salinities (to be referred to as CT10 ‰,
139 CT20 ‰, and CT30 ‰). Natural seawater, collected from 60 m depth at Solbergstrand,
140 Norway (ca. 34 ‰), was employed in the preparation of all media. All media were filtered to
141 0.2 µm prior to use. Test solutions of ENPs for use in the ecotoxicity assessments were
142 prepared by diluting the stock solution in the appropriate saline media. Dilutions were made
143 directly from the stock solution, where possible, in order to reduce the effects of potential
144 aggregation/agglomeration during serial dilution. All test solutions were stirred for 3 hours at
145 100 rpm prior to characterisation and use in ecotoxicology tests.

146 The reference chemical, analytical grade potassium dichromate (K₂Cr₂O₇; CAS
147 Registry No 778-50-9) was obtained from Sigma Aldrich. Polyvinylpyrrolidone (PVP; CAS
148 Registry No 9003-39-8) also obtained from Sigma Aldrich, was employed as a control for the
149 nanoparticle capping agent.

150

151 **2.2. Nanoparticle characterisation**

152 There are many techniques for measuring the size of nanoparticles, all of which are
153 based on fundamentally different scientific principles. It has previously been reported that no
154 single technique is sufficient to accurately estimate the size of nanoparticles in all media
155 types (Domingos et al., 2009 Scalf and West, 2006). Each individual technique has intrinsic
156 biases and artefacts produced during sample preparation, or from instrumental interference
157 (Domingos et al. 2009). Therefore, in this study, a battery of different characterisation
158 techniques was employed to investigate the size and behaviour of the Ag-NP_{PVP} in saline
159 media.

160

161 **2.2.1. Dynamic Light Scattering (DLS)**

162 Dynamic light scattering (DLS) was used to measure the hydrodynamic diameter (Z-
163 average), the intensity based size distribution and the zeta potential of Ag-NP_{PVP} in MilliQ
164 water and all 4 media types. The particle size distributions (PSD) of the Ag-NP_{PVP} were
165 analysed using a Malvern Instruments Zetasizer Nano Series (Malvern Instruments, UK).
166 Concentrations of 100 µg/L were prepared in all media and in MilliQ water. Samples were
167 analysed at 20 °C. Typically, 1 mL of solution was placed into a disposable sizing cuvette and
168 samples for all media were prepared and analysed in triplicate. Six measurements of each
169 triplicate sample were taken for all media.

170 Using the same Malvern Instrument, zeta potential measurements were also
171 performed on the same solutions described above. Folded capillary cells with ca. 3 mL of
172 each solution were employed. Measurements were conducted at 20 °C, using a concentration
173 of 100 µg/L silver ENP and 6 measurements on triplicate samples were made.

174

175 **2.2.2. Atomic Force Microscopy (AFM)**

176 Atomic force microscopy (AFM) was used to resolve particles and groups of particles
177 by offering visualisation in three dimensions. Samples of Ag-NP_{PVP} were prepared in all
178 media types and MilliQ water at a concentration of 10000 µg/L. Samples were drop cast onto
179 silicon wafers which were subsequently rinsed with de-ionised water and imaged in tapping
180 mode with a high performance AFM microscope (Asylum MFP-3D-BIO™).

181

182 **2.2.3. Transmission Electron Microscopy (TEM)**

183 Solutions of the highest working concentrations of silver ENPs were prepared in all
184 test media. The size and shape of the particles were determined by TEM. The ENP
185 suspensions were prepared on the day of analysis in appropriate test media and applied to
186 carbon coated grids (5 µl pipetted directly onto the grids). The grids were allowed to dry and
187 images were captured with a Philips CM 100 electron microscope. Using image analysis
188 software (Digimizer Version 3.4.1.0) to analyse micrograph images, 100 ENPs were
189 measured and the mean particle size and PSD of Ag-NP_{PVP} in all media described.

190

191 **2.3. Ecotoxicity testing**

192 **2.3.1. *Tisbe battagliai* culturing and testing**

193 *T. battagliai* were cultured in the laboratory at NIVA in accordance with standard
194 procedures (ISO, 1999) using NSW of ca. 34 %. Cultures were fed a single algal diet
195 consisting of *Rhodomonas baltica* once per week (additional feeding may be required
196 depending on the ability of the animals to clear the food) during culture water renewal.

197 *T. battagliai* toxicity tests were conducted with slight modifications according to the
198 ISO method (ISO, 1999). Toxicity tests were conducted with copepodids 6 ± 2 days old. The
199 bioassays were performed in 12-well polystyrene tissue culture plates (NUNC[®]) that had
200 been pre-treated overnight with appropriate exposure solutions. After initial range finding
201 studies, both Ag-NP_{PVP} and AgNO₃ were tested over the following range 1, 3.2, 5.6, 10, 32,
202 56, 100 µg/L (7 concentrations plus a negative control [NSW]). Controls for PVP were run
203 alongside to confirm that any observed toxicological effects with Ag-NP_{PVP} were not due to
204 the presence of the capping agent. All concentrations were run in quadruplicate (2.5 mL of
205 exposure solution per replicate with five test organisms per replicate, a total of 20 animals per
206 test concentration). Test plates were incubated in a temperature controlled room at 20 ± 2 °C
207 and a 16:8 hour light:dark photoperiod. Experiments were performed on three independent
208 occasions.

209 A reference substance, potassium dichromate (K₂Cr₂O₇), was run alongside the tests
210 in order to verify the sensitivity of the copepods and to ensure the test conditions were
211 reliable (ISO, 1999). At test initiation and termination, dissolved oxygen concentrations, pH
212 and salinity were measured in the control, top and bottom concentrations for all tests. Control
213 species mortality of less than or equal to 10 % was required as a further validation parameter.

214

215 **2.3.2 *Ceramium tenuicorne* adaptation**

216 A sample of *C. tenuicorne* (ITM, Stockholm) was supplied for use in this study (20
217 ‰), with a pre-testing adaptation period, to ensure good growth of the clone at all proposed
218 test salinities.

219 Adaptation of the plants to 10 and 30 ‰ was conducted over several weeks prior to
220 testing. This was done by successively transferring the algae every three days to an

221 increase/decrease of approximately 3 %. The algae were then cultivated for at least two
222 weeks in the final test salinity prior to test initiation.

223 Prior to testing, the growth rate of the algae at the varying test salinities was
224 determined to ensure sufficient growth during the test period. This was carried out by
225 conducting a 7 day growth experiment according to the ISO guideline (ISO, 2010) with
226 exponentially growing female tips (0.6 – 1.2 mm start length) in all media without test
227 chemical. All media growth experiments were conducted in triplicate and measurements, at
228 test initiation and termination, were taken using an Olympus 1x71 fluorescent microscope
229 (without fluorescent mode) (Tokyo, Japan) and Olympus Soft Imaging system: Cell[^]D,
230 Version 3.2 (Build 1700).

231

232 **2.3.3 *Ceramium tenuicorne* silver toxicity and salinity effects on growth rate**

233 The toxicity of Ag-NP_{PVP} and AgNO₃ as well as the effects of salinity on these
234 toxicities, to the female plants, were studied using pre-adapted *C. tenuicorne* (section 2.3.2).

235 After initial range finding studies, definitive toxicity tests at all salinities were
236 conducted within the range 1, 5, 10, 50, 100, 500, 1000, 5000, 10000 µg/L (9 concentrations
237 and appropriate negative controls), for both Ag-NP_{PVP} and AgNO₃. As with the *T. battagliai*
238 assay, a PVP control was incorporated into the test design. All toxicity testing was conducted
239 according to modified standard methods (Eklund, 2005; ISO, 2010). Due to the small amount
240 of test material available for Ag-NP_{PVP}, exposure volumes were reduced to 2.5 mL per
241 replicate (4 replicates/concentration). 12-well pre-treated tissue culture plates (NUNC[®]) were
242 used for the exposures. All concentrations were run in quadruplicate with two algal plants per
243 replicate. Microplates were incubated for seven days at a temperature of 22 ± 2 °C, a light
244 intensity of 70 ± 10 % µmol m⁻² s⁻¹ and a continuous light regime. In order to prevent
245 evaporation (due to the small volume size) plates were sealed with parafilm. Experiments

246 were conducted for all salinities, for both silver compounds, on three independent occasions
247 to ensure reproducibility and increase statistical power. A reference chemical (potassium
248 dichromate) was run alongside to ensure performance and reliability of test methods. At test
249 initiation and termination, dissolved oxygen concentrations, pH and salinity were measured
250 in the control, top and bottom concentrations.

251

252 **2.4 Statistical analysis**

253 The EC₁₀/EC₅₀ (concentration that elicits an estimated 10 %/50 % toxic effect (i.e.
254 growth inhibition, mortality) values for Ag-NP_{PVP}, AgNO₃ and K₂Cr₂O₇ were calculated
255 using REGTOX-EV6.xls (Eric Vindimian <http://eric.vindimian.9online.fr/>), a curve fitting
256 macro for Microsoft[®] Excel. Toxicity data for the algal and copepod tests were fitted to a
257 sigmoidal curve and the Weibull (algal assays) and Hill (copepods assays) models were used
258 to calculate Lethal Concentration (LC) and Effective Concentration (EC) values respectively.
259 Statistical analyses were carried out using analyses of variance (ANOVA) followed by
260 Dunnett's multiple comparison test. From these data, Lowest Observable Effects
261 Concentrations (LOEC) and No Observable Effect Concentrations (NOEC) were calculated.
262 These data analyses were performed using MINITAB[®] release 15 (MINITAB Inc. PA, USA).
263 Statistical significance was accepted at $p \leq 0.05$.

264

265 **3 Results**

266 **3.1. Nanoparticle characterisation**

267 **3.1.1. Dynamic Light scattering (DLS)**

268 The average particle sizes of Ag-NP_{PVP} in all media and MilliQ water, as measured by
269 DLS, are shown in Table 1. Ag-NP_{PVP} in MilliQ water showed hydrodynamic peaks at
270 around 6 and 60 nm (Z-average value of 56.9 nm). In the case of the saline media, the values
271 varied and there appeared to be no average particle size increase/decrease trend with
272 increasing salinity. The CT10 ‰ media showed peaks at around 40 and 80 nm (Z-average
273 value of 149 nm). The CT20 ‰ media showed two peaks in the size distribution of
274 approximately 5 and 60 nm (Z-average value of 57.28 nm). The CT30 ‰ media showed two
275 peaks in the size distribution at ca. 40 and 120 nm (Z-average value of 105.01 nm). The
276 highest salinity media (NSW) showed two main size distribution peaks, at around 28 and 154
277 nm (Z-average value of 76.44 nm). All size determination results are summarized in Table 1.
278 Although the polydispersity index was below 0.5 for all samples measured, indicating that Z-
279 average is a suitable mean size to use to compare these samples, the presence of multiple
280 peaks, indicates that a degree of aggregation/agglomeration occurs in the media and the Z-
281 average as determined by DLS may not be the most appropriate technique for representing
282 the size distribution of these particles in solution

283 For zeta potential measurements, it was only possible to obtain usable values in the
284 MilliQ, CT10 ‰ and CT20 ‰ media samples, as the high conductivity of the samples with a
285 salinity above 20 ‰ caused interference in the readings. The zeta potential of the particles
286 measured in the MilliQ water and *C. tenuicorne* media at 10 ‰ and 20 ‰ were -39.17 mV, -
287 8.68 mV and 10.32 mV respectively. The low zeta potential values are consistent with a
288 tendency for the nanoparticles to agglomerate or aggregate, as indicated by the size
289 distribution measurements.

290

291 3.1.2 Atomic Force Microscopy (AFM)

292 AFM images were captured and analysed for drop cast samples of Ag-NP_{PVP} from all
293 media and MilliQ water. Figure 1 shows an example of an AFM image and corresponding
294 topography of a cross section for Ag-NP_{PVP} from CT10%. As seen from Figure 1, the silver
295 nanoparticles form isolated aggregates when precipitated from the media. The cross section
296 of the image shows one of these aggregates with a height greater than 150 nm and a base
297 diameter of > 0.5 μm . However, a significant number of features of dimensions $\sim 10\text{nm}$ are
298 also observed, indicating the presence of isolated nanoparticles. The average size and particle
299 size distribution are described in Table 1.

300

301 **3.1.3 Transmission Electron Microscopy (TEM) analysis**

302 While both DLS and AFM indicate a significant degree of aggregation/agglomeration
303 in the dispersions, TEM images indicate the presence of particles in the nanometer range (<
304 100 nm, ASTM International, 2006) in the stock solution (Farkas et al., 2010) and in the
305 analysed exposure solutions 10000 $\mu\text{g/L}$. All TEM measurements are described in Table 1.
306 The presence of nanoparticles is clearly seen in the MilliQ water sample, in which
307 approximately 48 % of particles were < 5nm, while, 30 % were between 5 – 10 nm in size.
308 Of the exposure media, the smallest average size, based on TEM analysis, was for CT10 %
309 media with an average size of 16.0 (± 5.8) nm. The average size for the CT20 % media,
310 CT30 % media and the NSW (ca. 34 %) were 20.2 (± 4.1), 19.3 (± 6.3) and 18.9 (± 5.2) nm
311 respectively. As with the DLS analysis, there was no specific trend in particle size based on
312 increasing/decreasing salinity. The Ag-NP_{PVP} in all saline media (except in CT30 %, where
313 only dispersed particles were seen) were observed to form loose agglomerates at a
314 concentration of 10000 $\mu\text{g/L}$ as well as being present as monodisperse particles in solution
315 (Figure 2). In the calculated PSD (Figure 3), results indicated that there were no particles less

316 than 5 nm in diameter in any media type. The size distributions of the analysed particles were
317 similar for all media, 7.9 – 33.7, 11.4 – 36.3, 9.7 – 39.6 and 9.9 – 34.9 nm respectively, for
318 CT10 %, CT20 %, CT30 % and NSW. For all media types between 87 - 98 % of the
319 measured particles were in the range 10 – 30 nm in diameter (Figure 3).

320

321 **3.2 Ecotoxicity testing**

322 **3.2.1 *Tisbe battagliai***

323 The mean 48 hour EC₅₀ value for the reference chemical potassium dichromate with
324 *T. battagliai* was 7.2 (95 % CI = 5.8 – 9.1) mg/L, which corresponds to previously published
325 values for this species and chemical (Macken et al., 2008). No deleterious effects of PVP
326 were observed, therefore, toxicity due to the PVP capping agent in these experiments can be
327 ruled out.

328 Toxicity data for Ag-NP_{PVP} and AgNO₃ to *T. battagliai* are shown in Figure 4a and b
329 and Table 2. The LC₁₀ and LC₅₀ values indicate that the toxicity of Ag-NP_{PVP} was statistically
330 greater than the AgNO₃ toxicity. Despite the calculation of these values it can be seen from
331 Figure 4 that the silver salt had a greater toxic effect at higher concentrations, while the silver
332 ENP appeared to be more toxic at lower concentrations (see Figure 4c for a graphical
333 comparison of the mortality data). For all experiments, the test vessels were treated with
334 solutions of suitable concentrations overnight prior to test initiation and rinsed with MilliQ
335 water. This was done so as to remove any potential binding sites for the silver test solutions
336 on the plastic test vessel surfaces. The control mortality and physico-chemical parameters
337 measured at the start and end of all experiments were within the recommended limits (ISO,
338 1999).

339

340 3.2.2 *Ceramium tenuicorne*

341 The mean 7 day EC₅₀ value for the K₂Cr₂O₇ with *C. tenuicorne* was 3.9 mg/L (95 %
342 CI = 2.6 - 5.7), which was consistent with previous studies with this species (Macken,
343 unpublished) and no toxicity was observed with the PVP capping agent. The mean increase in
344 length over a 7 day test period, of six algal pieces, in the three proposed test salinities was
345 measured. Growth rate was observed to increase with increasing salinity, 1.12, 1.46 and 1.62
346 respectively for 10, 20 and 30 ‰ media. All salinities allowed for good growth of the algae,
347 sufficient to meet all validity criteria for standard guideline testing (ISO, 2010), however, for
348 continuous cultures, it was observed that the 20 ‰ cultures were the best performing for
349 long-term culture maintenance.

350 The control growth rates met with the validity criteria in all experiments (increase in
351 length over seven days by a factor of greater than 3, compared to the starting length [ISO,
352 2010]), therefore implying that the use of parafilm and reduced sample volume had no
353 adverse effects on the test system. Table 2 summarises the ecotoxicity data for all test
354 salinities with both Ag-NP_{PVP} and AgNO₃. As was observed with the *T. battalgiai* assays, the
355 toxicity of the silver ENPs was greater than that of AgNO₃ (Table 2 and Figure 5) to the test
356 organism when looking at EC values. Despite this, when NOEC/LOEC values were
357 calculated, the difference in toxicity changed and the ENPs can only be considered more
358 toxic at the highest salinity (CT30 ‰). The toxicity of the Ag-NP_{PVP} increased with
359 increasing salinity (EC₅₀ values of 1872.5 (95 % CI = 954.5 – 3757.6), 393.5 (95 % CI = 954.5 –
360 3757.6) and 26.7 (95 % CI = 12.6 – 69.8) µg/L, at CT10 ‰, CT20 ‰ and CT30 ‰,
361 respectively). The observed toxicity of AgNO₃ to *C. tenuicorne* at all salinities was similar,

362 with a slight, but not statistically significant, increase in toxicity between 10 ‰ and 20 ‰
363 (based on EC values).

364

365 **4 Discussion**

366

367 The toxicity of both the ionic silver and Ag-NP_{PVP} assayed in this study were high
368 compared to other similar studies in the literature. No comparable toxicity data for *T.*
369 *battagliai* and silver or silver ENPs are available, however, there is some information on
370 other marine copepods. Pedroso et al. (2007) found that at 30 ‰ the 48 h EC₅₀ for *Acartia*
371 *tonsa* was 173 µg/L (dissolved silver) indicating the harpacticoid, *T. battagliai* was more
372 sensitive to silver toxicity (48 h EC₅₀ at 34 ‰ = 90.9 µg/L). In relation to species specific
373 differences, there are limits to identifying the most sensitive of the two species assayed in this
374 study. Different salinities, exposure periods and endpoints (e.g. lethality and growth) are
375 being assessed. However, it is clear that the sensitivity of the two species differs depending
376 on these factors and it is important to incorporate a battery of endpoints and exposure periods
377 for ERA in order to allow for the protection of the marine and estuarine environment.

378 The higher toxicity of silver ENPs compared to ionic silver, with *T. battagliai* may be
379 indicative of a potential particle specific mode of action on the toxicity and has been
380 observed by other authors (Griffitt et al., 2008; Navarro et al., 2008). The increase in toxicity
381 could be as a result of increased surface area and therefore increased capacity to release ionic
382 silver from the surface of the particle.

383 *C. tenuicorne* was also observed to be slightly more sensitive to Ag-NP_{PVP} than ionic
384 silver, however, differences in sensitivities were not as evident as those observed with

385 *T. battagliai* at all salinities. After exposure in algal media of 10 ‰, the EC₁₀/EC₅₀ values
386 were similar for both Ag-NP_{PVP} and AgNO₃. The difference in sensitivity between the two
387 test chemicals became much more pronounced with increasing exposure media salinity.
388 These differences were particularly pronounced at 30 ‰. There did not appear to be a marked
389 increase in toxicity of AgNO₃ with increasing salinity. In contrast, there was a consistent
390 increase in toxicity up the salinity gradient with Ag-NP_{PVP}. The toxicity of metals in aquatic
391 systems is influenced by a variety of factors such as organic matter content (Nadella et al.,
392 2009) and salinity (Verslycke et al., 2003). These results may indicate that an increase of
393 chloride ions or other inorganic ligands within the media are changing the behaviour and
394 bioavailability of the silver ENPs to the test species. Previously published work, investigating
395 the effects of salinity on the toxicity of contaminants in the marine environment to
396 invertebrates, have observed an increase in toxicity with decreasing salinity (Verslycke et al.,
397 2003; Kwok and Leung, 2005; Pedroso et al. 2007). Ytreberg et al. (2011) investigated the
398 effects of salinity and organic matter on the toxicity of Cu to *C. tenuicorne* and concluded
399 that the effects of salinity on Cu toxicity were not clear, as both a positive and negative effect
400 was observed, a decrease in toxicity with increasing salinity in the presence of organic matter
401 but no decrease in the absence of organic matter.

402 As in this work, an increase in toxicity with increasing ionic strength has also been
403 observed in the literature (Erikson et al., 1998). This is the opposite of what would be
404 expected if only the free silver ions were responsible for the toxic response. Due to the high
405 ionic strength of seawater and the prevalence of inorganic ligands such as chlorine, silver
406 complexation will not be dominated by NOM, as in freshwater, but instead speciation is
407 dominated by the formation of strong chloro-complexes (Cowan et al., 1985; Cowan et al.,
408 1993; Reinfelder and Chang, 1999). Without definite information on the speciation and
409 complexation of silver in the media and seawater in our study, it is not possible to make clear

410 conclusions on the causes of the observed toxicity or the effects of changing ionic strength.
411 Unfortunately due to the low levels of silver, limitations in analytical capabilities and small
412 volumes of test material, speciation could not be investigated in greater detail in this
413 particular study.

414 As well as understanding the speciation of silver within seawater, an understanding of
415 the chemical nature of the exposure medium is also important in determining the behaviour
416 and bioavailability of silver and silver nanoparticles. In this study all of the exposure
417 solutions were prepared from NSW, however, the algal media was also enriched with a
418 variety of trace metals, salts and vitamins, including nitrogen, phosphorus, iron, and trace
419 amounts of cobalt, copper, zinc and manganese (ISO, 2010) One important factor for
420 consideration is the presence of chelating agents within exposure media (e.g. EDTA). In this
421 work the ISO *C. tenuicorne* media contained low levels of EDTA (final concentration of
422 approximately 300 µg/L). Despite this, it is unlikely that the presence of EDTA can
423 completely explain the difference in sensitivity between the two species, as EDTA has been
424 shown to have very a low affinity for silver (Zuiderveen and Birge, 1995).

425 In order to further try and explain the results of our study we have considered the
426 potential physiological susceptibility of the algae itself under differing environmental
427 regimes. Russell (1985) conducted an investigation of macroalgae in the Baltic Sea and found
428 *C. tenuicorne* was the least likely to cope with any large increase in salinity. Despite, the
429 algae used in our study going through an adaptation period of several weeks, this may help to
430 explain the increased sensitivity of this species with increasing salinity. In order to respond to
431 external salinity changes this particular seaweed, has to alter its water content and the
432 concentration of inorganic ions (Lobban and Harrison, 1994; Ferguson and Hogstrand, 1998).
433 Due to the fact that the Ag-NP_{PVP} toxicity increased with increasing salinity and the AgNO₃

434 toxicity did not increase, there could also be some sort of ENP effect that is enhanced in the
435 presence of increasing Cl⁻.

436 It is believed that no single technique for ENP characterisation is without its artefacts
437 or can be employed in all cases for nanoparticle characterisation (Domingos et al., 2009).
438 Therefore, in order to try and counteract this problem a battery of characterisation techniques
439 were employed in this study. The dispersion system employed in this study was intended to
440 reflect the realistic conditions and behaviour of the ENPs on entry into the marine
441 environment and so no manipulations to force the ENPs into solution were conducted (e.g.
442 use of solvents) (Kato et al., 2009).

443 In recent years, TEM has allowed scientists to obtain images of individual
444 nanoparticles. However, during this analysis only a small fraction of the total sample is
445 characterised. Therefore, the results may not be representative of the total sample being
446 assayed and it is easy to acquire incorrect information on exposure solutions (Nowack and
447 Bucheli, 2007). There are several potential artefacts associated with TEM and ENP
448 characterisation, e.g. meniscus-based artefacts and co-precipitation with inorganic salts
449 (Domingos et al., 2009). In saline media, the same effect could cause interference with the
450 visual analysis of the sample, as any impurities, or salt in the sample, could make it difficult
451 to distinguish the ENPs from these contributions on the air-dried grid.

452 Compared to the TEM results, the results of DLS measurements of the average
453 particle size of Ag-NP_{PVP} in all media types, were far greater. Similar results have been
454 observed by other authors (Domingos et al., 2009; Farkas et al., 2010; Hassellöv et al., 2008).
455 It is important to note, that in DLS the diameter being measured is the hydrodynamic
456 diameter, which is not only dependent on the core of the ENP but also on any surface

457 structures (i.e. PVP capping agent) and the concentration of ions in the medium. Therefore,
458 the particle size can be larger than that measured by electron microscopy techniques.

459 In this study, some of the raw data from the DLS measurements did not meet the
460 quality criteria (data generally showing high polydispersity). This polydispersity may have
461 been due to the fact that, with the addition of the silver ENPs to seawater, silver chloride
462 precipitated out of solution and caused the ENPs to sediment out. Immediately following the
463 addition of ENPs there was no visible sedimentation of particles in the test exposure
464 solutions, however, after microscopical examination (light microscope) at test termination
465 there was apparent fall out of particles on the bottom of the test wells in all experiments. For
466 zeta-potential measurement salinities above 20 ‰ were unable to provide usable,
467 reproducible values. The high conductivity of the saline samples was most likely the cause of
468 the interference in the readings. In the *C. tenuicorne* media there was a shift in zeta-potential
469 from a negative (CT10 ‰ = -8.68) to a positive (CT20 ‰ = 10.32) value with increasing
470 salinity. This shift of zeta-potential towards a positive value at higher salinities suggests that
471 a possible charge repulsion mechanism is becoming less dominant (Oo and Ong, 2010). It has
472 also been shown that inorganic ions (e.g. Cl⁻) in solution can interact with charged surfaces
473 by non-specific or specific ion adsorption, which will affect the isoelectric point. In some
474 situations, as shown in this study, specific ion adsorption can lead to charge reversal of the
475 surface (Zeta Sizer Nanoseries, 2004).

476 In order to conduct a more thorough risk assessment of nanoparticle exposure it is
477 important to have some specific information. Within the environment their quantity,
478 persistence and dispersion is of vital importance. Without this information it is impossible to
479 conduct a comprehensive risk assessment and herein lies the problem. As previously
480 mentioned, measured total/bulk silver levels in the marine and estuarine environment are very
481 low, 0.04 – 31 ng/L (Kramer et al., 2002) and it is, at present, impossible to distinguish bulk

482 silver released into the marine environment from silver ENPs present. Therefore, we have no
483 true understanding of what the realistic levels of these materials in the aquatic environment
484 are. It is also likely that the exposure levels used in the laboratory are far higher than the
485 natural levels in the environment (Ward et al., 2006). Potentially, these high exposure
486 concentrations, used to cause a toxic effect in some laboratory experiments, may result in
487 different uptake mechanisms and rates than those that would occur at environmentally
488 realistic levels of silver or ENP silver (Fortin and Campbell, 2000). Therefore, making the
489 results of laboratory based ENP studies difficult to incorporate into nanoparticle risk
490 assessment at present, however, studies on the sensitivity and effects of species exposed to
491 ENPs can help provide information that may, in time, lead to a better understanding and more
492 appropriate use of toxicity information for ERA of these materials. In addition, the guidelines
493 need to be clarified to incorporate definitive guidance on the characterisation of ENPs in
494 environmental media. Guidance on the preparation of ENP solutions for ecotoxicological
495 assessment needs to be harmonised in order to yield comparable interlaboratory results for
496 future ERA. The development of additional nano-specific guidance documents are at present
497 being discussed but it remains a complicated and developing area of regulatory
498 ecotoxicology.

499

500 **5 Conclusion**

501 Silver ENPs have the potential to cause toxic effects to marine macrophytes and
502 invertebrates. Given the sensitivity of the two species used in this study, and their widespread
503 distribution in the marine environment, their susceptibility is of utmost importance. In
504 developing methodologies for the assessment of ENP toxicity, it is vital that the most suitable
505 species and endpoints be identified. At present definitive guidelines on the assessment of the
506 risks associated with ENP release and effects in the environment do not exist. Considering

507 that increased salinity was observed to have such a marked effect on the toxicity of the ENPs
508 employed in this study, it may not be appropriate to use freshwater data to extrapolate risk
509 within the marine environment. In addition, as the ISO methods stand, they may not be
510 completely suitable for the assessment of ENPs. Despite the test design, organism's
511 sensitivity and environmental relevance, the methods lack specific information on sample
512 preparation, dosing and quantification of exposure concentrations. Although the
513 characterisation techniques employed were unable to correlate ENP behaviour and salinity it
514 is apparent that some process is causing an increase in toxicity with increasing salinity. As
515 similar increases in toxicity of AgNO₃ with increasing salinity were not observed, it can be
516 concluded that it is not merely the chemical nature of the ENPs responsible for the effect.
517 Therefore, the observed increase in toxicity is most likely due to a combination of the surface
518 properties of the ENPs and their reactivity within highly saline media, and the behaviour of
519 the silver ions and complexation. There is a definite need for a more thorough understanding
520 of the modifying effects of changing environmental parameters. This knowledge, along with
521 suitable guidelines, is required in order to contribute more accurately to an ERA of ENPs
522 introduced into the marine environment.

523

524 **Acknowledgements**

525 The authors sincerely wish to thank Dr. Britta Eklund at the University of Stockholm for
526 providing a starting culture of *C. tenuicorne*, Dr. Norbert Roos at the University of Oslo, for
527 helping with the TEM imaging and Dr. Luke O'Neill for his time and efforts with the AFM
528 imaging. This work was conducted under the framework of the INSPIRE programme, funded
529 by the Irish Government's Programme for Research in Third Level Institutions, Cycle 4,

530 National Development Plan 2007-2013, supported by the European Union Structural Fund.
531 and The Norwegian Research Council under the Yggdrasil mobility programme.

532

533 **6 References**

534

535 ASTM International., 2006 Terminology for nanotechnology. ASTM E2456 – 06.

536

537 Barriada, J.L., Tappin, A.D., Hywel Evans, E., Achterberg, EP., 2007. Dissolved silver
538 measurements in seawater. TrAC 26(8), 809-817.

539

540 Bhatt, I., Tripathi, B.N., 2011. Interaction of engineered nanoparticles with various
541 components of the environment and possible strategies for their risk assessment.
542 Chemosphere. 82(3), 308-317.

543

544 Bilberg, K., Malte, H., Wang, T., Baatrup, E., 2010. Silver nanoparticles and silver nitrate
545 cause respiratory stress in Eurasian perch (*Perca fluviatilis*). Aquat. Toxicol. 96(2), 159-165.

546

547 Canesi, L., Ciacci, C., Vallotto, D., Gallo, G., Marcomini, A., Pojana, G., 2010. *In vitro*
548 effects of suspensions of selected nanoparticles (C60 fullerene, TiO₂, SiO₂) on *Mytilus*
549 hemocytes. Aquat. Toxicol. 96, 151 – 158.

550

551 Cowan, C.E., Jeanne, E.A., Crecelius, E.A., 1985. Silver speciation in seawater: the
552 importance of sulphide and organic complexation. In: Marine and Estuarine Geochemistry.
553 Lewis Publishers, Inc., Chelsea, MI, USA.

554

555 Cowan, C.E., Jeune, E.A., Crecelius, E.A., 1993. Silver speciation in seawater: the
556 importance of sulfide and organic complexation. Proceedings of the First International
557 Conference on Transport, Fate, and Effects of Silver in the Environment, University of
558 Wisconsin-Madison, 15-18.

559

560 Diegoli, S., Manciuola, A.L., Begum, S., Jones, I.P., Lead, J.R., Preece, J.A., 2008.
561 Interaction between manufactured gold nanoparticles and naturally occurring organic
562 macromolecules. *Sci. Total. Environ.* 402, 51 – 61.

563

564 Domingos, R.F., Baalousha, M.A., Ju-Nam, Y., Reid, M.M., Tufenkji, N., Lead, J.R.,
565 Leppard, G.G., Wilkinson, K.J., 2009. Characterising manufactured nanoparticles in the
566 environment: multimethod determination of particles sized. *Environ. Sci. and Technol.* 43
567 (5), 1282 – 1286.

568

569 Eklund, E., 2005. Development of a growth inhibition test with the marine and brackish water
570 red alga *Ceramium tenuicorne*. *Mar. Pollut. Bull.* 50(9), 921 – 930.

571

572 Erikson, R.J., Brooke, L.T., Kahl, M.D., Venter, F.V., Hartling, S.L., Markee, T.P., Spehar,
573 R.L., 1998. Effects of laboratory test conditions on the toxicity of silver to aquatic organisms.
574 Environ. Toxicol. Chem. 17(4), 572 – 578.
575

576 Fabrega, J., Zhang, R., Renshaw, J.C., Liu, W-T., Lead, J.R., 2011. Impact of silver
577 nanoparticles on natural marine biofilm bacteria. Chemosphere. 85(6), 961-966.
578

579 Farkas, J., Christian, P., Gallego Urrea, J.A., Roos, N., Hassellöv, M., Tollefsen, K-E.,
580 Thomas, K.V., 2010. Effects of silver and gold nanoparticles on rainbow trout
581 (*Oncorhynchus mykiss*) hepatocytes. Aquat. Toxicol. 96, 44 – 52.
582

583 Farkas, J., Christian, P., Gallego Urrea, J.A., Roos, N., Hassellöv, M., Tollefsen, K-E.,
584 Thomas, K.V., 2011. Uptake and effects of manufactured silver nanoparticles in rainbow
585 trout (*Oncorhynchus mykiss*) gill cells. Aquat. Toxicol. 101(1), 117 – 125.
586

587 Ferguson, E.A., Hogstrand, C., 1998. Acute silver toxicity to seawater-acclimated rainbow
588 trout: Influence of salinity on toxicity and silver speciation. Environ. Toxicol. Chem. 17(4),
589 589 – 593.
590

591 Fortin, C., Campbell, P.G.C., 2000. Silver uptake by the green alga *Chlamydomonas*
592 *reinhardtii* in relation to chemical speciation: Influence of chloride. Environ. Toxicol. Chem.
593 19, 2769 – 2778.
594

595 Galloway, T., Lewis, C., Dolciotti, I., Johnston, B.D., Moger, J., Regoil, F., 2010. Sublethal
596 toxicity of nano-titanium dioxide and carbon nanotubes in a sediment dwelling marine
597 polychaete. *Environ. Pollut.* 158(5), 1748 – 1755.

598

599 Gebauer, J.S., Treuel, L., 2011. Influence of individual ionic components on the
600 agglomeration kinetics of silver nanoparticles. *J. Colloid. Interface. Sci.* 354(2), 546 – 554.

601

602 Griffitt, R.J., Luo, J., Gao, J., Bonzongo, J.C., Barber, D.S., 2008. Effects of particle
603 composition and species on toxicity of metallic nanomaterials in aquatic organisms. *Environ.*
604 *Toxicol. Chem.* 27, 1972 – 1978.

605

606 Hassellöv, M., Readman, J.W., Ranville, J.F., Tiede, K., 2008. Nanoparticle analysis and
607 characterization methodologies in environmental risk assessment of engineered nanoparticles.
608 *Ecotoxicology.* 17, 344 – 361.

609

610 Handy, R.D., Owen, R., Valsami-Jones, E., 2008. The ecotoxicology of nanoparticles and
611 nanomaterials: current status, knowledge gaps, challenges, and future needs. *Ecotoxicology.*
612 17, 315–325.

613

614 Humes, A.G., 1994. How many copepods? *Hydrobiologia.* 1 – 7, 292 – 293.

615

616 ISO, 14669., 1999. International Organisation for Standardisation. Water quality --
617 Determination of acute lethal toxicity to marine copepods (Copepoda, Crustacea). ISO
618 14669:1999.

619

620 ISO, 10710., 2010. International Organisation for Standardisation. Water quality -- Growth
621 inhibition test with the marine and brackish water macroalga *Ceramium tenuicorne*. ISO
622 10710:2010.

623

624 Ju-Nam, Y., Lead, J.R., 2008. Manufactured nanoparticles: An overview of their chemistry,
625 interactions and potential environmental implications. *Sci. Total. Environ.* 400(1-3), 396-414.

626

627 Kaegi, R., Ulrich, A., Sinnet, B., VonBank, R., Wichser, A., Zuleeg, S., Simmler, H.,
628 Brunner, S., Vonmont, H., Burkhardt, M., Bollner, M., 2008. Synthetic TiO₂ nanoparticle
629 emission from exterior facades into the aquatic environment. *Environ. Pollut.* 156 (2), 233 –
630 239.

631

632 Kato, H., Suzuki, M., Fujita, K., Horie, M., Endoh, S., Yoshida, Y., Iwahashi, H., Takahashi,
633 K., Nakamura, A., Kinugasa, S., 2009. Reliable size determination of nanoparticles using
634 dynamic light scattering method for in vitro toxicology assessment. *Toxicol. in Vitro.* 23 (5),
635 927 – 934.

636

637 Klaine, S.J., Alvarez, P.J.J., Batley, G.E., Fernandes, T.F., Handy, R.D., Lyon, D.Y.,
638 Mahendra, S., McLaughlin, M.J., Lead, J.R., 2008. Nanomaterials in the environment:
639 Behaviour, fate, bioavailability and effects. *Environ. Toxicol. Chemistry*. 27(9), 1825 – 1851.
640

641 Kramer, J.R., Benoit, G., Bowles, K.C., Di Toro, D.M., Herrin, R.T., Luther, G.W. III,
642 Manolopoulos, H., Robillard, K.A., Shafer, M.M., Shaw, J.R., 2002. Environmental
643 chemistry of silver. In: Andren AW, Bober TW. (Eds.) *Silver in the Environment: Transport,*
644 *Fate, and Effects*. SETAC, Pensacola, FL, USA, 1–20.
645

646 Kwok, K.W.H., Leung, K.M.Y., 2005. Toxicity of antifouling biocides to the intertidal
647 harpacticoid copepod *Tigriopus japonicus* (Crustacea, Copepoda): effects of temperature and
648 salinity. *Mar. Pollut. Bull.* 51, 830 – 837.
649

650 Lobban, C.S., Harrison, P.J., 1994. *Seaweed ecology and physiology*. Cambridge University
651 Press, USA. ISBN: 0-521-40334-0, 225–235.
652

653 Luoma, S.N., Ho, Y.B., Bryan, G.W., 1995. Fate, bioavailability and toxicity of silver in
654 estuarine environments. *Mar. Pollut. Bull.* 31(1-3), 44 – 54.
655

656 Macken, A., Giltrap, M., Foley, B., McGovern, E., McHugh, B., Davoren, M., 2008. A model
657 compound study: The ecotoxicological evaluation of five organic contaminants employing a
658 battery of marine bioassays. *Environ. Pollut.* 153(3), 627 – 637.

659

660 Matson, C.W., Auffan, M., Wiesner, M.R., Di Giulio, R.T., 2009. Silver nanoparticle
661 behavior and fish embryotoxicity across a salinity gradient, CEIN and I-CEINT Congress,
662 Washington, DC Matsen(4).

663

664 Meyer, J.N., Lord, C.A., Yang, X.Y., Turner, E.A., Badireddy, A.R., Marinakos, S.M.,
665 Chilkoti, A., Wiesner, M.R., Auffan, M., 2010. Intracellular uptake and associated toxicity of
666 silver nanoparticles in *Caenorhabditis elegans*. *Aquat. Toxicol.* 100, 140 – 150.

667

668 Miao, A-J., Schwehr, K.A., Xu, C., Zhang, S-J., Luo, Z., Quigg, A., Santschi, P.H., 2009.
669 The algal toxicity of silver engineered nanoparticles and detoxification by exopolymeric
670 substances. *Environ. Pollut.* 157, 3034 – 3041.

671

672 Moore, M.N., 2006. Do nanoparticles present ecotoxicological risks for the health of the
673 aquatic environment? *Environ. Internat.* 32(8), 967-976.

674

675 Murr, L.E., Esquivel, E.V., Bang, J.J., de la Rosa, G., Gardea-Torresdey, J.L., 2004.
676 Chemistry and nanoparticulate compositions of a 10 000 year-old ice core melt water. *Water.*
677 *Res.* 38, 4282 – 4296.

678

679 Naddy, R.B., Gorsuch, J.W., Rehner, A.B., McNerney, G.R., Bell, R.A., Kramer, J.R., 2007.
680 Chronic toxicity of silver nitrate to *Ceriodaphnia dubia* and *Daphnia magna*, and potential
681 mitigating factors. *Aquat. Toxicol.* 84(1), 1 – 10.

682

683 Nadella, S.R., Fitzpatrick, J.L., Franklin, N., Bucking, C., Smith, S., Wood, C.M., 2009. To
684 xicity of dissolved Cu, Zn, Ni and Cd to developing embryos of the blue mussel (*Mytilus*
685 *trossulus*) and the protective effect of dissolved organic carbon. *Comp. Biochem. Physiol.*
686 *Part C: Toxicology & Pharmacology.* 149(3), 340 – 348.
687

688 Nair, P.M.G., Park, S.Y., Lee, S-W., Choi, J., 2011. Differential expression of ribosomal
689 protein gene, gonadotrophin releasing hormone gene and Balbiani ring protein gene in silver
690 nanoparticles exposed *Chironomus riparius*. *Aquat. Toxicol.* 101(1), 31 – 37.
691

692 Navarro, E., Baun, A., Behra, R., Hartmann, N.B., Filser, J., Miao, A-J., Quigg, A., Santschi,
693 P.H., Sigg, L., 2008. Environmental behavior and ecotoxicity of engineered nanoparticles to
694 algae, plants, and fungi. *Ecotoxicology* 17(5), 372 – 386.
695

696 Nowack, B., Bucheli, T.D., 2007. Occurrence, behaviour and effects of nanoparticles in the
697 environment. *Environ. Pollut.* 150, 5 – 22.
698

699 Oo, M.H., Ong, S.L., 2010. Implication of zeta potential at different salinities on boron
700 removal by RO membranes. *J. Membr. Sci.* 352(1 - 2), 1 – 6.
701

702 Paterson, G., Macken, A.L., Thomas, K.V., 2011. The need for standardized methods and
703 environmental monitoring programs for anthropogenic nanoparticles. *Anal. Meth.* 3, 1461-
704 1467.
705

706 Pedroso, M.S., Pinho, G.L.L., Rodrigues, S.C., Bianchini, A., 2007. Mechanism of acute
707 silver toxicity in the euryhaline copepod *Acartia tonsa*. *Aquat. Toxicol.* 82, 173 – 180.

708

709 Petersen, E.J., Huang, Q., Weber, W.J., 2008. Ecological uptake and depuration of carbon
710 nanotubes by *Lumbriculus variegatus*. *Environ. Health. Persp.* 116(4), 496-500.

711

712 Petersen, E.J., Henry, T.B., 2012. Methodological considerations for testing the ecotoxicity of
713 carbon nanotubes and fullerenes: Review. *Env. Tox. Chem.* 31(1), 60 – 72.

714

715 Petit, A-N., Eullaffroy, P., Debenest, T., Gagné, F., 2010. Toxicity of PAMAM dendrimers to
716 *Chlamydomonas reinhardtii*. *Aquat. Tox.* 2, 187 – 193.

717

718 Quigg, A., 2008, Trace elements, in: S.E. Jørgensen and B.D. Fath, (Eds), *Ecological*
719 *Stoichiometry in the Encyclopedia of Ecology*, Elsevier, Oxford, pp. 3564–3573.

720

721 Reinfelder, J.R., Chang, S.I., 1999. Speciation and microalgal bioavailability of inorganic
722 silver. *Environ. Sci. Technol.* 33, 1860 – 1863.

723

724 Ringwood, A.H., McCarthy, M., Bates., T.C., Carroll, D.L., 2010. The effects of silver
725 nanoparticles on oyster embryos. *Mar. Environ. Res.* 69(1), 49 – 51.

726

727 Roebben, G., Ramirez-Garcia, S., Hackley, V.A., Roesslein, M., Klaessig, F., Kestens, V.,
728 Lynch, I., Garner, C.M., Rawle, A., Elder, Z.A., McNeil, S., Nel, A., Patri, A., Wick, P.,
729 Wiesner, M., Xia, T., Oberdorster, G., Dawson, K.A. 2011. Interlaboratory comparison of
730 size and surface charge measurements on nanoparticles prior to biological impact assessment.
731 *J. Nanopart. Res.* 13, 2675 – 2687.

732

733 Russel, G., 1985. Recent evolutionary changes in the algae of the Baltic Sea, *Br Phycol J*
734 20:87 – 104.

735

736 Scalf, J., West, P., 2006. Part I: Introduction to Nanoparticle Characterization with AFM.
737 Pacific Nanotechnology, Inc. Santa Clara, CA 95054, USA.

738

739 Tiede, K., Hassellöv, M., Breitbarth, E., Chaudhry, Q., Boxall, A.B.A., 2009. Considerations
740 for environmental fate and ecotoxicity testing to support environmental risk assessments for
741 engineered nanoparticles. *J. Chromatogr. A* 1216(3), 503-509.

742

743 Van Hoecke, K., De Schamphelaere, K.A.C., Van der Meeren, P., Smaghe, G., Janssen,
744 C.R., 2011. Aggregation and ecotoxicity of CeO₂ nanoparticles in synthetic and natural
745 waters with variable pH, organic matter concentration and ionic strength. *Environ. Pollut.*
746 159(4), 970 – 976.

747

748 Verslycke, T., Vangheluwe, M., Heijerick, D., DeSchampelaere, K., VanSprang, P., Janssen,
749 C.R., 2003. The toxicity of metal mixtures to the estuarine mysid *Neomysis integer*
750 (Crustacea: Mysidacea) under changing salinity. *Aquat. Toxicol.* 64, 307–315.

751

752 Vindimian, E., REGTOX-EV6.xls. Regtox 6.3 software program.Xls
753 <http://eric.vindimian.9online.fr>.

754

755 Ward, T.J., Boer, R.L., Hogstrand, C., Kramer, J.R., Lussier, S.M., Stubblefield, W.A.,
756 Wyskiel, D.C., Gorsuch, J.W., 2006. Influence of salinity and organic carbon on the chronic
757 toxicity of silver to mysids (*Americamysis bahia*) and silversides (*Menidia beryllina*).
758 *Environ. Toxicol. Chem.* 25(7), 1809 – 1816.

759

760 Ytreberg, E., Karlsson, J., Ndungu, K., Hasselöv, M., Breitbarth, E., Eklund, B., 2011.
761 Influence of salinity and organic matter on the toxicity of Cu to a brackish water and marine
762 clone of the red macroalga *Ceramium tenuicorne*. *Ecotox. Environ. Saf.* 74(4), 636 – 642.

763

764 Zeta Sizer Nanoseries., 2004. User Manual. Chapter 14. MAN0317, Issue 2.1. July 2004.

765

766 Zuiderveen, J.A., Birge, W. J., 1995. Session 3: Interaction of silver and metal chelators on
767 *Ceriodaphnia dubia* survival and reproduction, in: Andren, Anders W.; Bober, Thomas W.
768 (Eds.) / *The 3rd international conference proceedings: transport, fate and effects of silver in*
769 *environment* (August 1995)

770

771 **Figures**


772

773

774

775 **Figure 1** An example of AFM image of Ag-NP_{PVP} in CT10 ‰ with underlying topographic
776 profile.

777




778

779 **Figure 2** TEM images (scalebar: 200 nm) of Ag-NP_{PVP} in NSW (a - b), CT10 % (c - d),

780 CT20 % (e), CT30 % (g - h) and the corresponding average size in all media (f).

781

782




783

784 **Figure 3** Particle size distribution (PSD) from TEM image analysis of Ag-NP_{PVP} in different
 785 media (n = 100 for measurements in all media).


786

787 (a)



788

789 (b)



790


791

792

793 **Figure 4** Toxicity of (a) Ag-NP_{PVP} (b) AgNO₃ to *Tisbe battagliai* exposed for ■ 24 hours and
794 □ 48 hours.* indicates statistical significance from the control ($p \leq 0.05$). Data are mean \pm SD
795 (n = 3). CV for the controls ranged from 0.00 – 2.94 %.


796

797 (a)



798

799 (b)



800

801

802 **Figure 5** Growth inhibition of *Ceramium tenuicorne* after 7 days exposure to Ag-NP_{PVP} (a)
803 and AgNO₃ (b) at 10 ‰ (□), 20 ‰ (■) and 30 ‰ (▣). * indicates statistical significance from
804 the control ($p \leq 0.05$). Data are mean \pm SD (n = 3). CV for the control ranged from 0.15 –
805 1.33 % (10 ‰), 5.29 – 13.07 % (20 ‰) and 15.7 – 16.65 % (30 ‰)
806

807 **Table 1** Particle size analysis and characterisation results for Ag-NP_{PVP} in all test media.

Media type	Method	Concentration (mg/L)	Mean PS (nm)	SD (nm)	Min (nm)	Max (nm)
MilliQ water	DLS	0.1	56.9	7.8	-	-
	TEM	10	7.0*	-	1.0*	60.0*
	AFM	10	18.5	6.2	5.9	38.2
10 % 2A Media	DLS	0.1	149.0	29.8	-	-
	TEM	10	16.0	5.8	7.9	33.7
	AFM	10	24.7	38.7	6.5	280.1
20 % 2A Media	DLS	0.1	57.3	3.2	-	-
	TEM	10	20.2	4.1	11.4	36.3
	AFM	10	13.8	7.1	5.5	56.4
30 % 2A Media	DLS	0.1	105.0	18.5	-	-
	TEM	10	19.3	6.3	9.6	39.6
	AFM	10	24.0	23.7	5.7	120.3
NSW	DLS	0.1	76.4	39.2	-	-
	TEM	10	18.9	5.2	9.9	34.9
	AFM	10	12.2	7.8	55.4	71.7

808 SD = Standard deviation, NSW = Natural Seawater, DLS = Dynamic Light Scattering, TEM = Transmission Electron Microscopy, AFM =
809 Atomic Force Microscopy

810 * Approximately 48 % of particles were < 5nm, while, 30 % were between 5 – 10 nm in size (Farkas et al., 2010)

811

812 **Table 2** Ecotoxicity data for *Tisbe battagliai* and *Ceramium tenuicorne* assayed with silver nanoparticles (Ag-NP_{PVP}) and silver nitrate (AgNO₃).

Test Substance	Test Species	Salinity (‰)	Duration	EC ₁₀ /LC ₁₀ ^a (µg/L) ^a	EC ₅₀ /LC ₅₀ ^b (µg/L) ^b	NOEC ^c /LOEC ^d (µg/L)
Ag-NP _{PVP}	<i>T. battagliai</i>	34	24 h	3.8 (2.4 – 5.6)	30.8 (26.2 – 35.7)	3.2/5.6
		34	48 h	0.9 (0.6 – 1.7)	7.9 (6.2 – 10.3)	3.2/5.6
Ag-NP _{PVP}	<i>C. tenuicorne</i>	10	7 d	32.5 (4.0 – 129.6)	2120.4 (1199.9 – 3316.8)	100/500
Ag-NP _{PVP}	<i>C. tenuicorne</i>	20	7 d	1.5 (0.1 – 11.0)	373.9 (179.8 – 782.7)	50/100
Ag-NP _{PVP}	<i>C. tenuicorne</i>	30	7 d	0.1 (0.007 – 1.4)	26.7 (12.6 – 69.8)	5/10
AgNO ₃	<i>T. battagliai</i>	34	24 h	12.7 (0.8 – 258.7)	167.3 (52.8 – 481.8)	32/56
		34	48 h	8.7 (2.0 – 102.5)	90.9 (38.0 – 238.9)	10/32
AgNO ₃	<i>C. tenuicorne</i>	10	7 d	42.2 (6.4 – 143.8)	3606.5 (2318.2– 5378.6)	10/50
AgNO ₃	<i>C. tenuicorne</i>	20	7 d	10.9 (0.6 – 81.4)	2246.6 (1020.8 – 4056.4)	50/100
AgNO ₃	<i>C. tenuicorne</i>	30	7 d	11.4 (0.4 – 105.6)	2312.2 (1038.4 – 4819.6)	500/1000

813 ^aEC₁₀/LC₁₀ values and corresponding 95 % confidence intervals

814 ^bEC₅₀/LC₅₀ values and corresponding 95 % confidence intervals

815 ^cNOEC, no observed effect concentration, the highest observed concentration at which no significant effect ($p \leq 0.05$) was detected

816 ^dLOEC, lowest observed effect concentration, the lowest concentration of the tested concentration at which a significant ($p \leq 0.05$) effect was detected

817