

2009

## Application of Natural Antimicrobials for Food Preservation

Brijesh Tiwari

*University College Dublin*

Vasilis Valdramidis

*Technological University Dublin*

Colm O' Donnell

*University College Dublin*

*See next page for additional authors*

Follow this and additional works at: <https://arrow.tudublin.ie/schfsehart>



Part of the [Food Microbiology Commons](#)

---

### Recommended Citation

Cullen, P.J. et al. (2009) Application of Natural Antimicrobials for Food Preservation, *J. Agric. Food Chem.*, 2009, 57 (14), pp 5987–6000 DOI: 10.1021/jf900668n

This Article is brought to you for free and open access by the School of Food Science and Environmental Health at ARROW@TU Dublin. It has been accepted for inclusion in Articles by an authorized administrator of ARROW@TU Dublin. For more information, please contact [arrow.admin@tudublin.ie](mailto:arrow.admin@tudublin.ie), [aisling.coyne@tudublin.ie](mailto:aisling.coyne@tudublin.ie), [vera.kilshaw@tudublin.ie](mailto:vera.kilshaw@tudublin.ie).

Funder: National Development Plan, Dept of Agriculture, Food & Fisheries

---

**Authors**

Brijesh Tiwari, Vasilis Valdramidis, Colm O' Donnell, Kasiviswanathan Muthukumarappan, Patrick Cullen, and Paula Bourke

## Application of Natural Antimicrobials for Food Preservation

BRIJESH K. TIWARI,<sup>†</sup> VASILIS P. VALDRAMIDIS,<sup>§</sup> COLM P. O' DONNELL,<sup>†</sup>  
 KASIVISWANATHAN MUTHUKUMARAPPAN,<sup>†,‡</sup> PAULA BOURKE,<sup>§</sup> AND P. J. CULLEN<sup>\*,§</sup>

<sup>†</sup>Biosystems Engineering, School of Agriculture, Food Science and Veterinary Medicine,  
 University College Dublin, Belfield, Dublin 4, Ireland, <sup>‡</sup>Agricultural and Biosystems Engineering  
 Department, South Dakota State University, Brookings, South Dakota, and <sup>§</sup>School of Food Science &  
 Environmental Health, Dublin Institute of Technology, Dublin 1, Ireland

In this review, antimicrobials from a range of plant, animal, and microbial sources are reviewed along with their potential applications in food systems. Chemical and biochemical antimicrobial compounds derived from these natural sources and their activity against a range of pathogenic and spoilage microorganisms pertinent to food, together with their effects on food organoleptic properties, are outlined. Factors influencing the antimicrobial activity of such agents are discussed including extraction methods, molecular weight, and agent origin. These issues are considered in conjunction with the latest developments in the quantification of the minimum inhibitory (and noninhibitory) concentration of antimicrobials and/or their components. Natural antimicrobials can be used alone or in combination with other novel preservation technologies to facilitate the replacement of traditional approaches. Research priorities and future trends focusing on the impact of product formulation, intrinsic product parameters, and extrinsic storage parameters on the design of efficient food preservation systems are also presented.

**KEYWORDS:** Antimicrobial activity; chemical compounds; plant/animal/microbial antimicrobials mechanism; minimum inhibitory concentration

### INTRODUCTION

A number of nontraditional preservation techniques are being developed to satisfy consumer demand with regard to nutritional and sensory aspects of foods. Generally, foods are thermally processed by subjecting them to temperatures varying from 60 to 100 °C for the duration of a few seconds to a minute in order to destroy vegetative microorganisms. During this period of treatment, a large amount of energy is transferred to the food. However, this energy can trigger unwanted reactions, leading to undesirable organoleptic and nutritional effects (1). Ensuring food safety and at the same time meeting such demands for retention of nutrition and quality attributes has resulted in increased interest in alternative preservation techniques for inactivating microorganisms and enzymes in foods. Quality attributes of importance include flavor, odor, color, texture, and nutritional value. This increasing demand has opened new dimensions for the use of natural preservatives derived from plants, animals, or microflora. In biopreservation, storage life is extended, and/or safety of food products is enhanced by using natural or controlled microflora, mainly lactic acid bacteria (LAB) and/or their antibacterial products such as lactic acid, bacteriocins, and others (2). Typical examples of investigated compounds are lactoperoxidase (milk), lysozyme (egg white, figs), saponins and flavonoids (herbs and spices), bacteriocins (LAB), and chitosan (shrimp shells) (3). Antimicrobial compounds present in foods can

extend the shelf life of unprocessed or processed foods by reducing the microbial growth rate or viability (4). Originally, spices and herbs were added to change or to improve taste. Some of these substances are also known to contribute to the self-defense of plants against infectious organisms (5, 6).


Extensive research has investigated the potential application of natural antimicrobial agents in food preservation. In this review, antimicrobials and their chemical and biochemical components from a range of natural sources and their applications in food systems are reviewed. Natural antimicrobials in food preservation can be used alone or in combination with other nonthermal technologies. Naturally derived antimicrobial systems from plant, animal, and microbial origin are detailed, and the latest developments in the quantification of the minimum (and noninhibitory) concentration of antimicrobials and/or their components are presented.

### PLANT ORIGIN ANTIMICROBIAL AGENTS

Edible, medicinal, and herbal plants and their derived essential oils (EO) (and their hydrosols, i.e., byproducts of an essential oil purification procedure) and isolated compounds contain a large number of secondary metabolites that are known to retard or inhibit the growth of bacteria, yeast, and molds (7, 8). Many of these compounds are under investigation and are not yet exploited commercially. The antimicrobial compounds in plant materials are commonly found in the essential oil fraction of leaves (rosemary, sage, basil, oregano, thyme, and marjoram), flowers or buds (clove), bulbs (garlic and onion), seeds (caraway,

\*Corresponding author. Phone: +353 14027595. Fax: +35314024495. E-mail: pjculen@dit.ie.

## Scheme 1. Plant Origin Antimicrobial Agents



fennel, nutmeg, and parsley), rhizomes (asafetida), fruits (pepper and cardamom), or other parts of plants (9, 10). Plant EOs and their constituents have been widely used as flavoring agents in foods since the earliest recorded history, and it is well established that many have a wide spectra of antimicrobial action (11–15). These compounds may be lethal to microbial cells or they might inhibit the production of secondary metabolites (e.g., mycotoxins) (16). Plant essential oils are generally more inhibitory against Gram-positive than Gram-negative bacteria (10, 17, 18). While this is true for many EOs, there are some agents that are effective against both groups, such as oregano, clove, cinnamon, and citral (19–21). The major EO components with antimicrobial effects found in plants, herbs, and spices are phenolic compounds, terpenes, aliphatic alcohols, aldehydes, ketones, acids, and iso-flavonoids (8, 22–27). Chemical analysis of a range of EOs revealed that the principal constituents of many include carvacrol, thymol, citral, eugenol (see Scheme 1 for their chemical structure), and their precursors (8, 28–30). It has been reported that some nonphenolic constituents of EOs are more effective or quite effective against Gram-negative bacteria, e.g., allyl isothiocyanate (AIT) (31) and garlic oil (32), respectively. In addition, AIT is also effective against many fungi (33). Generally, the antimicrobial efficacy of EOs is dependent on the chemical structure of their components as well as the concentration. Many of the antimicrobial compounds present in plants can be part of their pre- or postinfectious defense mechanisms for combating infectious or parasitic agents (34). Consequently, plants that manifest relatively high levels of antimicrobial action may be sources of compounds that inhibit the growth of foodborne pathogens (35). Compounds are also generated in response to stress from inactive precursors (36), which may be activated by enzymes, hydrolases or oxidases, usually present in plant tissues (37). In mustard and horse radish, precursor glucosinolates are converted by enzyme myrosinase to yield a variety of isothiocyanates including the allyl form, which is a strong antimicrobial agent (38).

The application of plant EOs for controlling the growth of foodborne pathogens and food spoilage bacteria requires evaluation of the range of activity against the organisms of concern to a particular product, as well as effects on a food's organoleptic properties. Plant EOs are usually mixtures of several components. Oils with high levels of eugenol (allspice, clove bud and leaf, bay, and cinnamon leaf), cinnamamic aldehyde (cinnamon bark and cassia oil), and citral (lemon myrtle, *Litsea cubeba*, and lime) are usually strong antimicrobials (39, 40). The EOs from *Thymus* spp. possess significant quantities of phenolic monoterpenes and have reported antiviral (41), antibacterial (42, 43), and antifungal (44, 45) properties. The volatile terpenes carvacrol, *p*-cymene,  $\gamma$ -terpinene, and thymol contribute to the antimicrobial activity of oregano, thyme, and savory (18). The antimicrobial activity of sage and rosemary can be attributed to borneol and other phenolic compounds in the terpene fraction. Davidson and Naidu (40) reported that the terpene thejone was responsible

for the antimicrobial activity of sage, whereas in rosemary, a group of terpenes (borneol, camphor, 1,8 cineole, *a*-pinene, camphone, verbenonone, and bornyl acetate) was responsible. Plant EOs such as cumin, caraway, and coriander have inhibitory effects on organisms such as *Aeromonas hydrophila*, *Pseudomonas fluorescens*, and *Staphylococcus aureus* (46, 47), marjoram and basil have high activity against *B. cereus*, *Enterobacter aerogenes*, *Escherichia coli*, and *Salmonella*, and lemon balm and sage EOs appear to have adequate activity against *L. monocytogenes* and *S. aureus* (10). Gutierrez et al. (10) showed that oregano and thyme EOs had comparatively high activity against enterobacteria (minimum inhibitory concentration (MIC) of oregano and thyme at a range of 190 ppm and 440 ppm, respectively, for *E. cloacae*), lactic acid bacteria (MIC of oregano and thyme at a range of 55 ppm and 440 ppm, respectively, for *Lactobacillus brevis*), *B. cereus* (MIC of oregano and thyme at a range of 425 ppm and 745 ppm, respectively, for *Lactobacillus brevis*), and *Pseudomonas* spp (MIC of oregano and thyme at a range of 1500 ppm for *P. putida*), although in general *Pseudomonas* species are consistently highly resistant to plant antimicrobials (10, 48). One of the attributed factors can be the production of exopolysaccharide layers forming biofilms of the microorganism that can delay penetration of the antimicrobial agent (49).

Lee et al. (50) investigated the antibacterial activity of vegetables and juices and concluded that green tea and garlic extracts have broad applications as antibacterial agents against a wide range of pathogens. Arrowroot tea extract has reported antimicrobial activity against *E. coli* O157:H7 (19). Ibrahim et al. (35) reported the potential of caffeine at a concentration of 0.5% or higher as an effective antimicrobial agent for the inactivation of *E. coli* O157:H7 in a liquid system (i.e., brain heart infusion (BHI)).

**Mechanisms of Antimicrobial Action.** The possible modes of action for phenolic compounds (EO fractions) as antimicrobial agents have been previously reviewed (16, 24, 27, 36, 51–53). However, the exact mechanism of action is not clear. The effect of phenolic compounds can be concentration dependent (54). At low concentration, phenols affect enzyme activity, particularly those associated with energy production, while at high concentrations, they cause protein denaturation. The antimicrobial effect of phenolic compounds may be due to their ability to alter microbial cell permeability, thereby permitting the loss of macromolecules from the interior (for example ribose and Na glutamate) (55). They could also interfere with membrane function (electron transport, nutrient uptake, protein, nucleic acid synthesis, and enzyme activity) (55) and interact with membrane proteins, causing deformation in structure and functionality (56–58). The high antibacterial activity of phenolic components can be further explained in terms of alkyl substitution into the phenol nucleus (25). The formation of phenoxyl radicals that interact with alkyl substituents does not occur with more stable molecules such as the ethers myristicin or anethole, which was related to the relative lack of antimicrobial activity of fennel, nutmeg, or parsley EOs (10).

Delaquis and Mazza (38) reported that the antimicrobial activity of isothiocyanates derived from onion and garlic is related to the inactivation of extracellular enzymes through oxidative cleavage of disulfide bonds and that the formation of the reactive thiocyanate radical was proposed to mediate the antimicrobial effect. Carvacrol, (+)-carvone, thymol, and *trans*-cinnamaldehyde are reported to decrease the intracellular ATP (adenosine triphosphate) content of *E. coli* O157:H7 cells while simultaneously increasing extracellular ATP, indicating the disruptive action of these compounds on the plasma membrane (59). Inactivation of yeasts can be attributed to the disturbance of several enzymatic systems, such as energy production and structural component synthesis (60).

193 **Factors Affecting Antimicrobial Activity.** Antimicrobial activity  
194 of EOs is influenced by a number of factors including botanical  
195 source, time of harvesting, stage of development, and method of  
196 extraction (61). For example, Chorianopoulos et al. (62) reported  
197 that *Satureja* EOs obtained during the flowering period were the  
198 most potent with bactericidal properties. The composition, struc-  
199 ture as well as functional groups of the oils play an important role  
200 in determining their antimicrobial activity. Usually compounds  
201 with phenolic groups are the most effective (5, 25). Most studies  
202 related to the antimicrobial efficacy of EOs have been conducted  
203 in vitro using microbiological media (63–71). Consequently,  
204 there is less understanding related to their efficacy when applied  
205 to complex food systems. Key areas requiring further knowledge  
206 for optimized application of natural antimicrobials in food  
207 include targeting the microorganism of concern, the intelligent  
208 use of combinations to provide a synergy of activity, matching the  
209 activity of the compounds to the composition, and processing and  
210 storage conditions of the food (9, 72).

211 Plant EOs of thyme, clove, and pimento were tested against  
212 *Listeria monocytogenes* and were found to be highly effective in  
213 peptone water. However, when the EOs were applied in a food  
214 system, Singh et al. (73) concluded that efficacy of EOs was reduced  
215 due to interaction with food components. In general, higher  
216 concentrations of EOs are required in foods than in laboratory  
217 media. Combinations of EOs could minimize the application  
218 concentrations required, thereby reducing any adverse organolep-  
219 tical impact; however, their application for microbial control may  
220 also be affected by food composition (74). The antimicrobial  
221 efficacy of EOs was found to be a function of ingredient manipula-  
222 tion, for example, the antimicrobial activity of thyme is increased in  
223 high protein concentrations, concentrations of sugars above 5% on  
224 the microbial growth medium did not reduce EO efficacy, and high  
225 potato starch concentrations decreased the EO antimicrobial  
226 activity of oregano and thyme on *L. monocytogenes* in food model  
227 systems (74, 75). Finally, low pH values (of the range of 5) seemed to  
228 have the highest impact on the increase of the antimicrobial effect of  
229 EOs on *L. monocytogenes* (74). Low pH values appear to increase  
230 the hydrophobicity of EOs, consequently enabling easier dissolu-  
231 tion in the lipids of the cell membrane of target bacteria (54).

232 Accordingly, the challenge for practical application of EOs is  
233 to develop optimized low dose combinations to maintain product  
234 safety and shelf life, thereby minimizing the undesirable flavor  
235 and sensory changes associated with the addition of high con-  
236 centrations of EOs.

## 237 ANIMAL ORIGIN ANTIMICROBIAL AGENTS

238 There are numerous antimicrobial systems of animal origin,  
239 where they have often evolved as host defense mechanisms.  
240 Lysozyme is a bacteriolytic enzyme, commercially sourced from  
241 hen's egg white which is reported to inhibit the outgrowth of  
242 *Clostridium tyrobutyricum* spores in semihard cheeses (76). Lyso-  
243 zyme has found commercial applications; inovapure is said to be  
244 effective against a wide range of food spoilage organisms and can  
245 be successfully used to extend the shelf life of various food  
246 products, including raw and processed meats, cheese, and other  
247 dairy products. The lactoperoxidase system, which is naturally  
248 active in milk, has strong antimicrobial effects against both  
249 bacteria and fungi. A wide range of both Gram-negative (77)  
250 and Gram-positive bacteria (78) are inhibited by the lactoperox-  
251 idase system. However, studies have shown that Gram-negative  
252 bacteria were generally found to be more sensitive to lactoperox-  
253 idase mediated food preservation than Gram-positive species  
254 (79, 80). Many of the antimicrobial agents inherent to animals  
255 are in the form of antimicrobial peptides (polypeptides).

Antimicrobial peptides were first isolated from natural sources  
in the 1950s when nisin was isolated from lactic acid bacteria for  
potential application as a food preservative (81). Subsequently,  
antimicrobial peptides were isolated from other natural sources,  
such as plants, insects, amphibians, crustaceans, and marine  
organisms (82–84). Antimicrobial peptides (AMPs) are widely  
distributed in nature and are used by many if not all life forms as  
essential components of nonspecific host defense systems. The list  
of discovered AMPs has been constantly increasing, with much  
discovery in the last two decades. The list of AMPs produced by  
animal cells includes magainin (85), MSI-78 (86), PR-39 (87),  
spheniscin (88), pleurocidin (89), dermaseptin S4 (90), K4S4-  
(1-14) (91), cecropin P1 (92), melittin (93), LL-37 (94), clavanin  
A (92), and curvacin A (95). Antimicrobial peptides present a  
promising solution to the problem of antibiotic resistance because,  
unlike traditional antimicrobial agents, specific molecular sites are  
not targeted, and their characteristic rapid destruction of mem-  
branes does not allow sufficient time for even fast-growing  
bacteria to mutate. Some of the potential antimicrobials of animal  
origin which could be used as food additives are discussed below.

276 **Pleurocidin.** Pleurocidin, a 25 amino acid peptide isolated from  
277 the skin mucus membrane of the winter flounder (*Pleuronectes*  
278 *americanus*) is active against Gram-positive and Gram-negative  
279 bacteria. It is heat-stable, salt-tolerant, and insensitive to physio-  
280 logical concentrations of magnesium and calcium (96). Pleuroci-  
281 din has potential for use in food applications and was found to be  
282 effective against foodborne organisms including *Vibrio parahe-*  
283 *molyticus*, *L. monocytogenes*, *E. coli* O157:H7, *Saccharomyces*  
284 *cerevisiae*, and *Penicillium expansum* (97). The antimicrobial  
285 activity of pleurocidin against foodborne microorganisms was  
286 reported at levels well below the legal limit for nisin (10,000 IU/g)  
287 without significant effect on human red blood cells (97), thereby  
288 indicating its potential as a food preservative and a natural  
289 alternative to conventional chemicals. However, pleurocidin  
290 was inhibited by magnesium and calcium (96), which may limit  
291 the use of this AMP in environments rich in these cations.

292 **Defensins.** Defensins are another group of antimicrobial pep-  
293 tides widely found in nature including mammalian epithelial cells  
294 of chickens, turkeys, etc. They are abundant in cells and tissues  
295 active in host defense against microorganisms (98, 99). They are  
296 reported to have a broad spectrum of antimicrobial activity (100),  
297 including Gram-positive, Gram-negative bacteria, fungi, and  
298 enveloped viruses (101, 102).

299 **Lactoferrin.** Bovine and activated lactoferrin (ALF), an iron-  
300 binding glycoprotein present in milk, has antimicrobial activity  
301 against a wide range of Gram-positive and negative bacteria (102)  
302 fungi, and parasites (103). Lactoferrin has been applied in meat  
303 products (104–106) as it has recently received approval for  
304 application on beef in the USA (USDA-FSIS 2008. FSIS Direc-  
305 tive 7120.1 Amendment 15).

306 **Other AMPs.** Protamine, like salmine and clupeine, has been  
307 reported to be isolated from fish and is found to be effective against  
308 Gram-negative and Gram-positive bacteria, yeasts, and molds  
309 (108–111). Magainin peptides isolated from frogs (112) have been  
310 found effective against a range of food-related pathogens (113),  
311 implying a possible application as food preservatives (91, 114, 115).

312 **Chitosan.** Chitosan, a natural biopolymer obtained from the  
313 exoskeletons of crustaceans and arthropods, is known for its  
314 unique polycationic nature and has been used as active material  
315 for its antifungal activity (72, 116) and antibacterial activity (117–  
316 120). Liu et al. (121) studied the efficacy of chitosan against *E. coli*  
317 and concluded that low molecular weight chitosan is effective for  
318 controlling growth. The strong antibacterial activity of chitosan  
319 was also observed against *S. aureus*, while its molecular weight  
320 appeared to be a significant parameter defining its activity (122).

**Lipids.** Like lipids of plant origin, lipids of animal origin have antimicrobial activity against a wide range of microorganisms. Free fatty acids at mucosal surfaces have been shown to inactivate *S. aureus* (123). Milk lipids have recorded activity for inactivation of Gram-positive bacteria including *S. aureus*, *Cl. botulinum*, *B. subtilis*, *B. cereus*, *L. monocytogenes*, Gram-negative bacteria such as *P. aeruginosa*, *E. coli*, and *Salmonella enteritidis* (124–126), and also against various fungi such as *Aspergillus niger*, *Saccharomyces cerevisiae*, and *C. albicans* (36, 124). Lipids may serve to inhibit the proliferation as well as the prevention of the establishment of pathogenic or spoilage microorganisms in food matrices.

Shin et al. (127) studied eicosapentaenoic acid (EPA) and docosahexaenoic acid (DHA), which are formed in animal (including fish and shellfish) tissues but not plant tissues (18:3  $\omega$ -3). DHA is a component of membrane structural lipids that are enriched in certain phospholipid components of the retina and nonmyelin membranes of the nervous system in animals. Bioconverted EPA and DHA exhibited antibacterial activities against four Gram-positive bacteria, *B. subtilis*, *L. monocytogenes*, *Staphylococcus aureus* ATCC 6538, *S. aureus* KCTC 1916, and seven Gram-negative bacteria, *E. aerogenes*, *E. coli*, *E. coli* O157:H7, *E. coli* O157:H7 (human), *P. aeruginosa*, *Salmonella enteritidis*, and *S. typhimurium* (127). The growth inhibition by both EPA and DHA was similar against Gram-positive bacteria, while the bioconverted extract of DHA was more effective than EPA against Gram-negative bacteria.

**Mechanism of Antimicrobial Action.** The mechanism of action of AMPs seems to involve multiple targets. The plasma membrane is the most cited target; however, recent studies suggest intracellular targets at least for some peptides (128, 129). Although most AMPs act by nonspecific mechanisms, they often display some selectivity between different microorganisms, for example, Gram-negative compared with Gram-positive bacteria (130, 131) and susceptibility of fungal cells compared with other eukaryotic cells (132). Antimicrobial peptides can assume amphipathic structures, which are able to interact directly with the microbial cell membrane, rapidly disrupting the membrane in several locations, resulting in leaching out of vital cell components (96, 133). Previous studies conducted on the mechanism of action of pleurocidin revealed that it exhibits strong membrane translocation and pore-formation ability, reacting with both neutral and acidic anionic phospholipid membranes (134). Lipids inactivate microorganisms mainly by disruption of bacterial cell wall or membrane, inhibition of intracellular replication, or inhibition of an intracellular target (135). Monoacylglycerols lower the heat resistance of certain bacteria and fungi; therefore, they may find application in reducing the required heat treatment for certain foods (36). Lysozyme hydrolyses the  $\beta$ -1,4-glycosidic linkage in sugar polymers such as *N*-acetylmuramic acid and *N*-acetylglucosamine linkages found in bacterial peptidoglycan (136).

#### MICROBIAL ORIGIN ANTIMICROBIAL AGENTS

Bacteria produce many compounds that are active against other bacteria, which can be harnessed to inhibit the growth of potential spoilage or pathogenic microorganisms. These include fermentation end products such as organic acids, hydrogen peroxide, and diacetyl, in addition to bacteriocins and other antagonistic compounds such as reuterin (137). Both Gram-negative and Gram-positive bacteria produce bacteriocins. Bacteriocins are proteinaceous antibacterial compounds, which constitute a heterologous subgroup of ribosomally synthesized antimicrobial peptides (138). Bacteriocin production can be exploited by food processors to provide an additional barrier to undesirable bacterial growth in foods (Table 1).

Bacteriocins are cationic peptides that display hydrophobic or amphiphilic properties, and in most cases, the target for their activity is the bacterial membrane. Depending on the producer organism and classification criteria, bacteriocins can be categorized into several groups (139–142) with as many as five classes of bacteriocins proposed (143–145). The majority fall into classes I and II, which are the most intensively researched to date. The class I group, termed lantibiotics, are small peptides that are characterized by their content of several unusual amino acids (146). The class II bacteriocins are small, nonmodified, heat stable peptides (147). Another classification is with respect to the producing microorganism and is specifically named after the genus, species, or the group of microorganisms, e.g., lantibiotics for bacteriocins of lactic acid bacteria, colicins of *E. coli*, klebsins of *Klebsiella pneumoniae* (148). A large number of bacteriocins have been isolated and characterized from lactic acid bacteria, and some have acquired a status as potential food preservatives because of their antagonistic effect on important pathogens. Many bacteriocins are active against food borne pathogens and spoilage bacteria (149–152). The important ones include nisin, diplococcin, acidophilin, bulgarican, helveticin, lactacin, and plantaricin (153). Nisin is produced by various *Lactococcus lactis* strains, is the most thoroughly studied bacteriocin to date, and is applied as an additive in food worldwide (154). While the antimicrobial polypeptide nisin and related compounds such as pediocin are the only bacteriocins widely used for food preservation (155, 156), many other bacteriocins have been reported and have shown potential for food preservation and safety applications.

**Reuterin.** Reuterin ( $\beta$ -hydroxypropionaldehyde) is a water-soluble nonproteinaceous metabolite of glycerol (157). It is a broad spectrum antimicrobial compound produced by some strains of *Lactobacillus reuteri*, with recorded activity against Gram-negative and Gram-positive bacteria, yeasts, and filamentous fungi (158). Reuterin was isolated, purified, and identified by Talarico and Dobrogosz (159) and is active over a wide range of pH values and resistant to the action of proteolytic and lipolytic enzymes (160). Reuterin is reported to exhibit bacteriostatic activity against *Listeria monocytogenes* but was only slightly bactericidal against *Staphylococcus aureus* at 37 °C. However, higher bactericidal activity was reported against *E. coli* O157:H7, *S. choleraesuis* subsp. *Choleraesuis*, *Y. enterocolitica*, *A. hydrophila* subsp. *Hydrophila*, and *C. jejuni* (161).

**Pediocin.** Pediocin is produced by strains of *Pediococcus acidilactici* and *P. pentosaceus* and is designated generally recognized as a safe (GRAS). The organism is commonly isolated from and used in fermented sausage production. The bacteriocins produced by *P. acidilactici* are AcH, PA-1, JD, and 5, and those produced from *P. pentosaceus* are A, N5p, ST18, and PD1 (162). Most pediocins are thermostable proteins and function over a wide range of pH values. Pediocin AcH has proven efficacy against both spoilage and pathogenic organisms, including *L. monocytogenes*, *Enterococcus faecalis*, *S. aureus*, and *Cl. Perfringens* (163). Natamycin is an antifungal produced by *Streptomyces natalensis* that is effective against nearly all molds and yeasts but has little or no effect on bacteria.

**Nisin.** Nisin is the most widely used bacteriocin. To date, nisin is the only natural antimicrobial peptide (see Scheme 2 for its structure) approved by the FDA for use as a food preservative; however, it has a limited spectrum of activity, does not inhibit Gram-negative bacteria or fungi, and is only effective at low pH (164, 165). Nisin is produced by fermentation of a modified milk medium by certain strains of lactic acid bacterium, *Lactococcus lactis*. Nisin functions by interacting with the phospholipids

**Table 1.** Effect of Natural Antimicrobial Agents on Food Preservation and Quality<sup>a</sup>

food product	antimicrobial agent (concentrations)	microbial dynamics	quality attributes	reference
fruit yoghurt	vanillin (2000 ppm)	yeast, bacterial (delays growth)	shelf life (†)	(232)
tomato juice	clove oil (0.1%)	total plate count (3.9LR)	shelf life (†), vitamin C (~)	(208)
	mint extract (1.0%)	total plate count (8.34LR)		
	nisin (0.004%)	total plate count (↓)		
ready-to-eat fruit salad	citral (25–125 ppm)	yeasts and lactic acid bacteria (LAB) (delays growth)	shelf life (†)	(233)
	citron (300–900 ppm)			
	citron (600 ppm)	<i>Salmonella enteritidis</i> E4 (2 LR), <i>Escherichia coli</i> 555 (<4.5 LR) <i>Listeria monocytogenes</i> Scott A (4 LR)	sensory characteristics (~)	
raspberries	methyl jasmonate (MJ), allyl isothiocyanate (AITC) EO of <i>Melaleuca alternifolia</i> (tea tree oil)		AC (†) AC (↓) AC (†)	(234)
fresh cut water melon	nisin (25 µg/mL)	<i>L. monocytogenes</i> (0.8 LR)	quality (†)	(235)
lettuce	thyme oil (1 mL/l)	<i>E. coli</i> (6.32LR)		(236)
baby carrot		<i>E. coli</i> (5.57LR)		
minimally processed carrots	oregano oil (250 ppm)	background spoilage microflora total viable count (TVC) (>1 LR) lactic acid bacteria (LAB) (>1 LR) <i>Pseudomonas</i> (<1 LR)	sensory characteristics (~)	(205)
minimally processed vegetables	thyme oil (1%)	<i>Aeromonas spp</i> (2 LR)	sensory properties (↓), shelf life (†)	(237)
		psychrotrophic aerobic plate count (4.19 LR) plate count agar (5.44 LR)		
wine	nisin	LAB (minimum inhibitory concentration, MIC = 0.39 mg/mL) <i>Oenococcus oeni</i> (MIC 0.01 mg/mL) acetic acid bacteria (MIC 1.5 mg/mL)		(238)
milk	reuterin (8 AU/ml)	<i>L. monocytogenes</i> (4.59 LR)		(161)
	nisin (100 IU/ml)	<i>S. aureus</i> counts (5.45 LR)		
skimmed milk powder	nisin (100 IU/ml)	<i>L. innocua</i> (3.8 LR)		(240)
chicken meat	nisin	<i>E. coli</i> (<1 LR)	proximate composition (~), shelf life (†)	(209)
		<i>Brochothrix thermosphacta</i> (~) <i>Lactobacillus alimentarius</i> (~) <i>Brochothrix thermosphacta</i> (~) <i>Lactobacillus alimentarius</i> (delays growth)		
fish	EOs (0.5% carvacrol + 0.5% thymol)	TVC (2.5LR)	shelf life (†), lipid oxidation (↓) sensory characteristics (~)	(241)
red meat	tea catechins (300 mg/kg)		shelf life (†), lipid oxidation (↓)	(242)
beef hot dog	clove oil (5 mL/l)	<i>L. monocytogenes</i> (1.15–1.71LR)		(73)
	thyme oil (1 mL/l)	<i>L. monocytogenes</i> (0.67–1.05 LR)		
pork bologna	nisin (125 µg/mL)	<i>L. monocytogenes</i> (1.5LR)		(169)
minced beef	<i>Capsicum annum</i> extract	<i>Salmonella typhimurium</i> (Minimum lethal concentration, MLC 15 g/kg) <i>Pseudomonas aeruginosa</i> (MLC 30 g/kg)		(199)
chicken frankfurter	clove oil (1% v/w)	<i>L. monocytogenes</i> (4.5 LR)		(197)
cooked beef	grape seed extract (1%)	<i>Escherichia coli</i> (1.7 LR) <i>S. Typhimurium</i> (2.0 LR) <i>L. monocytogenes</i> (0.8 LR) <i>Aeromonas hydrophila</i> (0.4 LR)	color (~), lipid oxidation (↓)	(200)


<sup>a</sup> AU: arbitrary units were defined as the reciprocal of the highest two-fold dilution that did not allow the growth of the indicator strain. AC: anthocyanin content. † and ↓ indicate increase and decrease, respectively, while ~ shows no significant difference. LR: microbial log reduction.

449 in the cytoplasmic membrane of bacteria, thus disrupting  
450 membrane function and preventing outgrowth of spores by  
451 inhibiting the swelling process of germination. It is highly  
452 active against many of the Gram-positive bacteria and speci-  
453 fically used by the cheese industry to control the growth of  
454 *Clostridium spp.* (166). Substantial research has evaluated the  
455 efficacy of nisin against various pathogens and its use for  
456 different food products (167–174). Nisin has been used to  
457 inhibit microbial growth in beef (173), sausages (2), liquid  
458 whole egg (174), ground beef (175), and poultry (176). It has  
459 also been reported to reduce initial levels of *Listeria mono-*  
460 *cytogenes* and suppress subsequent growth in ready-to-eat  
461 (RTE) meat products (177, 178). Komitopoulou et al. (179)  
462 reported that nisin could be used for the effective control of

*Alicyclobacillus acidoterrestris* in fruit juices. A nisin level of 463  
464 6.25 µg/g could inhibit lactic acid bacteria (LAB) growth for  
465 over 28 days and for 35 days with 25 µg/g (180). The effects of  
466 three types of phosphate (used as emulsifiers) on nisin activity  
467 in sausage were compared, and LAB growth rate was fastest in  
468 samples containing orthophosphate and slowest in sausages  
469 containing diphosphate.

**Mechanism of Antimicrobial Action.** The antimicrobial action 470  
471 of bacteriocins is based on pore formation in the cytoplasmic  
472 membrane of the target microorganism. This leads to a loss of  
473 small intracellular molecules and ions and a collapse of the proton  
474 motive force (181). Nisin is less effective on Gram-negative  
475 bacteria, as the outer membrane disables the entry of this  
476 molecule to the site of action (50, 119, 182, 183). The first step

Scheme 2. Structure of Nisin



477 in the mode of action of nisin is to pass through the cell wall of  
 478 Gram-positive bacteria. Generally, it is assumed that nisin  
 479 passes the cell wall by diffusion. However, the Gram-positive  
 480 cell wall can act as a molecular sieve against nisin depending  
 481 on its composition, thickness, or hydrophobicity (184). The  
 482 removal of the cell wall from nisin-resistant *Listeria* resulted in  
 483 the removal of nisin resistance, suggesting that the cell wall plays  
 484 a role in the differences in susceptibility toward nisin (185). The  
 485 next step of the antimicrobial process of nisin is to associate with  
 486 the cytoplasmic membrane of the target microorganism. It has  
 487 been suggested that nisin interacts electrostatically with the  
 488 negatively charged phosphate groups of surface membrane  
 489 phospholipids (173).

490 **Factors Affecting Antimicrobial Activity.** Various factors can  
 491 impact the antimicrobial efficacy of bacteriocins. These include  
 492 the emergence of bacteriocin-resistant bacteria, conditions that  
 493 destabilize the biological activity of proteins such as proteases or  
 494 oxidation processes, binding to food components such as fat  
 495 particles or protein surfaces, inactivation by other additives, poor  
 496 solubility, and uneven distribution in the food matrix and/or pH  
 497 effects on bacteriocin stability and activity (137). The application  
 498 of bacteriocins in combination with other preservation hurdles  
 499 has been proposed to reduce the selection for resistance to  
 500 bacteriocins in target strains and/or to extend its inhibitory  
 501 activity to Gram-negative species (182). Interactions between  
 502 bacteriocin and the food matrix may result in a decrease in the  
 503 efficacy of the bacteriocin. The combination of bacteriocins with  
 504 other minimal or nonthermal preservation technologies may  
 505 prove useful for practical applications. This approach is of value  
 506 for the control of Gram-negative bacteria as their outer mem-  
 507 brane acts as an efficient barrier against hydrophobic solutes and  
 508 macromolecules, such as bacteriocins (119).

#### 509 QUANTIFICATION OF THE MINIMUM AND NONINHIBITORY 510 CONCENTRATION

511 The use of antimicrobials as preservatives in food systems  
 512 can be constrained when effective antimicrobial doses exceed  
 513 organoleptic acceptable levels (especially for essential oils) or  
 514 when they are added to complex food systems. Two specific  
 515 concentrations appear to be of interest, i.e., the noninhibitory  
 516 concentration, NIC, the concentration above which the inhibi-  
 517 tor begins to have a negative effect on growth, and the  
 518 minimum inhibitory concentration, MIC, which marks the  
 519 concentration above which no growth is observed by compar-  
 520 ison with the control (186). Therefore, these concentrations are  
 521 quantified with the aim of defining the boundaries of sensory  
 522 acceptability and antimicrobial efficacy of antimicrobials (26).  
 523 Most of the studies on the calculation of MIC and NIC are  
 524 semiquantitative, while quantitative approaches have been  
 525 mainly applied on studies concerning the antimicrobial activity  
 526 of plant origin antimicrobial agents, i.e., essential oils and their  
 527 components.

528 The MIC and NIC are dependent on experimental conditions.  
 529 The influencing conditions include the incubation temperature,  
 530 organism, and inoculum size, and therefore, they should be  
 531 reported in studies where MIC and NIC are evaluated (187,  
 532 188). In vitro studies for identifying the MIC can be divided into  
 533 groups such as diffusion, dilutions, impedance, and optical  
 534 density (or absorbance) methods (see for e.g., refs (189–191)).  
 535 Most of these evaluations are based on an end-point approach for  
 536 evaluating the MIC, i.e., end result in which no growth is obtained  
 537 for a test level of preservative, into which an inoculum of  
 538 microbes is added. This kind of approach is considered semi-  
 539 quantitative (188).

540 Lambert and Pearson (188) examined the inhibitory activity of  
 541 single compounds of EOs and developed a fully quantitative  
 542 approach. This is given by the Lambert–Pearson model (LPM)  
 543 inspired by a modified Gompertz equation (eq 1) to evaluate the  
 544 dose–responses of microorganisms against several inhibitors.  
 545 This modeling approach has already been examined for optical  
 546 density, O.D. (187, 188), and impedance microbial measure-  
 547 ments (62).

$$548 \quad fa = \exp \left[ - \left( \frac{x}{P_1} \right)^{P_2} \right] \quad (1)$$

549 In eq 1, *fa* is the fractional area which is defined as the ratio of  
 550 inhibited growth to uninhibited growth as measured by the  
 551 applied method (impedance, optical density, etc.), *x* is the  
 552 inhibitor concentration (mg/L), *P*<sub>1</sub> is the concentration at max-  
 553 imum slope (of a log *x* vs *fa* plot; see Figure 1 for a graphical  
 554 example of this equation), and *P*<sub>2</sub> is a slope parameter. Observe  
 555 that *fa* can be measured by using the trapezoidal rule under the O.  
 556 D. (or other microbial measurements)/time curves and then  
 557 taking the ratio of the test area to that of the control (187).  
 558 Therefore, the range of *fa* will be between 0 and 1 (Figure 1). The  
 559 routine, trapz, provided by Matlab is an example of a software  
 560 package that can be used for performing a trapezoidal numerical  
 561 integration.


562 The MIC (eq 2) and the NIC (eq 3) can then be calculated as the  
 563 intercept of the concentration axis to the tangent at the maximum  
 564 gradient of the *fa*/log concentration curve and the intercept of the  
 565 tangent at the maximum gradient of the *fa*/log concentration  
 566 curve to the *fa* = 1 contour.

$$567 \quad MIC = P_1 \cdot \exp \left( \frac{1}{P_2} \right) \quad (2)$$

$$568 \quad NIC = P_1 \cdot \exp \left( \frac{1-e}{P_2} \right) \quad (3)$$

569 Guillier et al. (192) developed another approach for evaluating  
 570 the MIC based on the use of growth rate models. After estimation  
 571 of the maximum specific growth rates ( $\mu_{max}$ ) from optical density  
 572 measurements, the MIC was defined as the concentration at which  
 573  $\mu_{max}$  was reduced by 50% compared to the control.





**Figure 1.** Hypothetical inhibition profile as can be described by eq 1 for increasing values of  $P_2$  and constant  $P_1$  (left panel) and increasing values of  $P_1$  and constant  $P_2$  (right panel). Inhibitor concentration is expressed on a logarithmic scale.

570 growth kinetics by a modified Gompertz model, they assessed the  
571 antimicrobial concentration dependence on  $\mu_{\max}$  (eq 4).

$$\sqrt{\mu_{\max}} = \sqrt{\mu_{\max}(c=0) \cdot f(c)} \quad (4)$$

572  $f(c)$  can be described either as eq 5, i.e., the  $SR_{\mu}$  model, or as eq 6,  
573 i.e., the  $LP_{\mu}$  model.

$$f(c) = \left(1 - \frac{c}{MIC}\right)^{\beta}, c < MIC \text{ or } 0, c \geq MIC \quad (5)$$

$$f(c) = \exp \left[ - \left( \frac{c}{MIC / \exp \left( \frac{\ln(NIC/MIC)}{-e} \right)} \right)^{-e / (\ln(NIC/MIC))} \right] \quad (6)$$

574  $\mu_{\max}(c=0)$  is the growth rate in the absence of the antimicrobial  
575 ( $c=0$ ) and  $\beta$  a shape parameter representing the sensitivity of the  
576 microorganism to an antimicrobial in eq 5. These two approaches  
577 appeared to give equivalent results. Observe that for estimating  
578 the parameters of MIC, NIC, and  $\mu_{\max}(c=0)$  of eq 6, a regression  
579 is performed for the data that relate the maximum specific growth  
580 rates ( $\mu_{\max}$ ) with the concentration of the inhibitor.

581 Lambert et al. (26) argued that the majority of antimicrobial  
582 activity could be attributed to two components acting independ-  
583 dently. Therefore, they also suggested another expression for a  
584 mixture of two inhibitors that could be extended in case there are  
585 more inhibitors as presented in eq 7:

$$fa_{x_1, \dots, x_k} = \exp \left\{ - \left[ \left( \frac{x_1}{C_{i,1}} \right)^{C_{i,2}} + \dots + \left( \frac{x_k}{C_{k,1}} \right)^{C_{k,2}} \right]^{C_Q} \right\} \quad (7)$$

586 where parameters  $C_{i,1}$  are the concentrations of the  $x_i$  inhibitors at  
587 the maximum slope. The main difference is that the current  
588 expression takes into account interactions between the antimi-  
589 crobials, which means that it could be considered for any additive,  
590 antagonistic, and synergistic activity between the studied inhibi-  
591 tors. For an example in which a mixture of two antimicrobials is  
592 studied reference is made to Lambert and Lambert (187). In that  
593 case, the MIC of any of the  $x_i$  antimicrobials is then given by eq 8.

$$MIC = C_{i,1} \cdot \exp \left( \frac{1}{C_{i,2} + C_Q} \right) \quad (8)$$

595 Another interesting quantitative approach for evaluating the  
596 bactericidal effect of different agents has been suggested by Lui  
597 et al. (193). This is based on a concentration killing curve  
598 approach and the estimation of the so-called median bactericidal  
599 concentration and bactericidal intensity. The developed method

is based on the correlation (by the use of a sigmoidal curve with an  
inflection point) of the population size (number CFU per plate)  
with respect to the concentration of the agent. This approach has  
been applied for quantifying the bactericidal potency of anti-  
biotics against *E. coli* and might have to be further investigated  
for different antimicrobials. Similar to the discussed approaches,  
novel modeling methods for quantitatively expressing the effect  
of antimicrobials through MIC and NIC values can be developed  
by knowledge coming from predictive microbiology. An overview  
of representative cases for different modeling expressions tackling  
the effect of both chemical and natural inhibitory compounds can  
be found in Devlieghere et al. (194).

Accurate quantitative evaluations of MIC and NIC are  
important for designing effective preservation methods that  
are based on the use of the discussed antimicrobials. These  
quantitative methods can be exploited to give insight to optimal  
concentrations or combinations for real food systems by direct  
comparison of the antimicrobial efficacy of different antimicro-  
bials, their individual or combined components, or their mix-  
tures, and for efficient design of preservation for food products  
based on the principles of hurdle technology. These approaches  
have not received much attention for evaluating the MIC or the  
minimum bactericidal concentration of the antimicrobials of  
animal and microbial origin, but their potential is evident.

## APPLICATIONS OF NATURAL ANTIMICROBIALS IN FOOD

The extrapolation of results obtained from in vitro experiments  
with laboratory media to food products is not straightforward as  
foods are complex, multicomponent systems consisting of differ-  
ent interconnecting microenvironments. Though there is vast  
potential for natural antimicrobial agents in food preservation,  
most of the literature presents inactivation data from model foods  
or laboratory media. **Table 1** reports inactivation studies in real  
food systems. The level of natural preservatives required for  
sufficient efficacy in food products in comparison with laboratory  
media may be considerably higher, which may negatively impact  
the organoleptic properties of food.

Monoacylglycerols have increased the shelf life of various  
foods including soy sauce, miso, sausages, cakes, and noodles (36).  
The lauric acid ester of monoacylglycerol has reported antimi-  
crobial potential in seafood salads and various flesh foods  
including deboned chicken meat, minced fish, refrigerated beef  
roasts, and frankfurter slurries (126, 195). Hao et al. (196) studied  
the efficacy of a range of plant extracts for inhibition of  
*A. hydrophila* and *L. monocytogenes* in refrigerated cooked  
poultry and found that eugenol reduced pathogen counts by  
4 log<sub>10</sub> cfu/g over a 14 day storage trial. Similarly, 1–2% w/w clove  
oil inhibited the growth of a range of *Listeria* spp. in chicken  
frankfurters over 2 weeks at 5 °C (197). Conversely, Shekarforoush

648 et al. (198) found that EOs of oregano and nutmeg were effective  
649 against *E. coli* O157:H7 in a broth system but had no effect in  
650 ready-to-cook chicken. Careaga et al. (199) recorded that 1.5 mL/  
651 100 g of capsicum extract was sufficient to prevent the growth of  
652 *S. typhimurium* in raw beef but that 3 mL/100 g was required for a  
653 bactericidal effect against *P. aeruginosa*. Ahn et al. (200) also found  
654 a range of plant extracts to be useful for reduction of pathogens  
655 associated with cooked beef and quality maintenance; however,  
656 Uhart et al. (201) concluded that when in direct contact, spices  
657 inactivated *S. typhimurium* DT104 but that the activity decreased  
658 considerably when added to a complex food system such as ground  
659 beef. Gutierrez et al. (74, 75) concluded that plant essential oils are  
660 more effective against food-borne pathogens and spoilage bacteria  
661 when applied to ready-to-use foods containing a high protein level  
662 at acidic pH as well as lower levels of fats or carbohydrates and  
663 moderate levels of simple sugars. The success of plant derived  
664 antimicrobials when applied to fruit and vegetable products is also  
665 documented in the literature. Karapinar et al. (202) recommended  
666 unripe grape juice as an alternative antimicrobial agent for  
667 enhancing the safety of salad vegetables, and Martinez-Romero  
668 et al. (203) suggested that carvacrol could be applied as a novel tool  
669 for the control of fungal decay on grapes. Although Valero and  
670 Frances (204) found that low concentrations of carvacrol, cinnamaldehyde, or thymol had a clear antibacterial effect against  
671 *B. cereus* in carrot broth, cinnamaldehyde retained a significant  
672 activity at storage temperatures of 12 °C. Gutierrez et al. (205)  
673 found that the efficacy of oregano EO was comparable with  
674 chlorine as a decontamination treatment for ready-to-eat carrots.  
675 Use of this essential oil contributed to the acceptability of sensory  
676 quality and appreciation. A novel application of plant extracts is  
677 for the production of chocolate; Kotzekidou et al. (206) reported  
678 enhanced inhibitory effects of plant extracts against an *E. coli*  
679 cocktail at 20 °C.

681 Antimicrobials from microbial sources, especially nisin, find  
682 application in a number of foods such as milk, orange juice (207),  
683 and tomato juice (208), and for increasing the shelf life of chicken  
684 meat without altering sensory properties of the product (209). The  
685 efficacy of enterocin AS-48 for inhibition of *B. cereus* in rice and  
686 *S. aureus* in vegetable sauces was investigated (210, 211) with  
687 bacteriocin levels in the range of 20–35 µg/mL and 80 µg/mL,  
688 respectively.

689 Investigation of the antimicrobial properties of preservatives  
690 from animal sources and their possible potential in food applica-  
691 tion is still in its infancy, with few published studies available as  
692 described above. A common conclusion that could be drawn  
693 from these studies is the fact that the significant potential of  
694 antimicrobials from animal sources is not being exploited.

695 Some other applications in foods that got attention in previous  
696 years are the use of bioactive packaging technologies. These  
697 systems can be applied for all of the discussed antimicrobials,  
698 i.e., plant, animal, and microbial origin agents either by adding a  
699 sachet (or possibly by encapsulating the agents (212)) into the  
700 package, dispersing bioactive agents in the packaging, coating  
701 bioactive agents on the surface of the packaging material, or  
702 utilizing antimicrobial macromolecules with film-forming prop-  
703 erties or edible matrixes (213, 214). Film-coating applications  
704 have been reported for meat, fish, poultry, bread, cheese, fruits,  
705 and vegetables (215).

#### 706 USE OF NATURAL ANTIMICROBIALS IN THE MULTIPLE- 707 HURDLE CONCEPT

708 Investigations based on combinations of natural antimicro-  
709 bials with other nonthermal processing technologies within the  
710 multiple-hurdle concept are warranted to counteract any poten-  
711 tial organoleptic or textural effects on food products as well as

712 optimizing microbial inactivation. The preservative action of  
713 bacteriocins alone in a food system is unlikely to ensure compre-  
714 hensive safety. This is of particular significance with regard to  
715 Gram-negative pathogenic bacteria that are protected from the  
716 antimicrobial action of bacteriocins by the presence of an outer  
717 membrane. When the outer membrane is disrupted by agents  
718 such as the food grade chelating agent ethylene diamine tetra-  
719 acetate (EDTA), which acts by binding to Mg<sup>2+</sup> ions in lipopo-  
720 lysaccharide, the outer membrane of Gram-negative bacteria are  
721 rendered sensitive to the antimicrobial action of bacterio-  
722 cins (181). Potential synergistic effects may be found with other  
723 chemical or physical inactivation technologies including dense  
724 phase carbon dioxide, ultrasound, pulsed-electric field, high  
725 pressure, and ozone treatment. As a consequence of applying  
726 these nonthermal methods, bacterial cell membranes can weaken  
727 or become susceptible to additional antimicrobial agents such as  
728 bacteriocins, causing lethality. The use of bacteriocins in combi-  
729 nation with organic acids or other antimicrobials can similarly  
730 result in enhanced inactivation (216). Studies reporting the  
731 effective use of nisin against Gram-negative organisms and fungi  
732 are those in which nisin was used in combination with traditional  
733 food preservatives such as organic acids and chelating  
734 agents (217). Rajkovic et al. (218) found that the activity of nisin  
735 combined with carvacrol was enhanced in a potato puree by  
736 comparison with BHI broth and that more obvious effects against  
737 *B. cereus* and *B. circulans* were observed at higher temperatures.  
738 The application of bacteriocins in combination with treatments  
739 that could enhance their effectiveness in foods requires investiga-  
740 tion. Examples of the synergistic effects that can be obtained  
741 using mild traditional preservation techniques in conjunction  
742 with novel food processing technologies are better studied in  
743 vitro but require further investigation in food products to ensure  
744 successful practical application. The antibacterial activity of  
745 inhibitory compounds, such as nisin, enterocin, monolaurin,  
746 and the lactoperoxidase system (LPS), can be enhanced if applied  
747 in combination (219–221), with chelating agents (182, 222, 223) or  
748 with preservative treatments such as high hydrostatic pressure,  
749 pulsed electric field, low pH, or freeze/thaw cycles (224–228). The  
750 combination of plant EOs with modified atmosphere packaging  
751 for control of spoilage species was reported by Skandamis and  
752 Nychas (229) and Matan et al., (230). Seydim and Sarikus (231)  
753 also investigated the use of EOs in an active packaging system  
754 based on an edible whey protein film and concluded that oregano  
755 was the most effective EO against a range of food pathogens.  
756 Allyl isothiocyanate was successfully applied to chopped, refrig-  
757 erated, nitrogen packed beef for the control of *E. coli* at levels in  
758 excess of 1000 ppm.

759 **Conclusions and Future Trends.** Interest in natural antimicro-  
760 bials has expanded in recent years in response to consumer  
761 demand for greener additives. During the last two decades,  
762 natural preservatives have been investigated for practical applica-  
763 tions. These technologies have been shown to inactivate micro-  
764 organisms and enzymes without significant adverse effects on  
765 organoleptic or nutritional properties. Reported studies have  
766 demonstrated that natural antimicrobial agents described in this  
767 review may offer unique advantages for food processing. In  
768 addition to improving the shelf life and safety of foods, natural  
769 antimicrobial agents may allow novel food products with en-  
770 hanced quality and nutritional properties to be introduced to the  
771 market.

772 The applications of natural antimicrobial agents are likely to  
773 grow steadily in the future because of greater consumer demands  
774 for minimally processed foods and those containing naturally  
775 derived preservation ingredients. More complex considerations  
776 arise for combinations of technologies, particularly with respect

777 to optimization of practical applications. Intelligent selection of  
778 appropriate systems based on detailed, sequential studies and  
779 quantitative approaches to evaluate the efficiency of antimicro-  
780 bials is necessary. The impact of product formulation, extrinsic  
781 storage parameters, and intrinsic product parameters on the  
782 efficacy of novel applications of combined nonthermal systems  
783 requires further study.

#### 784 ABBREVIATIONS USED

785 Abu, amino butyric acid; Ala, alanine; asn, asparagine; Dha,  
786 dehydroalanine; Dhb, dehydrobutyrine ( $\beta$ -methyldehydroala-  
787 nine); Gly, glycine; His, histidine; Ile, isoleucine; Leu, leucine ;  
788 Lys, lysine ; Met, methionine ; Pro, proline; Ser, serine; Val, valine.

#### 789 LITERATURE CITED

- 790 (1) Barbosa-Canovas, G. V.; Pothakamury, U. H.; Palou, E.; Swanson,  
791 B. G. *Nonthermal Preservation of Foods*; Marcel Dekker: New York,  
792 1997; p 304.
- 793 (2) Hugas, M.; Garriga, M.; Aymerich, M. T.; Monfort, J. M. Inhibition  
794 of *Listeria* in dry fermented sausages by the bacteriocinogeni  
795 *Lactobacillus sake* CTC494. *J. Appl. Bacteriol.* **1995**, *79* (3), 322–330.
- 796 (3) Devlieghere, F.; Vermeulen, A.; Debevere, J. Chitosan: antimicro-  
797 bial activity, interactions with food components and applicability as  
798 a coating on fruit and vegetables. *Food Microbiol.* **2004**, *21* (6),  
799 703–714.
- 800 (4) Beuchat, L. R.; Golden, D. A. Antimicrobials occurring naturally in  
801 foods. *Food Technol.* **1989**, *43* (1), 134–142.
- 802 (5) Deans, S. G.; Noble, R. C.; Hiltunen, R.; Wuryani, W.; Penzes, L. G.  
803 Antimicrobial and antioxidant properties of *Syzygium aromaticum*  
804 (L.) Merr. & Perry: impact upon bacteria, fungi and fatty acid levels in  
805 ageing mice. *Flavour Frag. J.* **1995**, *10*, 323–328.
- 806 (6) Kim, H. Y.; Lee, Y. J.; Hong, K.-H.; Kwon, Y.-K.; Sim, K.-C.; Lee,  
807 J.-Y.; Cho, H.-Y.; Kim, I.-S.; Han, S.-B.; Lee, C.-W.; Shin, I.-S.;  
808 Cho, J. S. Isolation of antimicrobial substances from natural  
809 products and their preservative effects. *Food Sci. Biotechnol.* **2001**,  
810 *10* (1), 59–71.
- 811 (7) Burt, S. A.; Reinders, R. D. Antibacterial activity of selected plant  
812 essential oils against *Escherichia coli* O157:H7. *Lett. Appl. Microbiol.*  
813 **2003**, *36*, 162–167.
- 814 (8) Chorianopoulos, N. G.; Giaouris, E. D.; Skandamis, P. N.;  
815 Haroutounian, S. A.; Nychas, G. J. E. Disinfectant test against  
816 monoculture and mixed-culture biofilms composed of technological,  
817 spoilage and pathogenic bacteria: bactericidal effect of essential oil  
818 and hydrosol of *Satureja thymbra* and comparison with standard acid-  
819 base sanitizers. *J. Appl. Microbiol.* **2008**, *104*, 1586–1869.
- 820 (9) Nychas, G. J. E.; Skandamis, P. N. Chapter 9: Antimicrobials from  
821 Herbs and Spices. In *Natural Anti-Microbials for the Minimal*  
822 *Processing of Foods*; Roller, S., Ed.; Woodhead Publishing: Cambridge,  
823 U.K., 2003; pp 176–200.
- 824 (10) Gutierrez, J.; Rodriguez, G.; Barry-Ryan, C.; Bourke, P. Efficacy of  
825 plant essential oils against foodborne pathogens and spoilage  
826 bacteria associated with ready-to-eat vegetables: Antimicrobial  
827 and sensory screening. *J. Food Prot.* **2008**, *71* (9), 1846–1854.
- 828 (11) Lis-Balchin, M.; Deans, S. G. Bioactivity of selected plant essential  
829 oils against *Listeria monocytogenes*. *J. Appl. Microbiol.* **1997**, *82*,  
830 759–762.
- 831 (12) Smith-Palmer, A.; Stewart, J.; Fyfe, L. Antimicrobial properties of  
832 plant essential oils and essences against five important foodborne  
833 pathogens. *Lett. Appl. Microbiol.* **1998**, *26*, 118–122.
- 834 (13) Kim, J.; Marshall, M. R.; Wei, C. Antimicrobial activity of some  
835 essential oil components against five food borne pathogens. *J. Agric.*  
836 *Food Chem.* **1995**, *43*, 2839–2845.
- 837 (14) Packiyasothy, E. V.; Kyle, S. Antimicrobial properties of some herb  
838 essential oils. *Food Aust.* **2002**, *54* (9), 384–387.
- 839 (15) Alzoreky, N. S.; Nakahara, K. Antimicrobial activity of extracts  
840 from some edible plants commonly consumed in Asia. *Int. J. Food*  
841 *Microbiol.* **2002**, *80*, 223–230.
- 842 (16) Davidson, P. M. Chemical Preservatives and Naturally Antimicro-  
843 bial Compounds. In *Food Microbiology: Fundamentals and Frontiers*,  
2nd ed.; Doyle, M. P., Beuchat, L. R., Montville, T. J., Eds.; ASM Press: 844  
Washington, DC, 2001; pp 593–628. 845
- (17) Marino, M.; Bersani, C.; Comi, G. Impedance measurement to study 846  
antimicrobial activity of essential oils from Lamiaceae and Compo- 847  
sitae. *Int. J. Food Microbiol.* **2002**, *67*, 187–195. 848
- (18) Chorianopoulos, N.; Kalpoutzakis, E.; Aligiannis, N.; Mitaku, S.; 849  
Nychas, G. J.; Haroutounian, S. A. Essential oils of *Satureja*, 850  
*Origanum*, and *Thymus* species: Chemical composition and anti- 851  
bacterial activities against foodborne pathogens. *J. Agric. Food* 852  
*Chem.* **2004**, *52*, 8261–8267. 853
- (19) Kim, S.; Fung, D. Y. Antibacterial effect of water soluble arrowroot 854  
(*Puerariae radix*) tea extract on foodborne pathogens in ground beef and 855  
mushroom soup. *J. Food Prot.* **2004**, *67*, 1953–1956. 856
- (20) Sivropoulou, A.; Papanikolaou, E.; Nikolaou, C.; Kokkini, S.; 857  
Lanaras, T.; Arsenakis, M. Antimicrobial and cytotoxic activities 858  
of *Origanum* essential oils. *J. Agric. Food Chem.* **1996**, *44*, 1202–1205. 859
- (21) Skandamis, P.; Tsigarida, E.; Nychas, G.-J.E. The effect of oregano 860  
essential oil on survival/death of *Salmonella typhimurium* in meat 861  
stored at 5 °C under aerobic, VP/MAP conditions. *Food Microbiol.* 862  
**2002**, *19*, 97–103. 863
- (22) Katayama, T.; Nagai, I. Chemical significance of the volatile 864  
components of spices in the food preservative view point. VI. 865  
Structure and antibacterial activity of Terpenes. *Bull. Jpn. Soc.* 866  
*Sci. Fish.* **1960**, *26*, 29–32. 867
- (23) Farag, R. S.; Daw, Z. Y.; Hewed, F. M.; El-Baroty, G. S. A. 868  
Antimicrobial activity of some Egyptian spice oils. *J. Food Prot.* 869  
**1989**, *52*, 665–667. 870
- (24) Nychas, G. J. E. Natural Antimicrobials from Plants. In *New* 871  
*Methods of Food Preservation*; Gould, G. W., Ed.; Blackie Academic 872  
& Professional: Glasgow, Scotland, 1995; pp 58–89. 873
- (25) Dorman, H. J. D.; Deans, S. G. Antimicrobial agents from plants: 874  
antibacterial activity of plant volatile oils. *J. Appl. Microbiol.* **2000**, 875  
*88*, 308–316. 876
- (26) Lambert, R. J. W.; Skandamis, P. N.; Coote, P. J.; Nychas, G.-J. E. 877  
A study of the minimum inhibitory concentration and mode of 878  
action of oregano essential oil, thymol and carvacrol. *J. Appl.* 879  
*Microbiol.* **2001**, *91*, 453–462. 880
- (27) Lopez-Malo, A.; Alzamora, S. M.; Palou, E. Naturally Occurring 881  
Compounds: Plant Sources. In *Antimicrobials in Food*, 3rd ed.; 882  
Davidson, P. M., Sofos, J. N., Branen, A. L., Eds.; CRC Press: New 883  
York, 2005; pp 429–251. 884
- (28) Juliano, C.; Mattana, A.; Usai, M. Composition and in vitro 885  
antimicrobial activity of the essential oil of *Thymus herba-borona* 886  
Loisel growing wild in Sardinia. *J. Essent. Oil Res.* **2000**, *12*, 516–522. 887
- (29) Demetzos, C.; Perdetzoglou, D. K. Composition and antimicrobial 888  
studies of the oils of *Origanum calcaratum* Juss and *O. scabrum* Boiss 889  
from Greece. *J. Essent. Oil Res.* **2001**, *13*, 460–462. 890
- (30) Shelef, L. A. Antimicrobial effects of spices. *J. Food Saf.* **1983**, *6*, 29–44. 891
- (31) Ward, S. M.; Delaquis, P. J.; Holley, R. A.; Mazza, G. Inhibition 892  
of spoilage and pathogenic bacteria on agar and pre-cooked roasted 893  
beef by volatile horseradish distillates. *Food Res. Int.* **1998**, *31*, 19–26. 894
- (32) Yin, M. C.; Cheng, W. S. Antioxidant and antimicrobial effects of 895  
four garlic-derived organosulfur compounds in ground beef. *Meat* 896  
*Sci.* **2003**, *63*, 23–28. 897
- (33) Nielsen, P. V.; Rios, R. Inhibition of fungal growth on bread by 898  
volatile components from spices and herbs, and their possible 899  
application in active packaging, with special emphasis on mustard 900  
essential oil. *Int. J. Food Microbiol.* **2000**, *60*, 219–229. 901
- (34) Rauha, J. P.; Remes, S.; Heinonen, M.; Hopia, A.; Kahkonen, M.; 902  
Kujala, T.; Pihlaja, K.; Vuorela, H.; Vuorela, P. Antimicrobial 903  
effects of Finnish plant extracts containing flavonoids and other 904  
phenolic compounds. *Int. J. Food Microbiol.* **2000**, *56*, 3–12. 905
- (35) Ibrahim, S. A.; Salameh, M. M.; Phetsomphou, S.; Yang, H.; Seo, C. 906  
W. Application of caffeine, 1,3,7-trimethylxanthine, to control 907  
*Escherichia coli* O157:H7. *Food Chem.* **2006**, *99*, 645–650. 908
- (36) Sofos, J.; Beuchat, L. R.; Davidson, P. M.; Johnson, E. A. *Naturally* 909  
*Occurring Antimicrobials in Food. Task Force Report*; Council of 910  
Agricultural Science and Technology: Ames, IA, 1998; p 103. 911
- (37) Holley, R. A.; Patel, D. Improvement in shelf-life and safety of 912  
perishable foods by plant essential oils and smoke antimicrobials. 913  
*Food Microbiol.* **2005**, *22*, 273–292. 914

- 915 (38) Delaquis, P. J.; Mazza, G. Antimicrobial properties of isothiocya- 986  
 916 nates in food preservation. *Food Technol.* **1995**, *49* (11), 73–84. 987
- 917 (39) Lis-Balchin, M.; Deans, S. G.; Eaglesham, E. Relationship between 988  
 918 bioactivity and chemical composition of commercial essential oils. 989  
 919 *Flavour Frag. J.* **1998**, *13*, 98–104. 990
- 920 (40) Davidson, P. M.; Naidu, A. S. Phyto-Phenols. In *Natural Food* 991  
 921 *Antimicrobial Systems*; Naidu, A. S., Ed.; CRC Press: Boca Raton, FL, 992  
 922 2000; pp 265–294. 993
- 923 (41) Wild, R. *The complete book of natural and medical cures*; Rodale Press, 994  
 924 Inc.: Emmaus, PA, 1994. 995
- 925 (42) Essawi, T.; Srour, M. Screening of some Palestinian medicinal 996  
 926 plants for antibacterial activity. *J. Ethnopharmacol.* **2000**, *70*, 343– 997  
 927 349. 998
- 928 (43) Cosentino, S.; Tuberoso, C. I. G.; Pisano, B.; Satta, M.; Mascia, V.; 999  
 929 Arzedi, E. In-vitro antimicrobial activity and chemical composition 1000  
 930 of Sardinian Thymus essential oils. *Lett. Appl. Microbiol.* **1999**, *29*, 1001  
 931 130–135. 1002
- 932 (44) Pina-Vaz, C.; Gonc, A.; Rodrigues, A.; Pinto, E.; Costa-de-Oliveira, 1003  
 933 S.; Tavares, C.; Salgueiro, L. Antifungal activity of Thymus oils and 1004  
 934 their major compounds. *J. Eur. Acad. Dermatol. Venereol.* **2004**, *18*, 1005  
 935 73–78. 1006
- 936 (45) Karaman, S.; Digrak, M.; Ravid, U.; Ilcim, A. Antibacterial and 1007  
 937 antifungal activity of the essential oils of *Thymus revolutus* Celak from 1008  
 938 Turkey. *J. Ethnopharmacol.* **2001**, *76*, 183–186. 1009
- 939 (46) Wan, J.; Wilcock, A.; Coventry, M. J. The effect of essential oils of 1010  
 940 basil on the growth of *Aeromonas hydrophila* and *Pseudomonas* 1011  
 941 *fluorescence*. *J. Appl. Microbiol.* **1998**, *84*, 152–158. 1012
- 942 (47) Fricke, G.; Hoyer, H.; Wermter, R.; Paulus, H. Staphylococcus 1013  
 943 aureus as an example of the influence of lipophilic components on 1014  
 944 the microbiological activity of aromatic extracts. *Arch. Lebensmit- 1015*  
 945 *telechn.* **1998**, *49*, 107–111. 1016
- 946 (48) Matasyoh, J. C.; Kiplimo, J. J.; Karubiu, N. M.; Hailstorks, T.P. 1017  
 947 Chemical composition and antimicrobial activity of essential oil of 1018  
 948 *Tarchonanthus camphorates*. *Food Chem.* **2007**, *101*, 1183–1187. 1019
- 949 (49) Mah, T.-F. C.; O'Toole, G.A. Mechanisms of biofilm resistance to 1020  
 950 antimicrobial agents. *Trends Microbiol.* **2001**, *9*, 34–39. 1021
- 951 (50) Lee, D. U.; Heinz, V.; Knorr, D. Effects of combination treatments 1022  
 952 of nisin and high-intensity ultrasound with high pressure on the 1023  
 953 microbial inactivation in liquid whole egg. *Innov. Food Sci. Emerg. 1024*  
 954 *Technol.* **2003**, *4*, 387–393. 1025
- 955 (51) Wilkins, K. M.; Board, R. G. Natural Antimicrobial Systems. In 1026  
 956 *Mechanisms of Action of Food Preservation Procedures*; Gould, G. W., 1027  
 957 Ed.; Elsevier: New York, 1989; pp 285–362. 1028
- 958 (52) Beuchat, L. R. Surface disinfection of raw produce. *Dairy Food 1029*  
 959 *Environ. Sanit.* **1992**, *12*, 6–9. 1030
- 960 (53) Lopez-Malo, A.; Alzamora, S. M.; Guerrero, S. Natural Antimicro- 1031  
 961 bials from Plants. In *Minimally Processed Fruits and Vegetables. 1032*  
 962 *Fundamentals Aspects and Applications* Alzamora, S. M.; Tapia, M. S.; 1033  
 963 Lopez-Malo, A., Eds.; Aspen Publishers: Gaithersburg, MD, 2000; 1034  
 964 pp 237–264. 1035
- 965 (54) Juven, B. J.; Kanner, J.; Schved, F.; Weisslowicz, H. Factors that 1036  
 966 interact with the antibacterial action of thyme essential oil and its 1037  
 967 active constituents. *J. Appl. Bacteriol.* **1994**, *76*, 626–631. 1038
- 968 (55) Bajpai, V. K.; Rahman, A.; Dung, N. T.; Huh, M. K.; Kang, S. C. In 1039  
 969 vitro inhibition of food spoilage and foodborne pathogenic bacteria 1040  
 970 by essential oil and leaf extracts of *Magnolia liliflora* Desr. *J. Food 1041*  
 971 *Sci.* **2008**, *73* (6), M314–M320. 1042
- 972 (56) Fung, D. Y. C.; Taylor, S.; Kahan, J. Effects of butylated hydro- 1043  
 973 xyanisole (BHA) and butylated hydroxitoluene (BHT) on growth 1044  
 974 and aflatoxin production of *Aspergillus flavus*. *J. Food Saf.* **1977**, *1*, 1045  
 975 39–51. 1046
- 976 (57) Rico-Munoz, E.; Bargiota, E.; Davidson, P. M. Effect of selected 1047  
 977 phenolic compounds on the membrane-bound adenosine triphos- 1048  
 978 phate of *Staphylococcus aureus*. *Food Microbiol.* **1987**, *4*, 239–249. 1049
- 979 (58) Kabara, J. J.; Eklund, T. Organic Acids and Esters. In *Food 1050*  
 980 *Preservatives*; Russel, N. J.; Gould, G. W., Eds.; Blackie & Son Ltd: 1051  
 981 Glasgow, Scotland, 1991; pp 44–71. 1052
- 982 (59) Helander, I. M.; Alakomi, H. L.; Latva-Kala, K.; Mattila-Sandholm, 1053  
 983 T.; Pol, I.; Smid, E. J.; Gorris, L. G. M.; von Wright, A. Characterisa- 1054  
 984 tion of the action of selected essential oil components on gram-negative 1055  
 985 bacteria. *J. Agric. Food Chem.* **1998**, *46*, 3590–3595. 1056
- (60) Connor, D. E.; Beuchat, L. R. Sensitivity of heat stressed yeasts to 986  
 essential oils of plants. *Appl. Environ. Microbiol.* **1984**, *47*, 229–233. 987
- (61) Janssen, A.; Scheffer, J.; Baerheim-Svensden, A. Antimicrobial 988  
 activity of essential oils: A 1976–1986 literature review on aspects 989  
 of test methods. *Planta Med.* **1986**, *53*, 395–398. 990
- (62) Chorianopoulos, N.; Evergetis, E.; Mallouchos, A.; Kalpoutzakis, 991  
 E.; Nychas, G. J.; Haroutounian, S. A. Characterization of the 992  
 essential oil volatiles of *Satureja thymbra* and *Satureja parnassica*: 993  
 Influence of harvesting time and antimicrobial activity. *J. Agric. Food 994*  
*Chem.* **2006**, *54*, 3139–3145. 995
- (63) Ting, W. T. E.; Diebel, K. E. Sensitivity of *L. monocytogenes* to 996  
 spices at two temperatures. *J. Food Saf.* **1992**, *12*, 129–137. 997
- (64) Remmal, A.; Bouchikhi, T.; Rhayour, K.; Ettayebi, M. Improved 998  
 method for the determination of antimicrobial activity of essential 999  
 oils in agar medium. *J. Essent. Oil Res.* **1993**, *5*, 179–184. 1000
- (65) Pandit, V. A.; Shelef, L. A. Sensitivity of *Listeria monocytogenes* 1001  
 to rosemary (*Rosemarinus officinalis* L.). *Food Microbiol.* **1994**, *11*, 1002  
 57–63. 1003
- (66) Firouzi, R.; Azadbakht, M.; Nabinedjad, A. Anti-listerial activity of 1004  
 essential oils of some plants. *J. Appl. Anim. Res.* **1998**, *14*, 75–80. 1005
- (67) Hammer, K. A.; Carson, C. F.; Riley, T. V. Antimicrobial activity of 1006  
 essential oils and other plant extracts. *J. Appl. Microbiol.* **1999**, *86*, 1007  
 985–990. 1008
- (68) Campo, J. D.; Amiot, M. J.; Nguyen-The, C. Antimicrobial effect of 1009  
 rosemary extracts. *J. Food Prot.* **2000**, *63*, 1359–1368. 1010
- (69) Griffin, S. G.; Markham, J. L.; Leach, D. N. An agar dilution 1011  
 method for the determination of the minimum inhibitory concentra- 1012  
 tion of essential oils. *J. Essent. Oil Res.* **2000**, *12*, 249–255. 1013
- (70) Elgayyar, M.; Draughon, F. A.; Golden, D. A.; Mount, J. R. 1014  
 Antimicrobial activity of essential oils from plants against selected 1015  
 pathogenic and saprophytic microorganisms. *J. Food Prot.* **2001**, *64*, 1016  
 1019–1024. 1017
- (71) Delaquis, P. J.; Stanich, K.; Girard, B.; Mazza, G. Antimicrobial 1018  
 activity of individual and mixed fractions of dill, cilantro, coriander 1019  
 and eucalyptus essential oils. *Int. J. Food Microbiol.* **2002**, *74*, 1020  
 101–109. 1021
- (72) Roller, S.; Covill, N. The antifungal properties of chitosan in 1022  
 laboratory media and apple juice. *Int. J. Food Microbiol.* **1999**, *47*, 1023  
 67–77. 1024
- (73) Singh, N.; Singh, R. K.; Singh, A.; Bhuniab, A. K. Efficacy of plant 1025  
 essential oils as antimicrobial agents against *Listeria monocytogenes* 1026  
 in hotdogs. *LWT-Food Sci. Technol.* **2003**, *36*, 787–794. 1027
- (74) Gutierrez, J.; Rodriguez, G.; Barry-Ryan, C.; Bourke, P. The 1028  
 antimicrobial efficacy of plant essential oil combinations and 1029  
 interactions with food ingredients. *Int. J. Food Microbiol.* **2008**, 1030  
*124*, 91–97. 1031
- (75) Gutierrez, J.; Barry-Ryan, C.; Bourke, P. Antimicrobial activity of 1032  
 plant essential oils using food model media: Efficacy, synergistic 1033  
 potential and interactions with food components. *Food Microbiol.* 1034  
**2009**, *26*, 142–152. 1035
- (76) Scott, D.; Hammer, F. E.; Szalkucki, T. J. Bioconversions: Enzyme 1036  
 Technology. In *Food Biotechnology*; Knorr, D., Ed.; Marcel Dekker: 1037  
 New York, 1987; p 625. 1038
- (77) Borch, E.; Wallentin, C.; Rosen, M.; Bjorck, L. Antibacterial effect 1039  
 of the lactoperoxidase/thiocyanate/hydrogen peroxide system 1040  
 against strains of *Campylobacter* isolated from poultry. *J. Food 1041*  
*Prot.* **1989**, *52*, 638–641. 1042
- (78) Siragusa, G. R.; Johnson, M. G. Inhibition of *Listeria monocyto-* 1043  
*genes* growth by the lactoperoxidase thiocyanate H<sub>2</sub>O<sub>2</sub> antimicroa- 1044  
 bial system. *Appl. Environ. Microbiol.* **1989**, *55*, 2802–2805. 1045
- (79) Russel, A. D. Mechanisms of bacterial resistance to nonantibiotics: 1046  
 food additives and food and pharmaceutical preservatives. *J. Appl. 1047*  
*Bacteriol.* **1991**, *71*, 191–201. 1048
- (80) de Wir, J. N.; van Hooydonk, A. C. M. Structure, functions and 1049  
 applications of lactoperoxidase in natural antimicrobial systems. 1050  
*Neth. Milk Dairy J.* **1996**, *50*, 227–244. 1051
- (81) Delvesbroughton, J. Nisin and its application as a food preservative. 1052  
*J. Soc. Dairy Technol.* **1990**, *43* (3), 73–76. 1053
- (82) Park, C.; Lee, J.; Park, I.; Kim, M.; Kim, S. A Novel antimicrobial 1054  
 peptide from loach, *Misgurnus anguillicaudatus*. *FEBS Lett.* **1997**, 1055  
*411*, 173–178. 1056

- 1057 (83) Tossi, A.; Sandri, L.; Giangaspero, A. Amphipathic, helical anti-  
1058 microbial peptides. *Biopolymers* **2000**, *55*, 4–30.
- 1059 (84) Silphaduang, U.; Noga, E. J. Peptide antibiotics in mast cells of fish.  
1060 *Nature* **2001**, *414*, 268–2699.
- 1061 (85) Zasloff, M.; Martin, B.; Chen, H. Antimicrobial activity of synthetic  
1062 magainin peptides and several analogues. *Proc. Natl. Acad. Sci. U.S.A.* **1988**, *85*, 910–913.
- 1063 (86) Ge, Y.; Yan, H. Extraction of natural Vitamin E from wheat germ  
1064 by supercritical carbon dioxide. *J. Agric. Food Chem.* **2002**, *50* (4),  
1065 685–689.
- 1066 (87) Shi, J.; Ross, C. R.; Chengappa, M. M.; Style, M. J.; McVey, D.  
1067 S.; Blecha, F. Antibacterial activity of a synthetic peptide (PR-26)  
1068 derived from PR-39, a prolinearginine-rich neutrophil anti-  
1069 microbial peptide. *Antimicrob. Agents Chemother.* **1996**, *40*, 115–  
1070 121.
- 1071 (88) Thouzeau, C.; Le Maho, Y.; Froget, G.; Sabatier, L.; Le Bohec, C.;  
1072 Hoffmann, J. A.; Bulet, P. Spheniscins, avian beta-defensins in  
1073 preserved stomach contents of the king penguin, *Aptenodytes pata-*  
1074 *gonicus*. *J. Biol. Chem.* **2003**, *278*, 51053–51058.
- 1075 (89) Cole, A. M.; Weis, P.; Diamond, G. Isolation and characterization of  
1076 pleurocidin, an antimicrobial peptide in the skin secretions of winter  
1077 flounder. *J. Biol. Chem.* **1997**, *272*, 12008–12013.
- 1078 (90) Mor, A.; Nicolas, P. Isolation and structure of novel defensive  
1079 peptides from frog skin. *Eur. J. Biochem.* **1994**, *219*, 145–154.
- 1080 (91) Rydlo, T.; Rotem, S.; Mor, A. Antibacterial properties of derma-  
1081 septin s4 derivatives under extreme incubation conditions. *Antimi-*  
1082 *cro. Agents Chemother.* **2006**, *50* (2), 490–497.
- 1083 (92) Lee, I. H.; Cho, Y.; Lehrer, R. Effects of pH and salinity on the  
1084 antimicrobial properties of clavanins. *Infect. Immunol.* **1997**, *65*,  
1085 2898–2903.
- 1086 (93) Wilcox, W.; Eisenberg, D. Thermodynamics of melittin tetrameriza-  
1087 tion determined by circular dichroism and implications for protein  
1088 folding. *Protein Sci.* **1992**, *1*, 641–653.
- 1089 (94) Johansson, J.; Gudmundsson, G. H.; Rottenberg, M. E.; Berndt, K.  
1090 D.; Agerberth, B. Conformation-dependent antibacterial activity of  
1091 the naturally occurring human peptide LL-37. *J. Biol. Chem.* **1998**,  
1092 *273* (6), 3718–3724.
- 1093 (95) Ganzle, M. G.; Hertel, C.; Hammes, W. P. Resistance of *Escherichia*  
1094 *coli* and *Salmonella* against nisin and curvacin A. *Int. J. Food*  
1095 *Microbiol.* **1999**, *48*, 37–50.
- 1096 (96) Cole, A.; Darouiche, R.; Legarda, D.; Connell, N.; Diamond, G.  
1097 Characterization of a fish antimicrobial peptide: gene expression,  
1098 subcellular localization and spectrum of activity. *Antimicrob. Agents*  
1099 *Chemother.* **2000**, *44*, 2039–2045.
- 1100 (97) Burrowes, O. J.; Hadjicharalambous, C.; Diamond, G.; Lee, T. C.  
1101 Evaluation of antimicrobial spectrum and cytotoxic activity of  
1102 pleurocidin for food applications. *J. Food Sci.* **2004**, *69* (3), 66–71.
- 1103 (98) Brockus, C. W.; Jackwood, M. W.; Hamon, B. G. Characterization  
1104 of  $\beta$ -defensin prepropeptide from chicken and turkey bone marrow.  
1105 *Anim. Genet.* **1998**, *29*, 283–289.
- 1106 (99) Zhao, C.; Nguyen, T.; Liu, L.; Sacco, R. E.; Brogden, K. A.; Lehrer,  
1107 R. I. Gallinacin-3, an inducible epithelial beta-defensin I chicken.  
1108 *Infect. Immunol.* **2001**, *69*, 2684–2691.
- 1109 (100) Ganz, T. Defensin: antimicrobial peptides of intimate immunity.  
1110 *Nat. Rev. Immunol.* **2003**, *3*, 710–720.
- 1111 (101) Higazi, A. A. R.; Ganz, T.; Kariko, K.; Cines, D. B. Defensin  
1112 modulates tissue-type plasminogen activator and plasminogen  
1113 binding to fibrin and endothelial cells. *J. Biol. Chem.* **1996**, *271*,  
1114 17650–17655.
- 1115 (102) Murdock, C. A.; Cleveland, J.; Matthews, K. R.; Chikindas, M. L.  
1116 the synergistic effect of nisin and lactoferrin on the inhibition of  
1117 *Listeria monocytogenes* and *Escherichia coli* O157:H7. *Let. Appl.*  
1118 *Microbiol.* **2007**, *44* (3), 255–261.
- 1119 (103) Naidu, A. S. Activated lactoferrin – A new approach to meat  
1120 safety. *Food Technol.* **2002**, *56* (3), 40–45.
- 1121 (104) Al-Nabulsi, A. A.; Holley, R. A. Effects of *Escherichia coli* O157:  
1122 H7 and meat starter cultures of bovine lactoferrin in broth and  
1123 microencapsulated lactoferrin in dry sausage batters. *Int. J. Food*  
1124 *Microbiol.* **2007**, *113*, 84–91.
- 1125 (105) Al-Nabulsi, A. A.; Ran, J. H.; Liu, Z. Q.; Rodrigues-Vleira, E. T.;  
1126 Holley, R. A. Temperature-sensitive microcapsules containing  
1127 lactoferrin and their action against *Carnobacterium viridans* on  
Bologna. *J. Food Sci.* **2006**, *71* (6), M208–M214.
- 1128 (106) Del Olmo, A.; Morales, P.; Nunez, M. Bactericidal activity of  
1129 lactoferrin and its amidated and pepsin-digested derivatives against  
1130 *Pseudomonas fluorescens* in ground beef and meat fractions.  
1131 *J. Food Protect.* **2009**, *72* (4), 760–765.
- 1132 (107) Kagan, B. L.; Ganz, T.; Lehrer, R. I. Defensins: a family of  
1133 antimicrobial and cytotoxic peptides. *Toxicology* **1994**, *87*, 131–149.
- 1134 (108) Islam, N. M. D.; Itakura, T.; Motohiro, T. Antibacterial spectra  
1135 and minimum inhibition concentration of clupeine and salmine.  
1136 *Bull. Jpn. Soc. Sci. Fish.* **1984**, *50*, 1705–1078.
- 1137 (109) Uyttendaele, M.; Debevere, J. Protamine evaluation of the anti-  
1138 microbial activity of protamine. *Food Microbiol.* **1994**, *11*, 417–427.
- 1139 (110) Johansen, C.; Gill, T.; Gram, L. Antibacterial effect of protamine  
1140 assayed by impedimetry. *J. Appl. Bacteriol.* **1995**, *78*, 297–303.
- 1141 (111) Conte, M.; Aliberti, F.; Fucci, L.; Piscopo, M. Antimicrobial  
1142 activity of various cationic molecules on foodborne pathogens.  
1143 *World J. Microbiol. Biotechnol.* **2007**, *23*, 1679–1683.
- 1144 (112) Zasloff, M. Antimicrobial peptides of multicellular organisms.  
1145 *Nature* **1987**, *415*, 389–395.
- 1146 (113) Abler, L. A.; Klapes, N. A.; Sheldon, B. W.; Klaenhammer, T. R.  
1147 Inactivation of food-borne pathogens with magainin peptides.  
1148 *J. Food Prot.* **1995**, *58*, 381–388.
- 1149 (114) Coote, P. J.; Holyoak, C. D.; Bracey, D.; Ferdinando, D. P.; Pearce,  
1150 J. A. Inhibitory action of a truncated derivative of the amphibian  
1151 skin peptide dermaseptin s3 on *Saccharomyces cerevisiae*. *Antimi-*  
1152 *cro. Agents Chemother.* **1998**, *42*, 2160–2170.
- 1153 (115) Yaron, S.; Rydlo, T.; Shachar, D.; Mor, A. Activity of dermaseptin  
1154 K4-S4 against foodborne pathogens. *Peptides* **2003**, *24* (11), 1815–  
1155 1821.
- 1156 (116) Ben-Shalom, N.; Ardi, R.; Pinto, R.; Aki, C.; Fallik, E. Controlling  
1157 gray mould caused by *Botrytis cinerea* in cucumber plants by means  
1158 of chitosan. *Crop Prot.* **2003**, *22*, 285–290.
- 1159 (117) Je, J. Y.; Kim, S. K. Chitosan derivatives killed bacteria by  
1160 disrupting the outer and inner membrane. *J. Agric. Food Chem.*  
1161 **2006**, *54* (18), 6629–6633.
- 1162 (118) Chung, Y. C.; Wang, H. L.; Chen, Y. M.; Li, S. L. Effect of abiotic  
1163 factors on the antibacterial activity of chitosan against waterborne  
1164 pathogens. *Bioresour. Technol.* **2003**, *88*, 179–184.
- 1165 (119) Helander, I. M.; Nurmiho-Lassila, E.-L.; Ahvenainen, R.;  
1166 Rhoades, J.; Roller, S. Chitosan disrupts the barrier properties of  
1167 the outer membrane of gram-negative bacteria. *Int. J. Food Micro-*  
1168 *biol.* **2001**, *71*, 235–244.
- 1169 (120) Liu, X. F.; Guan, Y. L.; Yang, D. Z.; Li, Z.; Yao, K. D.  
1170 Antibacterial action of chitosan and carboxymethylated chitosan.  
1171 *J. Appl. Polym. Sci.* **2001**, *79*, 1324–1335.
- 1172 (121) Liu, N.; Chen, Xi-G.; Park, H.; Liu, C.; Liu, C.; Hong Meng, X.;  
1173 Yu, L. Effect of MW and concentration of chitosan on antibacterial  
1174 activity of *Escherichia coli*. *Carbohydr. Polym.* **2006**, *64*, 60–65.
- 1175 (122) Fernandes, J. C.; Tavarira, F. K.; Soares, J. C.; Ramos, O. S.;  
1176 Monteiro, M. J.; Pintado, M. E.; Malcata, F. X. Antimicrobial  
1177 effects of chitosans and chitoooligosaccharides, upon *Staphylococcus*  
1178 *aureus* and *Escherichia coli*, in food model systems. *Food Microbiol.*  
1179 **2008**, *25*, 922–928.
- 1180 (123) Bibel, D. J.; Miller, S. J.; Brown, B. E.; Pandey, B. B.; Elias, P. M.;  
1181 Shinefield, H. M.; Aly, R. Antimicrobial activity of stratum  
1182 corneum lipids from normal and essential fatty acid-deficient mice.  
1183 *Dermatol.* **1989**, *92*, 632–638.
- 1184 (124) Isaacs, C. E.; Kim, K. S.; Thormar, H. Antiviral and antibacterial  
1185 lipids in milk and infant formula feeds. *Arch. Dis. Child* **1990**, *65*,  
1186 861–864.
- 1187 (125) Lampe, M. F.; Ballweber, L. M.; Isaacs, C. E.; Patton, D. L.;  
1188 Stamm, W. E. Inhibition of *Chlamydia trachomatis* by novel  
1189 antimicrobial lipids adapted from human breast milk. *Antimicrob.*  
1190 *Agents Chemother.* **1998**, *42* (5), 1239–1244.
- 1191 (126) Wang, L. L.; Johnson, E. A. Control of *Listeria monocytogenes* by  
1192 monoglycerides in food. *J. Food Prot.* **1997**, *60*, 131–138.
- 1193 (127) Shin, S. Y.; Bajpai, V. K.; Kim, H. R.; Kang, S. C. Antibacterial  
1194 activity of bioconverted eicosapentaenoic (EPA) and docosahex-  
1195 aenoic acid (DHA) against foodborne pathogenic bacteria. *Int. J.*  
1196 *Food Microbiol.* **2007**, *113*, 233–236.
- 1197  
1198  
1199  
1200

- 1199 (128) Zasloff, M. Antimicrobial peptides of multicellular organisms. *Nature* **2002**, *415*, 389–395. 1270
- 1200 (129) Brogden, K. A. Antimicrobial peptides: pore formers or metabolic 1271
- 1201 inhibitors in bacteria? *Nat. Rev. Microbiol.* **2005**, *3*, 238–250. 1272
- 1202 (130) Boman, H. G.; Faye, I.; Gudmundsson, G. H.; Lee, J. Y.; Lidholm, 1273
- 1203 D. A. Cell-free immunity in *Cecropia*. A model system for anti- 1274
- 1204 bacterial proteins. *Eur. J. Biochem.* **1991**, *201*, 23–31. 1275
- 1205 (131) Meister, M.; Lemaitre, B.; Hoffmann, J. A. Antimicrobial peptide 1276
- 1206 defense in *Drosophila*. *Bioessays* **1997**, *19*, 1019–1026. 1277
- 1207 (132) Tailor, R. H.; Acland, D. P.; Attenborough, S.; Cammue, B. P.; 1278
- 1208 Evans, I. J.; Osborn, R. W.; Ray, J. A.; Rees, S. B.; Broekaert, W. F. 1279
- 1209 A novel family of small cysteine-rich antimicrobial peptides from 1280
- 1210 seed of *Impatiens balsamina* is derived from a single precursor protein. 1281
- 1211 *J. Biol. Chem.* **1997**, *272*, 24480–24487. 1282
- 1212 (133) Hancock, R. E. W. Peptide antibiotics. *Lancet* **1997**, *349*, 418– 1283
- 1213 422. 1284
- 1214 (134) Yoshida, K.; Mukai, Y.; Niidome, T.; Takashi, C.; Tokunaga, Y.; 1285
- 1215 Hatakeyama, T.; Aoyagi, H. Interaction of pleurocidin and its 1286
- 1216 analogues with phospholipid membrane and their antimicrobial 1287
- 1217 activity. *J. Pept. Res.* **2001**, *57*, 119–126. 1288
- 1218 (135) Lampe, M. F.; Isaacs, C. E. Lactolipids. In *Natural Food Anti-* 1289
- 1219 *microbial Systems*; Naidu, A. S., Ed.; CRC Press: Boca Raton, FL, 1290
- 1220 2000; pp 159–163. 1291
- 1221 (136) Jones, M. V.; Anslow, P. A.; Anderson, W. A.; Cole, M. B.; Gould, 1292
- 1222 G. W. Food Preserving Combination of Lysozyme/12438–12442. 1293
- 1223 Nisin/Citrate, Unilever. EP 90307694.1, **1990**. 1294
- 1224 (137) Daeschel, M. A. Antimicrobial substances from lactic acid bacteria 1295
- 1225 for use as food preservatives. *Food Technol.* **1989**, *43* (1), 164–167. 1296
- 1226 (138) De Vugst, L.; Vandamme, E. J. *Bacteriocins of Lactic Acid* 1297
- 1227 *Bacteria: Microbiology, Genetics and Applications* Blackie Aca- 1298
- 1228 demics and Professional: London, 1994. 1299
- 1229 (139) Ennahar, S.; Sashihara, T.; Sonomoto, K.; Ishizaki, A. Class Iia 1300
- 1230 bacteriocins: Biosynthesis, structure, and activity. *FEMS Micro-* 1301
- 1231 *biol. Rev.* **2000**, *24*, 85–106. 1302
- 1232 (140) Jack, R. W.; Jung, G. Lantibiotics and microcins: polypeptides with 1303
- 1233 unusual hosp diversity. *Curr. Opin. Chem. Biol.* **2000**, *4*, 310–317. 1304
- 1234 (141) Cleveland, J.; Montville, T. J.; Nes, I. F.; Chikindas, M. L. 1305
- 1235 Bacteriocins: safe, natural antimicrobials for food preservation. 1306
- 1236 *Int. J. Food Microbiol.* **2001**, *71*, 1–20. 1307
- 1237 (142) McAuliffe, O.; Ross, R. P.; Hill, C. Lantibiotics: structure, 1308
- 1238 biosynthesis and mode of action. *FEMS Microbiol. Rev.* **2001**, *25*, 1309
- 1239 285–308. 1310
- 1240 (143) Cotter, P. D.; Hill, C.; Ross, R. P. Bacteriocins: Developing innate 1311
- 1241 immunity for food. *Nat. Rev. Microbiol.* **2005**, *3*, 777–788. 1312
- 1242 (144) Klaenhammer, T. R. Genetics of bacteriocins produced by lactic 1313
- 1243 acid bacteria. *FEMS Microbiol. Rev.* **1993**, *12*, 39–86. 1314
- 1244 (145) Nes, I. F.; Bao Diep, D.; Havarstein, L. S.; Brurberg, M. B.; Eijsink, 1315
- 1245 V.; Holo, H. Biosynthesis of bacteriocins of lactic aci bacteria. 1316
- 1246 *Antonie van Leeuwenhoek* **1996**, *70*, 113–128. 1317
- 1247 (146) Guder, A.; Wiedeman, I.; Sahl, H. G. Post translationally modified 1318
- 1248 bacteriocins the lantibiotics. *Biopolymers* **2000**, *55*, 62–73. 1319
- 1249 (147) Nes, I. F.; Holo, H. Class II antimicrobial peptides from lactic acid 1320
- 1250 bacteria. *Biopolymers* **2000**, *55*, 50–61. 1321
- 1251 (148) Riley, M. A.; Chavan, M. A. *Bacteriocins, Ecology and Evolution*; 1322
- 1252 Springer: Berlin, Germany, 2006. 1323
- 1253 (149) Vignolo, G.; Fadda, S.; DeKairuz, M. N.; De Ruiz Holgdo, A. A. 1324
- 1254 P.; Olivier, G. Control of *Listeria monocytogenes* in ground beef by 1325
- 1255 Lactocin 705, a bacteriocin prod by *L. casei* CRL 705. *Int. J. Food* 1326
- 1256 *Microbiol.* **1996**, *27*, 397–402. 1327
- 1257 (150) De Martinis, E. C. P.; Franco, D. G. M. Inhibition of *Listeria* 1328
- 1258 *monocytogenes* in a ponk prod by a *Lactobacillus sakei* strain. *Int. J.* 1329
- 1259 *Food Microbiol.* **1998**, *42*, 119–126. 1330
- 1260 (151) Bredholt, S.; Nesbakken, T.; Holck, A. Protective culture inhibit 1331
- 1261 growth of *Listeria monocytogenes* and *Escherichia coli* O157: H7 in 1332
- 1262 cooked, slied vacuum. And gas pacaged meat. *Int. J. Food* 1333
- 1263 *Microbiol.* **1999**, *53*, 43–52. 1334
- 1264 (152) Georgalaki, M. D.; Van den Berghe, E.; Kritikos, D.; Devreese, B.; 1335
- 1265 Van Beettmen, J.; Kalantzopoulos, G.; De Vuyst, L.; Tsakalidou, 1336
- 1266 E. Macedocin, a food-grade lantibiotic produced by *Streptococcus* 1337
- 1267 *macedonicus* ACA-DC 198. *Appl. Environ. Microb.* **2002**, *68* (12), 1338
- 1268 5891–5903. 1339
- 1269 (153) Nettles, C. G.; Barefoot, S. F. Biochem and genet characteristics of 1340
- 1270 bacteriocins of food associated lactic acid bacteria. *J. Food Prot.* 1271
- 1272 **1993**, *56*, 338–356. 1273
- 1273 (154) Delves Broughton, J.; Blackburn, P.; Evans, R. J.; Hugenholtz, J. 1274
- 1275 Applications of the bacteriocin, nisin. *Antonie van Leeuwenhoek* 1276
- 1276 **1996**, *69*, 193–202. 1277
- 1277 (155) Hansen, J. N. Nisin as a model food preservative. *Crit. Rev. Food* 1278
- 1278 *Sci. Nutr.* **1994**, *34*, 69–93. 1279
- 1279 (156) Montville, T. J.; Chen, Y. Mechanistic action of pediocin and nisin: 1280
- 1280 recent progress and unresolved questions. *Appl. Microbiol. Bio-* 1281
- 1281 *technol.* **1998**, *50*, 511–519. 1282
- 1282 (157) Axelsson, L. T.; Chung, T. C.; Dobrogosz, W. J.; Lindgren, S. E. 1283
- 1283 Production of a broad spectrum antimicrobial substance by *Lacto-* 1284
- 1284 *bacillus reuteri*. *Microbial Ecology in Health and Disease* **1989**, *2*, 1285
- 1285 131–136. 1286
- 1286 (158) Nom, M. J. R.; Rombouts, F. M. 'Fermentative Preservatron of 1287
- 1287 Plant Foods' in 1. *Appl. Bacterial Symp. Suppl.* **1992**, *73*, 1365– 1288
- 1288 1478. 1289
- 1289 (159) Talarico, T. L.; Dobrogosz, W. J. Chemical characterization of an 1290
- 1290 antimicrobial substance produced by *Lactobacillus reuteri*. *Anti-* 1291
- 1291 *microb. Agents Chemother.* **1989**, *33*, 674–679. 1292
- 1292 (160) El-Ziney, M. G.; van den Tempel, T.; Debevere, J. M.; Jakobsen, 1293
- 1293 M. Application of reuterin produced by *Lactobacillus reuteri* 12002 1294
- 1294 for meat decontamination and preservation. *J. Food Prot.* **1999**, *62*, 1295
- 1295 257–261. 1296
- 1296 (161) Arques, J. L.; Fernandez, J.; Gaya, P.; Nunez, M.; Rodriguez, E.; 1297
- 1297 Medina, M. Antimicrobial activity of reuterin in combination with 1298
- 1298 nisin against food-borne pathogens. *Int. J. Food Microbiol.* **2004**, 1299
- 1299 *95* (2), 225–229. 1300
- 1300 (162) Anastasiadou, S.; Papagianni, M.; Filioussis, G.; Ambrosiadis, I.; 1301
- 1301 Koidis, P. Pediocin SA-1, an antimicrobial peptide from *Pediococ-* 1302
- 1302 *cus acidilactici* NRRL B5627: Production conditions, purification 1303
- 1303 and characterization. *Bioresources Technol.* **2008**, *99*, 5384–5390. 1304
- 1304 (163) Bhunia, A. K.; Johnson, M. C.; Ray, B. Purification, characteriza- 1305
- 1305 tion and antimicrobial spectrum of a bacteriocin produced by 1306
- 1306 *Pediococcus acidilactici*. *J. Appl. Bacteriol.* **1988**, *65*, 261–268. 1307
- 1307 (164) Fella, T. J.; Karunaratne, D. N.; Hancock, R. E. W. Mode of action 1308
- 1308 of the antimicrobial peptide indolicidin. *J. Biol. Chem.* **1996**, *271*, 1309
- 1309 19298–303. 1310
- 1309 (165) Hancock, R. E. W.; Lehrer, R. Cationic peptides: a new source of 1311
- 1311 antibiotics. *Trends Biotechnol.* **1998**, *16*, 82–88. 1312
- 1312 (166) Branby-Smith, F. M. Bacteriocins: Applications in food preserva- 1313
- 1313 tions. *Trends Food Sci. Technol.* **1992**, *3*, 133–137. 1314
- 1314 (167) Coventry, M. J.; Muirhead, K.; Hickey, M. W. Partial character- 1315
- 1315 ization of pediocin PO2 and comparison with nisin for biopreserva- 1316
- 1316 tion of meat products. *Int. J. Food Microbiol.* **1995**, *26*, 133–145. 1317
- 1317 (168) Nettles-Cutter, C.; Siragusa, G. R. Decontamination of beef 1318
- 1318 carcass tissue with nisin using a pilot scale model carcass washer. 1319
- 1319 *Food Microbiol.* **1994**, *11* (6), 481–489. 1320
- 1320 (169) Samelis, J.; Bedie, G. K.; Sofos, J. N.; Belk, K. E.; Scanga, J. A.; 1321
- 1321 Smith, G. C. Combinations of nisin with organic acids or salts to 1322
- 1322 control *Listeria monocytogenes* on sliced pork bologna stored 1323
- 1323 at 4°C in vacuum packages. *LWT-Food Sci. Technol.* **2005**, *38* (1), 1324
- 1324 21–28. 1325
- 1325 (170) Geornaras, I.; Skandamis, P. N.; Belk, K. E.; Scanga, J. A.; 1326
- 1326 Kendall, P. A.; Smith, G. C.; Sofos, J. N. Postprocess control of 1327
- 1327 *Listeria monocytogenes* on commercial frankfurters formulated 1328
- 1328 with and without antimicrobials and stored at 10 degrees C. 1329
- 1329 *J. Food Protect.* **2006**, *69* (1), 53–61. 1330
- 1330 (171) El-khateib, T.; Yousef, A. E.; Ockerman, H. W. Inactivation and 1331
- 1331 attachment of *L. monocytogenes* on beef muscle treated with lactic 1332
- 1332 acid and selected bacteriocins. *J. Food Prot.* **1993**, *56* (1), 29–33. 1333
- 1333 (172) Hoover, D. G.; Steenson, L. R. *Bacteriocins of Lactic Acid Bacteria*; 1334
- 1334 Academic Press, Inc.: San Diego, CA, 1993. 1335
- 1335 (173) Eckner, K. F. Bacteriocins and food applications. *Dairy Food* 1336
- 1336 *Environ. Sanit.* **1992**, *12* (4), 204–209. 1337
- 1337 (174) Henning, S.; Metz, R.; Hammus, W. P. New aspects for the 1338
- 1338 application of nisin to food products based on its mode of action. 1339
- 1339 *Int. J. Food Microbiol.* **1986**, *3*, 141–155. 1340
- 1340 (175) Zhang, S. S.; Mustapha, A. Reduction of *Listeria monocytogenes* 1341
- 1341 and *Escherichia coli* O157: H7 numbers on vacuum-packaged fresh 1342

- 1341 beef treated with nisin or nisin combined with EDTA. *J. Food Prot.*  
1342 **1999**, *62*, 1123–1127.
- 1343 (176) Delves-Broughton, J.; Williams, G. C.; Wilkinson, S. The use of the  
1344 bacteriocin, nisin as a preservative in pasteurized liquid whole egg.  
1345 *Lett. Appl. Microbiol.* **1992**, *15*, 133–136.
- 1346 (177) Nassar, A.; Farrag, S. A. Nisin as inactivator to *L. monocytogenes*  
1347 in broth and in ground beef. *Assiut-Veter. Med. J.* **1995**, *32*,  
1348 198–206.
- 1349 (178) Mahadeo, M.; Tatini, S. R. The potential use of nisin to control  
1350 *L. monocytogenes* in poultry. *Lett. Appl. Microbiol.* **1994**, *18*,  
1351 323–326.
- 1352 (179) Komitopoulou, E.; Boziaris, I. S.; Davies, E. A.; Delves-Broughton,  
1353 J.; Adams, M. R. *Alicyclobacillus acidoterrestris* in fruit juices and  
1354 its control by nisin. *Int. J. Food Sci. Technol.* **1999**, *34* (1), 81–85.
- 1355 (180) Davies, E. A.; Milne, C. F.; Bevis, H. E.; Potter, R. W.; Harris, J.  
1356 M.; Williams, G. C.; Thomas, L. V.; Delves-Broughton, J. Effective  
1357 use of nisin to control lactic acid bacterial spoilage in vacuum-  
1358 packed bologna-type sausage. *J. Food Prot.* **1999**, *62* (9), 1004–1010.
- 1359 (181) Driessen, A. J. M.; van den Hooven, H. W.; Kuiper, W.; van der  
1360 Kamp, M.; Sahl, H. G.; Konnigs, R. N. H.; Konnigs, W. N.  
1361 Mechanistic studies of lantibiotic-induced permeabilization of  
1362 phospholipids vesicles. *Biochemistry* **1995**, *34*, 1606–1614.
- 1363 (182) Stevens, K. A.; Sheldon, B. W.; Klapes, N. A.; Klaenhammer, T. R.  
1364 Nisin treatment for inactivation of *Salmonella* species and other  
1365 Gram-negative bacteria. *App. Environ Microbiol.* **1991**, *57*, 3613–  
1366 3615.
- 1367 (183) Boziaris, I. S.; Adams, M. R. Temperature shock, injury and  
1368 transient sensitivity to nisin in Gram negatives. *J. Appl. Microbiol.*  
1369 **2001**, *91*, 715–724.
- 1370 (184) Crandall, A. D.; Montville, T. J. Nisin resistance in *Listeria*  
1371 *monocytogenes* ATCC 700302 is a complex phenotype. *Appl.*  
1372 *Environ. Microbiol.* **1998**, *64*, 231–237.
- 1373 (185) Davies, E. A.; Adams, M. R. Resistance of *Listeria monocytogenes*  
1374 to the bacteriocin nisin. *Int. J. Food Microbiol.* **1994**, *21* (4),  
1375 341–347.
- 1376 (186) Carson, C. F.; Hammer, K. A.; Riley, T. V. Broth microdilution  
1377 method for determining the susceptibility of *Escherichia coli* and  
1378 *Staphylococcus aureus* to the essential oil of *Melaleuca alternifolia*  
1379 (tea tree oil). *Microbios* **1995**, *82*, 181–185.
- 1380 (187) Lambert, R. J. W.; Lambert, R. A model for the efficacy of  
1381 combined inhibitors. *J. Appl. Microbiol.* **2003**, *95*, 734–743.
- 1382 (188) Lambert, R. J. W.; Pearson, J. Susceptibility testing: accurate and  
1383 reproducible minimum inhibitory concentration (MIC) and non-  
1384 inhibitory concentration (NIC) values. *J. Appl. Microbiol.* **2000**,  
1385 *91* (3), 453–462.
- 1386 (189) Koutsoumanis, K.; Lambropoulou, K.; Nychas, G. J. E. A predictive  
1387 model for the non-thermal inactivation of *Salmonella*  
1388 *enteritidis* in a food model system supplemented with a natural  
1389 antimicrobial. *Int. J. Food Microbiol.* **1999**, *49*, 67–74.
- 1390 (190) Tassou, C. C.; Koutsoumanis, K.; Nychas, G.-J. E. Inhibition of  
1391 *Salmonella enteritidis* and *Staphylococcus aureus* in nutrient broth  
1392 by mint essential oil. *Food Res. Int.* **2000**, *33*, 273–280.
- 1393 (191) Walsh, S. E.; Maillard, J.-Y.; Russell, A. D.; Catrenich, C. E.;  
1394 Charbonneau, D. L.; Bartolo, R.G. Activity and mechanism of  
1395 action of selective biocidal agents on Gram-positive and -negative  
1396 bacteria. *J. Appl. Microbiol.* **2003**, *94*, 240–247.
- 1397 (192) Guillier, L.; Naser, A. I.; Dubois-Brissonnet, F. Growth response  
1398 of salmonella typhimurium in the presence of natural and synthetic  
1399 antimicrobials: Estimation of MICs from three different models.  
1400 *J. Food Prot.* **2007**, *70*, 2243–2250.
- 1401 (193) Lui, Y. Q.; Zhang, Y. Z.; Gao, P. J. Novel concentration-killing  
1402 curve method for estimation of bactericidal potency of antibiotics in  
1403 an in vitro dynamic model. *Antimicrob. Agents Ch.* **2004**, 3884–  
1404 3891.
- 1405 (194) Devglieghere, F.; Francois, K.; Vereecken, K. M.; Geeraerd, A. H.;  
1406 Van Impe, J. F.; Debevere, J. Effect of chemicals on the microbial  
1407 evolution in foods. *J. Food Protect.* **2004**, 1977–1990.
- 1408 (195) Unda, J. R.; Molins, R. A.; Walker, H. W. *Clostridium sporogenes*  
1409 and *Listeria monocytogenes*: survival and inhibition in microwave-  
1410 ready beef roasts containing selected antimicrobials. *J. Food Sci.* **1991**,  
1411 *56*, 198–205.
- (196) Hao, Y. Y.; Brackett, R. E.; Doyle, M. P. Efficacy of plant extracts  
1412 in inhibiting *Aeromonas hydrophila* and *Listeria monocytogenes* in  
1413 refrigerated cooked poultry. *Food Microbiol.* **1998**, *15*, 367–378. 1414
- (197) Mytle, N.; Anderson, G. L.; Doyle, M. P.; Smith, M. A. Anti-  
1415 microbial activity of clove (*Syzygium aromaticum*) oil in inhibiting  
1416 *Listeria monocytogenes* on chicken frankfurters. *Food Control* **2006**,  
1417 *17*, 102–107. 1418
- (198) Shekarforoush, S. S.; Nazer, A. H. K.; Firouzi, R.; Rostami, M.  
1419 Effects of storage temperatures and essential oils of oregano  
1420 and nutmeg on the growth and survival of *Escherichia coli* O157:  
1421 H7 in barbecued chicken used in Iran. *Food Control* **2007**, *18*, 1428–  
1429 1433. 1423
- (199) Careaga, M.; Fernandez, E.; Dorantes, L.; Mota, L.; Jaramillo,  
1424 M. E.; Hernandez-Sancheza, H. Antibacterial activity of Capsicum  
1425 extract against *Salmonella typhimurium* and *Pseudomonas aerugi-*  
1426 *nosa* inoculated in raw beef meat. *Int. J. Food Microbiol.* **2003**, *83*,  
1427 331–335. 1428
- (200) Ahn, J.; Grun, I. U.; Mustapha, A. Effect of plant extracts on  
1429 microbial growth, color change, and lipid oxidation in cooked beef.  
1430 *Food Microbiol.* **2007**, *24*, 7–14. 1431
- (201) Uhart, M.; Maki, N.; Ravishankar, S. Effect of spices on growth  
1432 and survival of *Salmonella typhimurium* DT 104 in ground beef  
1433 stored at 4 and 8 C. *J. Food Saf.* **2006**, *26*, 115–125. 1434
- (202) Karapinar, M.; Sengun, I. Y. Antimicrobial effect of koruk juice  
1435 against *Salmonella typhimurium* on salad vegetables. *Food Control*  
1436 **2007**, *18*, 702–706. 1437
- (203) Martinez-Romero, D.; Guillen, F.; Valverde, J. M.; Bailen, G.;  
1438 Zapata, P.; Serrano, M.; Castillo, S.; Valero, D. Influence of  
1439 carvacrol on survival of *Botrytis cinerea* inoculated in table grapes.  
1440 *Int. J. Food Microbiol.* **2007**, *115*, 144–148. 1441
- (204) Valero, M.; Francés, E. Synergistic bactericidal effect of carvacrol,  
1442 cinnamaldehyde or thymol and refrigeration to inhibit *Bacillus*  
1443 *cereus* in carrot broth. *Food Microbiol.* **2006**, *23* (1), 68–73. 1444
- (205) Gutierrez, J.; Bourke, P.; Lonchamp, J.; Barry-Ryan, C. Impact of  
1445 plant essential oils on microbiological, organoleptic and quality  
1446 markers of minimally processed vegetables. *Innov. Food Sci. Emerg.*  
1447 *Technol.* **2009**, *10*, 195–202. 1448
- (206) Kotzekidou, P.; Giannakidis, P.; Boulamatsis, A. Antimicrobial  
1449 activity of some plant extracts and essential oils against foodborne  
1450 pathogens in vitro and on the fate of inoculated pathogens in  
1451 chocolate. *LWT Food Sci. Technol.* **2007**, *41*, 119–127. 1452
- (207) Lee, C. H.; Park, H. J.; Lee, D. S. Influence of antimicrobial  
1453 packaging on kinetics of spoilage microbial growth in milk and  
1454 orange juice. *J. Food Eng.* **2004**, *65*, 527–531. 1455
- (208) Nguyen, P.; Mittal, G. S. Inactivation of naturally occurring  
1456 microorganisms in tomato juice using pulsed electric field (PEF)  
1457 with and without antimicrobials. *Chem. Eng. Process.* **2007**, *46*,  
1458 360–365. 1459
- (209) Lemay, M.-J.; Choquette, J.; Delaquis, P. J.; Garipey, C.; Rodrigue,  
1460 N.; Saucier, L. Antimicrobial effect of natural preservatives in a  
1461 cooked and acidified chicken meat model. *Int. J. Food Microbiol.*  
1462 **2002**, *78*, 217–226. 1463
- (210) Grande, M. J.; Lucas, R.; Abriouel, H.; Valdivia, E.; Omar, N. B.;  
1464 Maqueda, M.; Martínez-Bueno, M.; Martínez-Cañamero, M.;  
1465 Gálvez, A. Inhibition of toxicogenic *Bacillus cereus* in rice-based  
1466 foods by enterocin AS-48. *Int. J. Food Microbiol.* **2006**, *106* (2),  
1467 185–194. 1468
- (211) Grande, M. J.; Lucas, R.; Abriouel, H.; Valdivia, E.; Omar, N. B.;  
1469 Maqueda, M. M.; Martínez-Cañamero, M.; Gálvez, A. Treatment  
1470 of vegetable sauces with enterocin AS-48 alone or in combination  
1471 with ehenolic compounds to inhibit proliferation of *Staphylococcus*  
1472 *aureus*. *J. Food Prot.* **2007**, *70* (2), 405–411. 1473
- (212) Coma, V. Bioactive packaging technologies for extended shelf life  
1474 of meat-based products. *Meat Sci.* **2008**, *78* (1–2), 90–103. 1475
- (213) Seydim, A. C.; Sarikus, G. Antimicrobial activity of whey protein  
1476 based edible films incorporated with oregano, rosemary and garlic  
1477 essential oils. *Food Res. Int.* **2006**, *39*, 639–644. 1478
- (214) Oussalah, L.; Caillet, S.; Salmieri, S.; Saucier, L.; Lacroix, M.  
1479 Antimicrobial and antioxidant effects of milk protein-based film  
1480 containing essential oils for the preservation of whole beef muscle.  
1481 *J. Agr. Food Chem.* **2004**, *52* (18), 5598–5605. 1482

- 1483 (215) Vermeiren, L.; Devlieghere, F.; van Beest, M.; de Kruijf, N.;  
1484 Debevere, J. Developments in the active packaging of foods. *Trends*  
1485 *Food Sci. Technol.* **1999**, *10*, 77–86. 1540
- 1486 (216) Hill, C.; Deegan, L. H.; Paul, D.; Cotter, P. D.; Ross, P. Bacteriocins:  
1487 Biological tools for bio-preservation and shelf-life extension. *Int.*  
1488 *Dairy J.* **2006**, *16*, 1058–1071. 1541
- 1489 (217) Stevens, K.; Sheldon, B.; Klapes, N.; Klaenhammer, T. Effect  
1490 of treatment conditions on nisin inactivation of Gram-negative  
1491 bacteria. *J. Food Prot.* **1992**, *55*, 763–766. 1542
- 1492 (218) Rajkovic, A.; Uyttendaele, M.; Courtens, T.; Debevere, J. Anti-  
1493 microbial effect of nisin and carvacrol and competition between  
1494 *Bacillus cereus* and *Bacillus circulans* in vacuum-packed potato puree.  
1495 *Food Microbiol.* **2005**, *22*, 189–197. 1543
- 1496 (219) Mansour, M.; Feicht, E. A.; Behechti, A.; Schramm, K. W.;  
1497 Kettrup, A. Determination photostability of selected agrochemi-  
1498 cals in water and soil. *Chemosphere* **1999**, *39* (4), 575–585. 1544
- 1499 (220) Mansour, M.; Milliere, J.-B. An inhibitory synergistic effect of a  
1500 nisin-monolaurin combination on *Bacillus* sp. Vegetative cells in  
1501 milk. *Food Microbiol.* **2001**, *18*, 87–94. 1545
- 1502 (221) McLay, J. C.; Kennedy, M. J.; Orourke, A. L.; Elliot, R. M.;  
1503 Simmonds, R. S. Inhibition of bacterial foodborne pathogens by  
1504 the lactoperoxidase system in combination with monolaurin. *Int. J.*  
1505 *Food Microbiol.* **2002**, *73* (1), 1–9. 1546
- 1506 (222) Nikaido, H.; Vaara, M. Outer Membrane. In *Escherichia coli and*  
1507 *Salmonella Typhimurium: Cellular and Molecular Biology*; Neidhardt,  
1508 F. C., Ed.; American Society of Microbiological Institute of Brewing:  
1509 Washington, DC, 1987; Vol. 92, pp 379–383. 1547
- 1510 (223) Molinos, A. C.; Abriouel, H.; Lopez, R. L.; Valdivia, E.; Omar, N.  
1511 B.; Galvez, A. Combined physico-chemical treatments based on  
1512 enterocin AS-48 for inactivation of Gram-negative bacteria in  
1513 soybean sprouts. *Food Chem. Toxicol.* **2008**, *46*, 2912–2921. 1548
- 1514 (224) Oh, D.; Marshall, D. L. Effect of pH on the minimum inhibitory  
1515 concentration of monolaurin against *Listeria monocytogenes*. *J.*  
1516 *Food Prot.* **1992**, *55*, 449–450. 1549
- 1517 (225) Roberts, C. M.; Hoover, D. G. Sensitivity of *Bacillus coagulans*  
1518 spores to combinations of high hydrostatic pressure, heat, acidity and  
1519 nisin. *J. Appl. Bacteriol.* **1996**, *81* (4), 363–368. 1550
- 1520 (226) Garcia-Graells, C.; Valckx, C.; Michiels, C. W. Inactivation of  
1521 *Escherichia coli* and *Listeria innocua* in milk by combined treatment  
1522 with high hydrostatic pressure and the lactoperoxidase system. *Appl.*  
1523 *Environ. Microbiol.* **2000**, *66* (10), 4173–4179. 1551
- 1524 (227) Cressy, H. K.; Jerrett, A. R.; Osborne, C. M.; Bremer, P. J. A novel  
1525 method for the reduction of numbers of *Listeria monocytogenes* by  
1526 freezing in combination with an essential oil in bacteriological  
1527 medium. *J. Food Prot.* **2003**, *66*, 390–395. 1552
- 1528 (228) Viedma, P. M.; López, A. S.; Omar, N. B.; Abriouel, H.; López, R.  
1529 L.; Valdivia, E.; Belloso, O. M.; Gálvez, A. Enhanced bactericidal  
1530 effect of enterocin AS-48 in combination with high-intensity  
1531 pulsed-electric field treatment against *Salmonella enterica* in apple  
1532 juice. *Int. J. Food Microbiol.* **2008**, *128*, 244–249. 1553
- 1533 (229) Skandamis, P.; Nychas, G.-J. Effect of oregano essential oil on  
1534 microbiological and physico-chemical attributes of minced meat  
1535 stored in air and modified atmospheres. *J. Appl. Microbiol.* **2001**,  
1536 *91*, 1011–1022. 1554
- 1537 (230) Matan, N.; Rimkeeree, H.; Mawson, A. J.; Chompreeda, P.;  
1538 Haruthaithanasan, V.; Parker, M. Antimicrobial activity of  
1539 cinnamon and clove oils under modified atmosphere conditions. *Int. J. Food Microbiol.* **2006**, *107* (2), 180–185. 1555
- (231) Seydim, A. C.; Sarikus, G. Antimicrobial activity of whey protein  
1542 based edible films incorporated with oregano, rosemary and garlic  
1543 essential oils. *Food Res. Int.* **2006**, *39* (5), 639–644. 1556
- (232) Penney, V.; Gemma Henderson, G.; Blum, C.; Johnson-Green, P.  
1544 The potential of phytopreservatives and nisin to control microbial  
1545 spoilage of minimally processed fruit yogurts. *Innov. Food Sci.*  
1546 *Emerg. Technol.* **2004**, *5*, 369–375. 1557
- (233) Belletti, N.; Lanciotti, R.; Patrignani, F.; Gardini, F. Antimicrobial  
1548 efficacy of citron essential oil on spoilage and pathogenic micro-  
1549 organisms in fruit-based salads. *J. Food Sci.* **2008**, *73* (7), M331–  
1550 M338. 1551
- (234) Chanjirakul, K.; Wang, C. Y.; Wang, S. Y.; Siriphanich, J. Effect of  
1552 natural volatile compounds on antioxidant capacity and antioxi-  
1553 dant enzymes in raspberries. *Postharvest Biol. Technol.* **2006**, *40*,  
1554 106–115. 1552
- (235) Ukuku, D. O.; Bari, M. L.; Kawamoto, S.; Isshiki, K. Use of  
1556 hydrogen peroxide in combination with nisin, sodium lactate and  
1557 citric acid for reducing transfer of bacterial pathogens from whole  
1558 melon surfaces to fresh-cut pieces. *Int. J. Food Microbiol.* **2005**, *104*,  
1559 225–233. 1553
- (236) Singh, N.; Singh, R. K.; Bhunia, A. K.; Stroshine, R. L. Efficacy of  
1561 chlorine dioxide, ozone, and thyme essential oil or a sequential  
1562 washing in killing *Escherichia coli* O157:H7 on lettuce and baby  
1563 carrots. *LWT-Food Sci. Technol.* **2002**, *35* (8), 720–729. 1554
- (237) Uyttendaele, M.; Neyts, K.; Vanderswalmen, H.; Notebaert, E.;  
1565 Debevere, J. Control of *Aeromonas* on minimally processed vege-  
1566 tables by decontamination with lactic acid, chlorinated water, or  
1567 thyme essential oil solution. *Int. J. Food Microbiol.* **2004**, *90* (3),  
1568 263–271. 1555
- (238) Beatriz, R.; Yolanda, S.; Myriam, Z.; Carmen, T.; Fernanda, R.  
1570 Antimicrobial activity of nisin against *Oenococcus oeni* and other  
1571 wine bacteria. *Int. J. Food Microbiol.* **2007**, *116* (1), 32–36. 1556
- (239) Calderon Miranda, M. L.; Barbosa Canovas, G. V.; Swanson, B. G.  
1573 Inactivation of *Listeria innocua* in skim milk by pulsed electric fields  
1574 and nisin. *Int. J. Food Microbiol.* **1999**, *51* (1), 19–30. 1557
- (240) Cava, R.; Nowak, E.; Taboada, A.; Marin-Iniesta, F. Antimicro-  
1576 bial activity of clove and cinnamon essential oils against *Listeria*  
1577 *monocytogenes* in pasteurized milk. *J. Food Prot.* **2007**, *70*, 2757–  
1578 2763. 1558
- (241) Mahmoud, B. S. M.; Yamazaki, K.; Miyashita, K.; Miyashita, K.;  
1580 Shin, I. I.; Suzuki, T. A new technology for fish preservation by  
1581 combined treatment with electrolyzed NaCl solutions and essential  
1582 oil compounds source. *Food Chem.* **2006**, *99* (4), 656–662. 1559
- (242) Tang, S.; Kerry, J. P.; Sheehan, D.; Buckley, J.; Morrissey, P. A.  
1584 Antioxidative effect of added tea catechins on susceptibility of  
1585 cooked red meat, poultry and fish patties to lipid oxidation. *Food*  
1586 *Res. Int.* **2001**, *34* (8), 651–657. 1560

---

Received February 26, 2009. Revised manuscript received May 22, 2009. 1589  
Accepted May 28, 2009. Funding for this research was provided under 1590  
National Development Plan, through Food Institutional Research 1591  
Measure, administered by Department of Agriculture, Fisheries & 1592  
Food, Ireland. 1593