

2014

## Atmospheric Cold Plasma Inactivation of Escherichia Coli, Salmonella Enterica Serovar Typhimurium and Listeria Monocytogenes Inoculated on Fresh Produce

Dana Ziuzina

*Technological University Dublin, dana.ziuzina@tudublin.ie*

Sonal Patil

*Technological University Dublin, sonalpatil81@gmail.com*

Patrick Cullen

*Technological University Dublin, pj.cullen@tudublin.ie*

*See next page for additional authors*

Follow this and additional works at: <https://arrow.tudublin.ie/schfsehart>



Part of the [Food Microbiology Commons](#), and the [Food Processing Commons](#)

---

### Recommended Citation

Ziuzina, D. et al. (2014). Atmospheric Cold Plasma inactivation of Escherichia coli, Salmonella enterica serovar Typhimurium and Listeria monocytogenes inoculated on fresh produce. *Food Microbiology*, 42, pp.109-116 DOI: 10.1016/j.fm.2014.02.007

This Article is brought to you for free and open access by the School of Food Science and Environmental Health at ARROW@TU Dublin. It has been accepted for inclusion in Articles by an authorized administrator of ARROW@TU Dublin. For more information, please contact [arrow.admin@tudublin.ie](mailto:arrow.admin@tudublin.ie), [aisling.coyne@tudublin.ie](mailto:aisling.coyne@tudublin.ie), [vera.kilshaw@tudublin.ie](mailto:vera.kilshaw@tudublin.ie).

Funder: The research leading to these results has received funding from the European Community

---


**Authors**

Dana Ziuzina, Sonal Patil, Patrick Cullen, Kevin Keener, and Paula Bourke

# Accepted Manuscript

Atmospheric Cold Plasma inactivation of *Escherichia coli*, *Salmonella enterica* serovar Typhimurium and *Listeria monocytogenes* inoculated on fresh produce

D. Ziuzina, S. Patil, P.J. Cullen, K.M. Keener, P. Bourke



PII: S0740-0020(14)00032-X

DOI: [10.1016/j.fm.2014.02.007](https://doi.org/10.1016/j.fm.2014.02.007)

Reference: YFMIC 2105

To appear in: *Food Microbiology*

Received Date: 24 June 2013

Revised Date: 13 January 2014

Accepted Date: 11 February 2014

Please cite this article as: Ziuzina, D., Patil, S., Cullen, P.J., Keener, K.M., Bourke, P., Atmospheric Cold Plasma inactivation of *Escherichia coli*, *Salmonella enterica* serovar Typhimurium and *Listeria monocytogenes* inoculated on fresh produce, *Food Microbiology* (2014), doi: 10.1016/j.fm.2014.02.007.

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

1 **Atmospheric Cold Plasma inactivation of *Escherichia coli*, *Salmonella enterica* serovar**  
2 **Typhimurium and *Listeria monocytogenes* inoculated on fresh produce**

3 D. Ziuzina<sup>1</sup>, S. Patil<sup>1</sup>, P.J. Cullen<sup>1,2</sup>, K.M. Keener<sup>3</sup> and P. Bourke<sup>1</sup>

4 1. School of Food Science and Environmental Health, Dublin Institute of Technology,  
5 Dublin, Ireland

6 2. School of Chemical Engineering, University of New South Wales, Sydney, Australia.

7 3. Purdue University, Nelson Hall of Food Science, West Lafayette, IN, USA

8 **Corresponding Author:**

9 Paula Bourke, School of Food Science and Environmental Health, Dublin Institute of  
10 Technology, Cathal Brugha Street, Dublin 1, Ireland.

11 E-mail: [paula.bourke@dit.ie](mailto:paula.bourke@dit.ie)

12 **Abstract**

13 Atmospheric cold plasma (ACP) represents a potential alternative to traditional methods for  
14 non-thermal decontamination of foods. In this study, the antimicrobial efficacy of a novel  
15 dielectric barrier discharge ACP device against *Escherichia coli*, *Salmonella enterica*  
16 Typhimurium and *Listeria monocytogenes* inoculated on cherry tomatoes and strawberries,  
17 was examined. Bacteria were spot inoculated on the produce surface, air dried and sealed  
18 inside a rigid polypropylene container. Samples were indirectly exposed (i.e. placed outside  
19 plasma discharge) to a high voltage (70kV<sub>RMS</sub>) air ACP and subsequently stored at room  
20 temperature for 24 h. ACP treatment for 10, 60 and 120 s resulted in reduction of *Salmonella*,  
21 *E. coli* and *L. monocytogenes* populations on tomato to undetectable levels from initial  
22 populations of 3.1, 6.3, and 6.7 log<sub>10</sub> CFU/sample, respectively. However, an extended ACP  
23 treatment time was necessary to reduce bacterial populations attached on the more complex

24 surface of strawberries. Treatment time for 300 s resulted in reduction of *E. coli*, *Salmonella*  
25 and *L. monocytogenes* populations by 3.5, 3.8 and 4.2 log<sub>10</sub> CFU/sample, respectively, and  
26 also effectively reduced the background microflora of tomatoes.

27 Highlights:

28 A key advantage of this in-package non-thermal decontamination approach is the possibility  
29 to eliminate of post-processing contamination of the produce, thus increasing microbiological  
30 food safety and extension of produce shelf life. Inactivation was dependent on fresh produce  
31 surface features.

32 Key words: Atmospheric cold plasma, decontamination efficacy, pathogenic bacteria, fresh  
33 produce, ozone.

## 34 1. Introduction

35 The benefits associated with consumption of fresh produce maintain a high consumer demand  
36 for a wide range of pre-packed ready to use products. Nevertheless, fresh produce may  
37 contribute to the transmission of bacterial, parasitic and viral pathogens (Abadias *et al.* 2008).  
38 In recent years, foodborne human illnesses resulting from contaminated fresh produce have  
39 been widely reported globally. Most reporting countries identified *Escherichia coli* O157:H7,  
40 *Listeria monocytogenes* and *Salmonella* spp. as the target pathogens capable of causing  
41 severe human infection and deaths (Rangel *et al.* 2005; Raybaudi-Massilia *et al.* 2009;  
42 Olaimat and Holley 2012; CDC, 2012). A wide range of fresh fruit and vegetable products  
43 have been implicated in foodborne infections, such as lettuce, sprouted seed, melon,  
44 tomatoes, radish, pepper, basil and other mixed salads (Fernandez *et al.* 2013; Olaimat and  
45 Holley 2012). Pathogens, such as *E. coli* O157:H7 and *Salmonella* may reside in protected  
46 sites on surface of the fresh produce and be able to survive for long periods of time beyond  
47 the expected shelf-life (Olaimat and Holley 2012). Flessa *et al.* (2005) reported that *L.*  
48 *monocytogenes* is capable of survival on the surface of fresh intact or cut strawberries

49 throughout the shelf life of the fruit and can survive on frozen strawberries for periods of 4  
50 weeks. A health hazard to the consumers may also arise due to the possible presence of  
51 microbial toxins as a consequence of produce contamination with spoilage bacteria (Issa-  
52 Zacharia *et al.* 2010).

53 Raw fruits and vegetables can become contaminated while growing or during harvesting,  
54 postharvest processing, storage or distribution (Cevallos-Cevallos *et al.* 2012). How bacteria  
55 attach and the strength of attachment has not been well understood, but once attached to the  
56 surface of fresh produce it is difficult to remove the pathogens by washing (Berger *et al.*  
57 2010; Warning and Datta 2013). Conventional postharvest washing and sanitising treatments  
58 are not highly effective for produce, often resulting in less than 2 log unit reductions of  
59 pathogens (Niemira 2012). Moreover, some low pH based preservation techniques may  
60 contribute to the bacterial adaption to acidic environment and subsequently increase their acid  
61 resistance (Roberts and Wiedmann 2005). Disinfection can become less effective when  
62 microorganisms are attached to produce surface include biofilm formation, concentration  
63 reduction of sanitizer near produce surface and accessibility of sanitizer to cells attached to  
64 rough surfaces (Wang *et al.* 2012). Pathogens can also attach to surface through interaction  
65 with epiphytic microflora and may be further protected by internalising which itself  
66 dependant on many produces phyllosphere characteristics (Erickson 2012).

67 Non-thermal antimicrobial treatments of fruits, vegetables and other food produce have been  
68 the subject of much research. Atmospheric cold plasma (ACP) technology is a relatively new  
69 approach aiming to improve microbiological safety in conjunction with maintenance of  
70 sensory attributes of the treated foods. A key process advantage is the minimal water usage.  
71 However, apart from issues associated with water mediated decontamination, it is likely that  
72 many of the features associated with minimal processing and phyllosphere of produce that  
73 impact on traditional washing decontamination, may also interact with the optimum

74 application of ACP. The antimicrobial efficacy and design of ACP systems including  
75 producer gas composition, electrode configuration as well as the type of bacteria and  
76 substrate varies widely among research studies (Fernandez *et al.* 2013, Niemira 2012;  
77 Noriega *et al.* 2011; Niemira and Sites 2008). The use of indirect plasma in conjunction with  
78 utilisation of closed chambers for decontamination of meat produce have been highlighted in  
79 recent studies conducted by Rod *et al.* (2012) and Frohling *et al.* (2012b). Our previous study  
80 also demonstrated the antimicrobial efficiency of indirect ACP exposure, where *E. coli* in a  
81 sealed package was readily inactivated within seconds (Ziuzina *et al.* 2013). However, there  
82 are limited numbers of reports based on in-package plasma decontamination of fresh fruits  
83 and vegetables (Fan *et al.* 2012; Klockow and Keener 2009). Therefore, the objective of this  
84 study was to evaluate the efficacy of indirect ACP generated inside a sealed package against  
85 *E. coli*, *Salmonella* and *L. monocytogenes* inoculated on cherry tomatoes and strawberries  
86 and to evaluate its potential to reduce background microflora present on cherry tomatoes and  
87 strawberries in order to increase the produce shelf life.

## 88 **2. Materials and methods**

### 89 **2.1. Bacterial strains and inocula preparation**

90 Three bacterial strains were used in this study. *Escherichia coli* NCTC 12900 was obtained  
91 from National Collection of type cultures of the Health Protection Agency (HPA, UK),  
92 *Salmonella enterica* Typhimurium ATCC 14028 and *Listeria monocytogenes* NCTC 11994  
93 were obtained from the microbiology stock culture of the School of Food Science and  
94 Environmental Health of the Dublin Institute of Technology. Stock cultures were maintained  
95 at -70°C in the form of protective beads (Technical Services Consultants Ltd, UK). One  
96 protective bead of each culture was streaked onto separate tryptic soy agar (TSA,  
97 ScharlauChemie, Spain), incubated at 37°C for 24 h and further maintained at 4°C. A single  
98 isolated colony of each culture was inoculated in tryptic soy broth without glucose (TSB-G,

99 ScharlauChemie, Spain) and incubated at 37°C for 18 h. The cells were harvested by  
100 centrifugation at 10,000 rpm for 10 min, washed twice in sterile phosphate buffered solution  
101 (PBS, Oxoid LTD, UK) and finally resuspended in PBS, resulting in concentration of 8-9  
102 Log<sub>10</sub> CFU/ml, which were further used as the working inoculum. The concentration of  
103 inoculum was confirmed by plating appropriate dilutions on TSA, followed by incubation at  
104 37°C for 24 h for *E. coli* and *Salmonella* and 48 h for *L. monocytogenes*.

## 105 **2.2. Preparation of produce**

106 Whole fresh cherry tomatoes and strawberries (Class 1, Origin: Spain) were purchased from  
107 the local supermarket and stored at 4°C until use. The tomatoes were 2±0.5 cm in diameter  
108 and 5-15 g in weight. Strawberries weight was approximately 10-20 g. The same produce  
109 cultivar was used for each experiment. Cherry tomatoes were sterilized with 70% of ethanol  
110 (Klerwipe 70/30, Shield Medicare LTD, Farnham, UK) in order to reduce the background  
111 microbial load before surface inoculation of respective bacterial strain. Sterilized samples  
112 were then washed with sterile deionized water to remove any remaining ethanol residue and  
113 allowed to dry in the laminar flow safety cabinet at 23°C for 1 h prior to inoculation (Mattson  
114 *et al.* 2011). In order to assess ACP treatment efficacy for reduction of the background  
115 microflora, unsterilized tomatoes were also used.

## 116 **2.3. Fresh produce inoculation procedure**

117 For inoculation, tomatoes and strawberries were placed with the blossom end down on sterile  
118 petri dishes. The samples were spot-inoculated with bacteria applying either 50 µl or 100 µl  
119 of a culture on the tomato or strawberry surface, respectively (Das *et al.* 2006; Mahmoud *et*  
120 *al.* 2007). The droplets were deposited in several different locations, ensuring that the  
121 inoculum did not flow to the side of the samples. Inoculated samples were dried for 1 h in  
122 laminar flow safety cabinet to allow the attachment of bacteria on the surface of produce  
123 prior to the ACP treatment.



#### 124 **2.4. Experimental design**

125 The ACP system utilised was a dielectric barrier discharge system previously described in  
126 Ziuzina *et al.* (2013), with a maximum high voltage output of 120 kV at 50 Hz. The distance  
127 between the two 15 cm diameter aluminium disk electrodes was 40 mm which was equal to  
128 the height of the polypropylene container (310 x 230 x 40 mm) utilised as both a sample  
129 holder and as a dielectric barrier.

130 Inoculated samples (four of either tomatoes or strawberries) were aseptically transferred on  
131 one of the corner of the container so as to expose the samples to indirect ACP discharge  
132 (Fig.1). The distance between the samples and centre of the electrodes was within the range  
133 from 140 mm to 160 mm. In order to evaluate ACP treatment efficacy against background  
134 microflora, uninoculated samples were used. After product loading, each container was  
135 sealed within a high barrier polypropylene film (Cryovac, B2630, USA) and placed between  
136 the aluminum electrodes of the transformer. The inoculated and uninoculated samples were  
137 treated with 70 kV<sub>RMS</sub> for 30 s - 300 s in air and at atmospheric pressure. All samples were  
138 subjected to a post-treatment storage time of 24 h at room temperature. In order to evaluate  
139 any possible effect of storage on the bacterial growth, inoculated control samples were stored  
140 for 24 h under similar conditions. All experiments were performed in duplicate and replicated  
141 three times to ensure reproducibility of the experimental data and are reported as log<sub>10</sub>  
142 CFU/sample.

#### 143 **2.5. Microbiological analysis**

144 For microbiological analysis, inoculated untreated control samples (to estimate initial  
145 attached bacterial population), inoculated untreated samples stored for 24 h (to assess the  
146 effect of storage on microbial growth), uninoculated untreated control samples (to determine  
147 initial background microflora), and either inoculated or uninoculated ACP treated samples  
148 were analyzed. The samples were aseptically transferred into separate sterile stomacher bags

149 (BA6041, Seward LTD, UK) with 10 ml of sterile MRD and hand rubbed for 2-3 min. The  
150 resulting suspension was serially diluted in MRD. The surviving *E. coli*, *Salmonella* and *L.*  
151 *monocytogenes* populations were determined by agar overlay method (Mahmoud 2010).  
152 Briefly, aliquots of an appropriate dilution were surface plated on TSA, incubated for 2-4 h,  
153 and overlaid with the appropriate selective media: Sorbitol MacConkey agar (SMAC,  
154 ScharlauChemie, Spain) supplemented with Cefixime-Tellurite (CT, Oxoid LTD, England)  
155 for *E. coli*, Xylose Lysine Deoxycholate agar (XLD, ScharlauChemie, Spain) for *Salmonella*,  
156 and polymyxin-acriflavine-LiCl-ceftazidime-aesculin-mannitol (PALCAM, ScharlauChemie,  
157 Spain) supplemented with PALCAM Listeria Selective Supplement (Oxoid LTD, England)  
158 for *L. monocytogenes*. Plates were then incubated for 24-48 h at 37°C.  
159 Surviving background microflora of the uninoculated samples was evaluated using non-  
160 selective media TSA for estimation of aerobic mesophilic bacteria and Potato Dextrose agar  
161 (PDA, ScharlauChemie, Spain) for estimation of yeasts and moulds, with further incubation  
162 of agar plates at 37°C and 25°C, for 48 h and 5 days, respectively. The limit of detection for  
163 bacterial recovery on food samples was 1.0 Log<sub>10</sub> CFU/sample.

164

## 165 **2.6. Ozone measurements**

166 Ozone concentration inside the sealed package was measured using Gastec ozone detector  
167 tubes (Product #18M, Gastec Corporation, Japan). Measurements were taken immediately  
168 after plasma treatment and after 24 h of post treatment storage.

169

## 170 **2.7. Scanning Electron Microscopy (SEM)**

171 Attachment of different bacteria, namely *E. coli* and *L. monocytogenes*, attached on tomato  
172 and strawberry samples was observed using SEM. Inoculated strawberry samples were  
173 prepared as described by Gratao *et al.* (2008) with minor modifications. Briefly, the samples

174 were spot inoculated with either bacterium and dried under laminar flow at 23°C. The tissue  
175 from the inoculated sites of the fruit was excised forming 1 cm in diameter and 1 mm of  
176 thickness pieces. The cells were fixed in ice-cold 2.5% glutaraldehyde in 0.05 M sodium  
177 cacodylate buffer (pH7.4) (SCB) for 2 h. The cells were washed with the same buffer three  
178 times and fixed in 1% osmium tetroxide for 2 h at 4°C. After 2 h of fixation, bacterial cells  
179 were washed with SCB followed by three washes with distilled water. The samples were  
180 dehydrated using increasing concentrations of ethanol (30%, 50%, 70%, 80%, 95%, and  
181 99.5%) and freeze dried (Labconco, FreeZone 6; Mason Technology, Dublin, Ireland). In  
182 order to prevent surface charging by the electron beam, the samples were sputter-coated with  
183 gold particles using Emitech K575X Sputter Coating Unit resulting in a coating of 10 nm  
184 after 30 s. The samples were examined visually using a FEI Quanta 3D FEG Dual Beam  
185 SEM (FEI Ltd, Hillsboro, USA) at 5 kV.

## 186 **2.8. Statistical Analysis**

187 Statistical analysis was performed using SPSS 19.0 (SPSS Inc., Chicago, USA). The  
188 surviving population of *E. coli*, *Salmonella* and *L. monocytogenes* and ozone concentration  
189 following ACP treatment were subjected to analysis of variance (ANOVA). Means were  
190 compared according to the method of Fisher's Least Significant Difference-LSD at the 0.05  
191 level.

## 192 **3. Results**

193 Generally, indirect ACP treatment with subsequent 24 h of storage effectively reduced the  
194 numbers of microorganisms on either produce surface studied. On cherry tomatoes,  
195 treatments for 10 s, 60 s, and 120 s reduced populations of *Salmonella*, *E. coli* and *L.*  
196 *monocytogenes* to undetectable levels, respectively. However, an extended treatment time of  
197 300 s was necessary to reduce bacterial populations attached on the more complex surface of  
198 strawberries.

### 199 3.1. Inactivation of bacteria on cherry tomatoes

200 The influence of ACP treatments on viability of *E. coli*, *Salmonella* and *L. monocytogenes* is  
201 represented in Figure 2. Tomato samples were inoculated with an average of  $3.1 \pm 0.6 \log_{10}$   
202 CFU/sample for *E. coli*,  $6.3 \pm 0.6 \log_{10}$  CFU/sample for *Salmonella* and  $6.7 \pm 0.6 \log_{10}$   
203 CFU/sample for *L. monocytogenes*. After treatment for 10 s and above *Salmonella*  
204 populations on tomato were undetectable. Treatment for 45 s reduced populations of *E. coli*  
205 and *L. monocytogenes* by  $2 \pm 1.2$  and  $4.5 \pm 0.2 \log_{10}$  CFU/sample, respectively. Further  
206 increasing treatment time from 45 s to 60 s reduced populations of *L. monocytogenes* by  $5.1$   
207  $\pm 0.5 \log_{10}$  CFU/sample and reduced populations of *E. coli* to undetectable levels. Populations  
208 of *L. monocytogenes* were reduced to levels below detection limits after extended treatment  
209 for 120 s.

### 210 3.2. Inactivation of bacteria on strawberries

211 Reductions of *E. coli*, *Salmonella* and *L. monocytogenes* inoculated on strawberries are  
212 represented on Figure 3. The average initial attached population of *E. coli*, *Salmonella* and *L.*  
213 *monocytogenes* was  $4.4 \pm 1.7$ ,  $6.6 \pm 1.2$  and  $7.3 \pm 0.3 \log_{10}$  CFU/sample, respectively. After 60  
214 s and 120 s of ACP treatment, populations of *E. coli* were reduced by  $1.2 \pm 1.6$  and  $1.6 \pm 0.1$   
215  $\log_{10}$  CFU/sample, respectively, with significantly different reductions of  $3.5 \pm 0.7 \log_{10}$   
216 CFU/sample achieved after treatment for 300 s ( $P \leq 0.05$ ). Similarly, populations of  
217 *Salmonella* were reduced by  $1.7 \pm 0.1$  and  $3.8 \pm 0.4 \log_{10}$  CFU/sample after ACP exposure for  
218 120 s and 300 s, respectively. No significant difference in antimicrobial efficacy of ACP  
219 treatments for either 120 s or 300 s against *L. monocytogenes* was observed where average  
220 reductions of approximately  $4.2 \pm 0.5 \log_{10}$  CFU/sample were recorded. No changes were  
221 noticed in the levels of bacterial populations attached on the untreated control tomato or  
222 strawberries samples after storage for 24 h.

### 223 3.3. Inactivation of background microflora on produce

224 The reductions of background microflora on cherry tomatoes and strawberries due to indirect  
225 ACP treatments are represented on Figure 3.

226 An average of initial background microflora on cherry tomatoes was  $5 \pm 0.1 \log_{10}$   
227 CFU/sample (Fig. 4a). After 60 s of ACP treatment the aerobic mesophilic counts were  
228 reduced by  $3 \pm 0.7 \log_{10}$  CFU/sample while yeasts and moulds were reduced by  $2.5 \pm 0.6 \log_{10}$   
229 CFU/sample. Further increase in treatment time to 120 s resulted in reductions of yeasts and  
230 moulds to undetectable levels while population of mesophilic bacteria was reduced of by  $4.2$   
231  $\pm 0.8 \log_{10}$  CFU/sample. Mesophilic bacteria were not detected when the treatment time was  
232 increased to 300 s. Untreated and stored for 24 h samples showed no changes in the growth  
233 levels of background microflora on tomato samples.

234 Lower reduction levels of spoilage microorganisms by ACP treatment were observed in the  
235 case of strawberry samples (Fig. 4b). Significant decrease in mesophilic counts was observed  
236 after 60 s of ACP treatment, resulting in reductions by  $1.6 \pm 0.9 \log_{10}$  CFU/sample ( $P \leq 0.05$ )  
237 from the control  $3.6 \pm 0.3 \log_{10}$  CFU/sample. Populations of mesophilic bacteria did not  
238 decrease further when treatment time was extended from 60 s to either 120 s or 300 s.  
239 Populations of yeasts and moulds initially present on strawberries were  $5.5 \pm 0.1 \log_{10}$   
240 CFU/sample. These levels decreased by  $1.0 \pm 0.8 \log_{10}$  CFU/sample after 120 s of ACP  
241 treatment. Extending the treatment time from 120 s to 300 s resulted in an additional  $0.4 \pm 0.4$   
242 log reduction in the population of yeasts and moulds. It should be noted that the levels of  
243 mesophilic bacteria of untreated control strawberry samples increased by  $1.8 \pm 1.0 \log_{10}$   
244 CFU/sample during 24 h storage, whereas populations of yeasts and moulds remained the  
245 same.

#### 246 **3.4. Ozone generation**

247 Generation of ozone inside the sealed package containing either cherry tomatoes or  
248 strawberry samples as a function of ACP treatment time is represented in Figure 5. The ozone

249 concentration inside the package containing cherry tomatoes increased gradually with  
250 increasing the treatment time. All ACP treatment times studied resulted in significant  
251 increase of ozone concentration ( $P \leq 0.05$ ) with maximum concentration of 5600 ppm  
252 achieved after 300 s of treatment. However, no significant difference in ozone concentration  
253 generated during the treatment of strawberry samples was observed. ACP treatment for 60 s  
254 resulted in an average of 2800 ppm, and further increasing treatment time from 60 s to 120  
255 and 300 s resulted in an average of 3200 and 3500 ppm of ozone, respectively.

### 256 3.5. Scanning Electron Microscopy (SEM)

257 In order to examine if the complex substrate surface features had any effect on the bacterial  
258 adherence, and thus effect antimicrobial efficacy of ACP treatment, SEM analysis of  
259 untreated *E. coli* and *L. monocytogenes* inoculated on produce surface was conducted. Figure  
260 6(a,b) represents the surface of strawberry and tomato, respectively, inoculated with *L.*  
261 *monocytogenes* where strong bacterial attachment in the form of clusters was noticed. On the  
262 contrary, only a small amount of individually attached bacterial cells of *E. coli* on the rough  
263 surface of strawberry was found (Fig. 6c).

## 264 4. Discussion

265 The indirect ACP treatment showed better inactivation efficacy against inoculated challenge  
266 bacteria and background microflora present on the surface of the two different products  
267 tested. Cherry tomatoes were selected as they have been associated with recent foodborne  
268 illness outbreaks and represent common raw food ingredients of commercial salads.  
269 Strawberries are also popular fruits and consumed raw. Moreover, these produce types  
270 present different surface decontamination challenges to the ACP system, i.e. tomato surface  
271 which is smooth, and the more complex surface of strawberry - uneven with numerous seeds.  
272 In general, higher inactivation rates due to ACP treatment were achieved for bacteria  
273 inoculated on smooth surface of tomatoes. *Salmonella* and *E. coli* were more rapidly

274 inactivated on tomato than *L. monocytogenes*. Among the three bacteria studied, *Salmonella*  
275 was the most sensitive to ACP, where 10 s of treatment time reduced bacterial population to  
276 undetectable levels. For tomato, increasing treatment time enhanced the inactivation efficacy  
277 of ACP in the case of *E. coli* and *L. monocytogenes*. Increasing treatment time from 45 s to  
278 60 s inactivated *E. coli* populations present on tomatoes, whereas inactivation to undetectable  
279 levels of *L. monocytogenes* was obtained only after an extended treatment time of 120 s. It is  
280 reported that Gram positive bacteria are more resistant to ACP treatments than Gram negative  
281 (Montie *et al.* 2000; Lee *et al.* 2006; Ermolaeva *et al.* 2011; Frohling *et al.* 2012a), which was  
282 also clearly demonstrated in the current study. *Salmonella* and *E. coli* are Gram negative  
283 bacteria with a thinner outer membrane compared to the Gram positive *L. monocytogenes*.  
284 The thicker membrane of the Gram positive bacteria may present a barrier to the diffusion of  
285 plasma reactive species through the bacterial cell wall, thus impacting antimicrobial efficacy.  
286 On the contrary, Fan *et al.* (2012) revealed greater sensitivity of Gram positive *L. innocua*  
287 than Gram negative *Salmonella* and *E. coli* inoculated on tomato surface. Interestingly, other  
288 comparative studies reported similar susceptibility between Gram positive and Gram negative  
289 bacteria to ACP with respect to inactivation (Kostov *et al.* 2010; Olmez and Temur 2010;  
290 Klampfl *et al.* 2012). Clearly, the target cell characteristics are important factors for  
291 inactivation efficacy, but no clear trend is apparent and complex interactions with the system,  
292 process, surface or medium may also impact on efficacy in combination with cell type.

293 In this study we observed that the difference in the initial levels of the attached bacterial  
294 populations complicates the comparison of the bacterial sensitivity to the ACP treatments  
295 based on bacterial cell membrane characteristics. It is widely accepted that high initial  
296 bacterial concentration may affect inactivation efficacy of plasma treatment. The study  
297 conducted by Fernandez *et al.* (2012) clearly demonstrated that increasing the concentration  
298 of *S. Typhimurium* from 5 to 8 log<sub>10</sub> CFU/filter reduced the inactivation efficiency of ACP,

299 suggesting that the initial concentration of microorganisms present on foods plays an  
300 important role in the efficacy of plasma treatment. In the present work, the lower initial  
301 populations of *E. coli* attached on tomatoes surface did not necessarily contribute to the  
302 increased ACP bactericidal characteristics. Within 45 s of treatment populations of *E. coli*  
303 were reduced by 2 log from the initial 3.1 log<sub>10</sub> CFU/sample, whereas this treatment time  
304 resulted in the reductions of *L. monocytogenes* populations by 4.5 log from the initial 6.7  
305 log<sub>10</sub> CFU/sample, and only 10 s was required to reduce *Salmonella* by 6.3 log<sub>10</sub> CFU/sample.  
306 This indicates the importance of the mechanisms and strengths of bacterial attachment with  
307 respect to a decontamination procedure. It has also been demonstrated that the resistance to  
308 ACP may also vary between bacteria species. Despite the higher inoculation levels on tomato  
309 surface, *Salmonella* appeared to be more sensitive than *E. coli*. Similar results were achieved  
310 in the research conducted by Niemira and Sites (2008) where *Salmonella* Stanley was more  
311 sensitive to ACP than *E. coli* inoculated on both agar and apple surfaces.

312 The influence of the produce type on the overall antimicrobial efficacy of ACP was observed  
313 when results are compared with the strawberry decontamination study. Treatment for 120 s  
314 significantly reduced *L. monocytogenes* inoculated on strawberries. Increasing treatment time  
315 to 300 s did not yield any further reductions of bacteria. However, after 300 s of treatment, a  
316 proportional reduction of *E. coli* and *Salmonella* was achieved. Strawberry surface is more  
317 porous than the surface of tomato. Irregularities of the fruit surface may provide many niche  
318 areas for bacteria, providing physiological barrier or protection against ACP treatments. This  
319 factor probably contributed to the reduced ACP bactericidal effect on Gram negative bacteria  
320 on strawberries by comparison with tomatoes.

321 The influence of the complexity of the produce surface structure on inactivation efficacy of  
322 ACP was observed when treatments were evaluated for the reduction of background  
323 microflora naturally present on the produce. The causative agents of microbial spoilage in



324 fruits and vegetables can be bacteria (*Erwinia* spp., *Enterobacter* spp., *Propionibacterium*  
325 *chlohexanicum*, *Pseudomonas* spp., and lactic acid bacteria) as well as moulds and yeasts  
326 (*Penicillium* spp., *Aspergillus* spp., *Alternaria* spp., and *Saccharomyces* spp., *Cryptococcus*  
327 spp., *Rhodotorula* spp.) (Raybaudi-Massilia *et al.* 2009). In recent study conducted by Jensen  
328 *et al.* (2013), 34 different species from 23 different genera for bacteria and 22 different  
329 species from 9 different genera for yeasts were identified in strawberry samples. Despite this  
330 potential diversity of indigenous microflora, an ACP treatment time of 120 s significantly  
331 reduced the numbers on smooth surface of tomatoes in our study. However, again ACP was  
332 not very effective for the reduction of background microflora on more complex surface of  
333 strawberries, although tomato and strawberries tend to share similar bacterial communities  
334 (Leff and Fierer 2013).

335 Current information available for characterisation of ACP suggests that plasma is a source of  
336 heat, UV radiation, charged particles and reactive oxygen and nitrogen based species (ROS  
337 and RNS, respectively) with a main role given to the ROS as prime plasma disinfectants  
338 (Laroussi and Leipold 2004; Laroussi 2009). In this study, it was demonstrated that  
339 increasing the treatment time resulted in increased antimicrobial efficacy of ACP against  
340 bacteria inoculated on produce. Moreover, the inoculated samples were indirectly exposed to  
341 plasma, i.e., at some distance to the plasma discharge (~160 mm from the centre of the  
342 plasma discharge). In case of indirect treatment the charged particles and the short-lived  
343 species would not be expected to play a role due to their potential to recombine before  
344 reaching the sample (Laroussi 2009). Therefore, ozone was expected to be one of the key  
345 factors contributing to antimicrobial efficacy of ACP treatments. It has been demonstrated  
346 earlier that considerable reductions of bacteria by indirect ACP occurred within seconds  
347 when extended post treatment storage was applied, suggesting diffusion of the reactive  
348 species into liquids during post-treatment storage, thereby affecting microbial cells (Ziuzina

349 *et al.* 2013). Extended 24 h post treatment storage time was also employed in the current  
350 study. It is likely that 24 h post treatment storage time facilitated ACP action on the bacterial  
351 cells by retaining generated reactive species within closed container, thus, promoting  
352 diffusion of the species inside the product tissue.

353 In the current work, as the treatment time increased, a significant increase in the ozone  
354 generated by plasma inside the package containing produce was noted ( $P \leq 0.05$ ). However, it  
355 was also observed that the produce type influenced the concentration of ozone, where lower  
356 ozone levels were recorded for strawberry samples. Strawberries surface exhibit numerous  
357 pores, likely making the surface contact area larger than the area of tomato surface. This  
358 surface area differential may contribute to the increased dissolution rate of ozone generated  
359 inside the strawberry package, with subsequent reduced antimicrobial efficacy of ACP with  
360 regard to the all bacteria tested.

361 Considering the lower ozone concentrations and the consequent lower reductions of the  
362 challenge bacteria and background microflora on strawberries, it is likely that protection by  
363 more complex produce structures could be a critical parameter determining plasma treatment  
364 efficacy. Similarly, Fernandez *et al.* (2013) demonstrated that antimicrobial efficacy of  
365 plasma was influenced by produce surface features with higher bacterial reduction levels  
366 achieved on microbial filters than on more complex biotic surfaces.

367 As mentioned earlier, in this study, variations between initial populations of bacteria were  
368 apparent, with *Salmonella* and *L. monocytogenes* more readily attaching on the surface of  
369 either produce than *E. coli*. Regardless of the different surface features of the produce  
370 studied, SEM images confirmed the larger populations of *L. monocytogenes* adherent cells in  
371 addition to clusters of cells present. Despite the irregular nature of strawberry surface, which  
372 would probably facilitate bacterial attachment, *E. coli* populations visualised by SEM on the  
373 fruit surface were still less dense by comparison with *L. monocytogenes* images. A possible

374 explanation for the lower levels of attached *E. coli* is the presence and interaction with  
375 naturally existing indigenous epiphytic bacteria. Depending on the types of epiphyte present  
376 the survival of pathogens can be either enhanced or inhibited (Erickson 2012). For example,  
377 Cooley *et al.* (2006) demonstrated that one epiphyte *Enterobacter asburiae* isolated from  
378 lettuce inhibited colonisation of *E. coli*, whereas another epiphyte *Wausteria paucula* had the  
379 opposite effect; enhancing *E. coli* survival.

380 Other factors that may affect microbial attachment to fresh produce are the different  
381 morphology and chemistry of the produce as different fruits and vegetables offer different  
382 microniches for the attachment, penetration and proliferation of bacteria (Keeratipibul *et al.*  
383 2011). Motility of microorganisms facilitates pathogen entry into wounds, stomata and other  
384 existing fruit surface openings (Deering *et al.* 2012). We observed in SEM images that  
385 bacterial cells were likely adhered inside the natural crevices of produce surface or close to  
386 these regions. Naturally existing crack and pits on the surface of produce provide bacteria  
387 opportunity to internalise. Internalisation through the naturally existing opening is widely  
388 described in literature and considered as one of the major route of pathogens entry to plant  
389 tissue (Deering *et al.* 2012). Incidences of internalisation dependent upon concentration of  
390 bacteria, their location on the plant, age, integrity and stages of plant development, as well as  
391 indigenous agonistic/antagonistic bacteria present on plant have been reported (Erickson  
392 2012; Shi *et al.* 2009). This study indicated that the decontaminating effect of ACP is a  
393 function of produce type and the contaminating pathogen. The produce surface has an  
394 influence on pathogen attachment, with the potential for internalisation particularly  
395 associated with minimally processed fresh produce. Therefore the depth to which the plasma  
396 generated chemical species are able to diffuse through a tissue in order to affect internalised  
397 cells or those within a biofilm requires further investigation to elucidate how that diffusion  
398 capability of ACP can be effectively harnessed. Overall, the results of this study indicated

399 that bacterial attachment and increased survivability on more complex surfaces following  
400 ACP treatments should be considered as very important factors influencing treatment design.

#### 401 **Conclusion**

402 In summary, the high voltage indirect ACP treatment was highly efficient for  
403 decontamination of fresh produce inside a sealed package. Short treatment times of 10, 60 s  
404 and 120 s resulted in reductions to undetectable levels of *Salmonella*, *E. coli* and *L.*  
405 *monocytogenes*, respectively on cherry tomatoes. However, treatment times of up to 300 s  
406 were required to attain substantial reductions on strawberry surfaces. Similarly,  
407 yeasts/moulds and mesophiles on tomato surface were not detected after 120 to 300 s,  
408 respectively. Thus, it can be concluded that ACP treatment with 24 h post treatment storage  
409 can eliminate microorganisms on fresh produce surfaces inside a sealed package. In order to  
410 achieve optimum decontamination efficiency by ACP, factors including type of produce,  
411 their inherent surface characteristics, bacterial type, the strength and the nature of their  
412 attachment as well as the diffusion capacity of the plasma species, to holistically address the  
413 food safety issues associated with fresh produce, should be considered.

#### 414 **Acknowledgements**

415 The research leading to these results has received funding from the European Community's  
416 Seventh Framework Program (FP7/2207-2013) under grant agreement number 285820.

#### 417 **References:**

- 418 Abadias, M., Usall, J., Anguera, M., Solsona, C., and Viñas, I. (2008) Microbiological  
419 quality of fresh, minimally-processed fruit and vegetables, and sprouts from retail  
420 establishments. *International Journal of Food Microbiology*, 123, 121-129.
- 421 Berger, C. N., Sodha, S. V., Shaw, R. K., Griffin, P. M., Pink, D., Hand, P. and Frankel,  
422 G. (2010). Fresh fruit and vegetables as vehicles for the transmission of human  
423 pathogens. *Environmental Microbiology*, 12, 2385–2397.

- 424 Centers for Disease Control and Prevention (CDC). (2012). Multistate Outbreak of  
425 Listeriosis Linked to Whole Cantaloupes from Jensen Farms, Colorado. Available at  
426 <http://www.cdc.gov/listeria/outbreaks/cantaloupes-jensen-farms/082712/index.html>  
427 [Accessed on 26 March 2013].
- 428 Cevallos-Cevallos, J.M., Gu, G., Danyluk, M.D., Dufault, N.S., Ariena van Bruggen,  
429 H.C. (2012). *Salmonella* can reach tomato fruits on plants exposed to aerosols formed by  
430 rain. *International Journal of Food Microbiology*, 158, 140–146.
- 431 Cooley, M.B, Chao, D. and Mandrell, R.E. (2006). *Escherichia coli* O157:H7 survival  
432 and growth on lettuce is altered by the presence of epiphytic bacteria. *Journal of food*  
433 *protection*, 69, 2329-2335.
- 434 Das, E., Gürakan, G.C. and Bayindirli, A. (2006). Effect of controlled atmosphere  
435 storage, modified atmosphere packaging and gaseous ozone treatment on the survival of  
436 *Salmonella* Enteritidis on cherry tomatoes. *Food Microbiology*, 23, 430 - 8.
- 437 Deering, A.J., Mauer, L.J. and Pruitt, R.E. (2012). Internalization of *E. coli* O157:H7 and  
438 *Salmonella* spp. in plants: A review. *Food Research International*, 45, 567–575.
- 439 Erickson, M. C. (2012). Microbial ecology. Produce contamination. In Decontamination  
440 of Fresh and Minimally Processed Produce ed. Gomez-Lopez, V. M. pp. 3-29. USA:  
441 Wiley-Blackwell Publishing.
- 442 Ermolaeva, S.A., Varfolomeev, A.F., Chernukha, M. Yu., Yurov, D.S., Vasiliev, M.M.,  
443 Kaminskaya, A.A., Moisenovich, M.M., Romanova, J.M., Murashev, A.N., Selezneva,  
444 I.I., Shimizu, T., Sysolyatina, E.V., Shaginyan, I.A., Petrov, O.F., Mayevsky, E.I., Fortov,  
445 V.E., Morfill, G.E., Naroditsky, B.S. and Gintsburg, A.L. (2011). Bactericidal effects of  
446 non-thermal argon plasma in vitro, in biofilms and in the animal model of infected  
447 wounds. *Journal of Medical Microbiology*, 60, 75–83.
- 448 Fan, X., Sokorai, K.J., Engemann, J., Gurtler, J.B., Liu, Y. (2012). Inactivation of *Listeria*  
449 *innocua*, *Salmonella* Typhimurium, and *Escherichia coli* O157:H7 on surface and stem  
450 scar areas of tomatoes using in-package ozonation. *Journal of Food Protection*, 75, 1611-  
451 8.

- 452 Fernández, A., Noriega, E., and Thompson, A. (2013). Inactivation of *Salmonella*  
453 *enterica* serovar Typhimurium on fresh produce by cold atmospheric gas plasma  
454 technology. *Food Microbiology*, 33, 24 – 9.
- 455 Fernandez, A., Shearer, N., Wilson, D. R. and Thompson, A. (2012) Effect of microbial  
456 loading on the efficiency of cold atmospheric gas plasma inactivation of *Salmonella*  
457 *enterica* serovar Typhimurium. *International Journal of Food Microbiology* 152, 175 –  
458 180.
- 459 Flessa, S., Lusk, D.M., Harris, L.J. (2005). Survival of *Listeria monocytogenes* on fresh  
460 and frozen strawberries. *International Journal of Food Microbiology*, 101, 255– 262.
- 461 Frohling, A., Baier, M., Ehlbeck, J., Knorr, D. and Schluter, O. (2012a) Atmospheric  
462 pressure plasma treatment of *Listeria innocua* and *Escherichia coli* at polysaccharide  
463 surfaces: inactivation kinetics and flow cytometric characterisation. *Innovative food*  
464 *science and emerging technologies* 13, 142 – 150.
- 465 Frohling, A., Durek, J., Schnabel, U., Ehlbeck, J., Bolling, J., Schlüter, O. (2012b).  
466 Indirect plasma treatment of fresh pork: Decontamination efficiency and effects on  
467 quality attributes. *Innovative Food Science & Emerging Technologies*, 16, 381–390.
- 468 Gratao, P.L., Monteiro, C.C., Rossi, M.L., Martinelli, A.P., Peresc, L.E.P., Medici. L.O.,  
469 Lea, P.J. and Azevedo, R.A. (2009). Differential ultrastructural changes in tomato  
470 hormonal mutants exposed to cadmium. *Environmental and Experimental Botany*, 67,  
471 387–394.
- 472 Issa-Zacharia, A., Kamitani, Y., Muhimbula, H., and Iwasaki, K. (2010). Antimicrobial  
473 effect of slightly acidic electrolyzed water for inactivation of *Salmonella* spp. and  
474 *Escherichia coli* on fresh strawberries (*Fragaria L.*). *African Journal of Microbiology*  
475 *Research*, 4, 2174-2180.
- 476 Jensen, B., Knudsen, I.M.B., Andersen, B., Nielsen, K.F., Thrane, U., Jensen, D.F.,  
477 Larsen, J. (2013). Characterization of microbial communities and fungal metabolites on  
478 field grown strawberries from organic and conventional production. *International Journal*  
479 *of Food Microbiology*, 160, 313–322.

- 480 Keeratipibul, S., Phewpan, A., and Lursinsap, C. (2011). Prediction of coliforms and  
481 *Escherichia coli* on tomato fruits and lettuce leaves after sanitizing by using Artificial  
482 Neural Networks. *LWT - Food Science and Technology*, 44, 130 – 138.
- 483 Klampfl, T.G., Isbary, G., Shimizu, T., Li, Y.F., Zimmermann, J.L., Stolz, W., Schlegel,  
484 J., Morfilla, G.E. and Schmidt, H.U. (2012). Cold Atmospheric Air Plasma Sterilization  
485 against Spores and Other Microorganisms of Clinical Interest. *Applied and*  
486 *Environmental Microbiology*, 78(15):5077. doi: 10.1128/AEM.00583-12. Available at  
487 <http://aem.asm.org/content/78/15/5077.full> [Accessed on 17 April 2013].
- 488 Klockow, P.A., and Keener, K.M. (2009). Safety and quality assessment of packaged  
489 spinach treated with a novel ozone-generation system. *LWT - Food Science and*  
490 *Technology*, 42, 1047–1053.
- 491 Kostov, K.G., Rocha, V., Koga-Ito, C.Y., Matos, B.M., Algatti, M.A., Honda, R.Y.,  
492 Kayama, M.E. and Mota, R.P. (2010). Bacterial sterilization by a dielectric barrier  
493 discharge (DBD) in air. *Surface & Coatings Technology*, 204, 2954–2959.
- 494 Laroussi, M. 2009. Low-temperature plasmas for medicine? *Plasma Science, IEEE*  
495 *Transactions on plasma science* 37, 714-725.
- 496 Laroussi, M. and Leipold, F. (2004) Evaluation of the roles of reactive species, heat, and  
497 UV radiation in the inactivation of bacterial cells by air plasmas at atmospheric pressure.  
498 *International Journal of Mass Spectrometry* 233, 81 – 86
- 499 Lee, K., Paek, K., Ju, W.T, and Lee, Y. (2006). Sterilization of Bacteria, Yeast, and  
500 Bacterial Endospores by Atmospheric-Pressure Cold Plasma using Helium and Oxyge.  
501 *The Journal of Microbiology*, 44, 269-275
- 502 Leff, J.W and Fierer, N. (2013). Bacterial Communities Associated with the Surfaces of  
503 Fresh Fruits and Vegetables. *PLoS ONE* 8: e59310. doi:10.1371/journal.pone.0059310
- 504 Mahmoud, B.S.M. (2010). The effects of X-ray radiation on *Escherichia coli* O157:H7,  
505 *Listeria monocytogenes*, *Salmonella enterica* and *Shigella flexneri* inoculated on whole  
506 Roma tomatoes. *Food Microbiology*, 27, 1057-1063.

- 507 Mahmoud, B.S.M., Bhagat, A.R., and Linton, R.H. (2007). Inactivation kinetics of  
508 inoculated *Escherichia coli* O157:H7, *Listeria monocytogenes* and *Salmonella enterica*  
509 on strawberries by chlorine dioxide gas. *Food Microbiology*, 24, 36–744.
- 510 Mattson, T.E., Johny, A.K., Amalaradjou, M.A.R., More, K., Schreiber, D.T., Patel, J.,  
511 and Venkitanarayanan, K. (2011). Inactivation of *Salmonella* spp. on tomatoes by plant  
512 molecules. *International Journal of Food Microbiology*, 144, 464–468.
- 513 Montie, T. C., Kelly-Wintenber, K. and Roth, J. R. (2000). An overview of research  
514 using the one atmosphere uniform glow discharge plasma (OAUGDP) for sterilisation of  
515 surfaces and materials. *IEEE Transactions on plasma science* 28, 1
- 516 Niemira, B.A. (2012). Cold Plasma Decontamination of Foods. *Annual Review of Food*  
517 *Science and Technology*, 3, 125-142.
- 518 Niemira, B.A. (2012). Cold plasma reduction of *Salmonella* and *Escherichia coli*  
519 O157:H7 on almonds using ambient pressure gases. *Journal of Food Science*, 77, 171-  
520 175.
- 521 Niemira, B.A. and Sites, J. (2008). Cold plasma inactivates *Salmonella Stanley* and  
522 *Escherichia coli* O157:H7 inoculated on golden delicious apples. *Journal of Food*  
523 *Protection*. 71, 1357-65.
- 524 Noriega, E., Shama, G., Laca, A., Díaz, M., Kong, M.G. (2011). Cold atmospheric gas  
525 plasma disinfection of chicken meat and chicken skin contaminated with *Listeria*  
526 *innocua*. *Food Microbiology*, 7, 1293–1300.
- 527 Olaimat, A.N. and Holley R.A. (2012). Factors influencing the microbial safety of fresh  
528 produce: A review. *Food Microbiology* 32, 1 – 19.
- 529 Olmez H. and Temur S.D. (2010). Effects of different sanitizing treatments on biofilms  
530 and attachment of *Escherichia coli* and *Listeria monocytogenes* on green leaf lettuce.  
531 *LWT - Food Science and Technology* 43, 964–970.
- 532 Rangel, J. M., Sparling, P. H., Crowe, C., Griffin, P. M. and Swerdlow, D. L. (2005).  
533 Epidemiology of *Escherichia coli* O157:H7 Outbreaks, United States, 1982–2002.  
534 *Emerging infectious diseases*, 11, 603–609. doi: 10.3201/eid1104.040739 Available at  
535 <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC3320345/> [Accessed on 26 March 2013].



536 Raybaudi-Massilia, R.M., Mosqueda-Melgar, J., Soliva-Fortuny, R. and Martin-Belloso,  
537 O. (2009). Control of Pathogenic and Spoilage Microorganisms in Fresh-cut Fruits and  
538 Fruit Juices by Traditional and Alternative Natural Antimicrobials. *Comprehensive*  
539 *reviews in food science and food safety*, 8, 157 – 180.

540 Roberts, A. and Wiedmann, M. (2005). Host-pathogen interaction. In Understanding  
541 pathogen behaviour. Virulence, stress response and resistance ed. Griffiths, M. pp.98-114.  
542 UK: Woodhead Publishing Limited.

543 Rod, S. K., Hansen, F., Leipold, F. and Knochel, S. (2012) Cold atmospheric pressure  
544 plasma treatment of ready-to-eat meat: inactivation of *Listeria innocua* and changes in  
545 product quality. *Food microbiology* 30, 1-6

546 Shi, X., Wu, Z., Namvar, A., Kostrzynska, M., Dunfield, K. and Warriner, K. (2009).  
547 Microbial population profiles of the microflora associated with pre- and postharvest  
548 tomatoes contaminated with *Salmonella* Typhimurium or *Salmonella* Montevideo.  
549 *Journal of Applied Microbiology*, 107, 329–338

550 Wang, H., Zhou, B. and Feng, H. (2012). Surface characteristics of fresh produce and  
551 their impact on attachment and removal of human pathogens. Produce contamination. In  
552 Decontamination of Fresh and Minimally Processed Produce ed. Gomez-Lopez, V. M.  
553 pp. 43-55. USA: Wiley-Blackwell Publishing.

554 Warning, A. and Datta, A.K. (2013). Interdisciplinary engineering approaches to study  
555 how pathogenic bacteria interact with fresh produce. *Journal of Food Engineering*, 114,  
556 426–448.

557 Ziuzina, D., Patil, S., Cullen, P.J., Keener, K.M. and P. Bourke. (2013). Atmospheric cold  
558 plasma inactivation of *Escherichia coli* in liquid media inside a sealed package. *Journal*  
559 *of applied microbiology*, 114, 778-787.


560

561

562

563

564 **Figures:**



565

566 Fig.1: Schematic diagram of samples distributed within polypropylene container with respect  
567 to the electrodes.

568

569

570

571

572

573


574

575

576

577

578



579

580 Fig.2: ACP inactivation efficacy against *E. coli* (□), *Salmonella* (△) and *L. monocytogenes* (◇)  
581 inoculated on cherry tomatoes. Vertical bars represent standard deviation. Limit of detection  
582 1.0 log<sub>10</sub> CFU/sample.

583

584

585

586

587

588

589

590

591


592

593

594

595

596



597

598 Fig.3: ACP inactivation efficacy against *E. coli* (□), *Salmonella* (△) and *L. monocytogenes* (◇)  
599 inoculated on strawberries. Vertical bars represent standard deviation. Limit of detection 1.0  
600 log<sub>10</sub> CFU/sample.

601

602

603

604

605

606

607

608

609


610

611


612

613

614



615



616

617 Fig.4: ACP inactivation efficacy against aerobic mesophilic bacteria (◇) and yeasts and  
 618 moulds (□) on (a) cherry tomatoes and (b) strawberries. Vertical bars represent standard  
 619 deviation. Limit of detection 1.0 log<sub>10</sub> CFU/sample.


620

621

622

623

624



625

626 Fig.5: Generation of ozone inside a sealed package during ACP treatment of either inoculated  
627 or uninoculated samples of cherry tomatoes (■) and strawberries (□). Vertical bars represent  
628 standard deviation.

629

630

631

632

633

634

635

636

637

638

639

640

641

642

643

644



645

646

647

648

649



650

651


652

653

654

655

656



657

658 Fig.6: SEM images of untreated *L. monocytogenes* on (a) strawberries and (b) cherry

659 tomatoes, and (c) *E. coli* inoculated on strawberry.

660

661

662

663

664

**YFMIC 2105 Food Microbiology**

**Atmospheric Cold Plasma inactivation of *Escherichia coli*, *Salmonella enterica* serovar**

**Typhimurium and *Listeria monocytogenes* inoculated on fresh produce**

**Highlights**

- In this study antimicrobial efficacy of ACP against *Escherichia coli*, *Salmonella enterica* Typhimurium and *Listeria monocytogenes* inoculated on cherry tomatoes and strawberries was evaluated.
- A key advantage of this high voltage level treatment for in-package non-thermal decontamination approach is the possibility to eliminate post-processing contamination of the produce.
- This approach has potential to provide both increased microbiological food safety and extension of produce shelf life.
- Inactivation was however, dependent on fresh produce surface features and pathogen type.