

2023

Strabismus and Amblyopia in Africa – A Systematic Review and Meta-Analysis

Prince Kwaku Akowuah
University of Houston, USA

Samuel Adade
Kwame Nkrumah University of Science and Technology, Ghana

Andrews Nartey
University of Houston, USA

See next page for additional authors

Follow this and additional works at: <https://arrow.tudublin.ie/scschphyart>



Part of the [Optometry Commons](#)

Recommended Citation

Akowuah, Prince Kwaku; Adade, Samuel; Nartey, Andrews; Owusu, Ebenezer; Donkor, Richard; and Ankamah-Lomotey, Stephen, "Strabismus and Amblyopia in Africa – A Systematic Review and Meta-Analysis" (2023). *Articles*. 190.

<https://arrow.tudublin.ie/scschphyart/190>

This Article is brought to you for free and open access by the School of Physics, Clinical and Optometric Science at ARROW@TU Dublin. It has been accepted for inclusion in Articles by an authorized administrator of ARROW@TU Dublin. For more information, please contact arrow.admin@tudublin.ie, aisling.coyne@tudublin.ie, vera.kilshaw@tudublin.ie.



This work is licensed under a [Creative Commons Attribution-Share Alike 4.0 International License](#).

Funder: This research received no external funding

Authors

Prince Kwaku Akowuah, Samuel Adade, Andrews Nartey, Ebenezer Owusu, Richard Donkor, and Stephen Ankamah-Lomotey



Strabismus and amblyopia in Africa – a systematic review and meta-analysis

Prince Kwaku Akowuah, Samuel Adade, Andrews Nartey, Ebenezer Owusu, Richard Donkor, Stephen Ankamah-Lomotey, Asafo Agyei Frimpong, Joseph Adjei-Anang & Emmanuel Kobia-Acquah

To cite this article: Prince Kwaku Akowuah, Samuel Adade, Andrews Nartey, Ebenezer Owusu, Richard Donkor, Stephen Ankamah-Lomotey, Asafo Agyei Frimpong, Joseph Adjei-Anang & Emmanuel Kobia-Acquah (2023) Strabismus and amblyopia in Africa – a systematic review and meta-analysis, *Strabismus*, 31:1, 31-44, DOI: [10.1080/09273972.2022.2157023](https://doi.org/10.1080/09273972.2022.2157023)

To link to this article: <https://doi.org/10.1080/09273972.2022.2157023>



Published online: 28 Dec 2022.



Submit your article to this journal [↗](#)



Article views: 136











View related articles [↗](#)



View Crossmark data [↗](#)



Strabismus and amblyopia in Africa – a systematic review and meta-analysis

Prince Kwaku Akowuah, OD, PhD ^{a,b}, Samuel Adade, OD, PhD ^a, Andrews Nartey, OD, MSc, MPH^a, Ebenezer Owusu, OD ^a, Richard Donkor, OD, PhD ^c, Stephen Ankamah-Lomotey, OD, MSc ^b, Asafo Agyei Frimpong, OD ^b, Joseph Adjei-Anang, OD, MSc ^b, and Emmanuel Kobia-Acquah, OD, MSc ^{b,d}

^aCollege of Optometry, University of Houston, Houston, TX, USA; ^bDepartment of Optometry and Visual Science, Kwame Nkrumah University of Science and Technology; ^cSchool of Optometry and Vision Science, University of Waterloo, Waterloo, ON, Canada; ^dCentre for Eye Research Ireland, School of Physics, Clinical and Optometric Sciences, College of Sciences and Health, Technological University Dublin

ABSTRACT

The objective of this review was to estimate the prevalence of strabismus and amblyopia in Africa. A systematic online literature search was conducted for articles on strabismus and amblyopia in Africa. Meta-analysis was performed, using the Freeman-Tukey double arcsine transformation, to estimate the prevalence of strabismus and amblyopia in Africa. Subgroup analyses were conducted according to age, gender, study year, and type of amblyopia. Meta-regression was used to evaluate the influence of predetermined factors on the prevalence of amblyopia. 8 (1 population-based & 7 school-based) and 21 (3 population-based & 18 school-based) studies on strabismus and amblyopia with sample sizes of 22,355 and 46,841, respectively, were included in the review. Overall prevalence of strabismus in Africa was estimated to be 0.8% (95% CI: 0.4% – 1.4%); exotropia was 0.2% (95% CI: 0.1% – 0.5%) and esotropia was 0.5% (95% CI: 0.1% – 1.2%). Overall prevalence of amblyopia was estimated to be 0.6% (95% CI: 0.3% – 0.9%); refractive and strabismic amblyopia were 1.1% (95% CI: 0.2% – 2.5%) and 0.4% (95% CI: 0.2% – 0.6%), respectively. Prevalence estimate of amblyopia in males was 1.8% (95% CI: 0.7% – 3.3%) and in females was 1.3% (95% CI: 0.4% – 2.6%). There was a significant association between the prevalence of amblyopia and the type of amblyopia ($p = .007$) and the study year ($p = .006$). Although there appears to be a relatively low prevalence of strabismus and amblyopia in Africa, there is a dearth of well-designed population-based studies on strabismus and amblyopia in Africa, resulting in the lack of epidemiological information on strabismus and amblyopia within the general African population. Information about the prevalence of strabismus and amblyopia across Africa can inform policy making and design and implementation of public health intervention program.

KEYWORDS

Africa; amblyopia; meta-analysis; strabismus; systematic review

Introduction

Strabismus is a developmental disorder in which the two eyes are not aligned when viewing an object.¹ Strabismus is a leading cause of visual impairment especially in children, with an estimated prevalence of 2–5% globally.^{2–7} An estimated 0.80% of children in Singapore,⁸ 5.65% children in China⁹ and 2.47% children in the United States of America¹⁰ have strabismus. Strabismus can have significant effects on the development of children if not treated early, affecting learning and impairing performance.^{8,11} In adulthood, strabismus can affect the vision-related quality of life and have significant economic and public health significance.^{12–14} Given the significance of strabismus, it is important to know the burden and the public health implications of the

strabismus problem globally and across different geographical regions.

Strabismus is a common cause of amblyopia – which is a common developmental disorder of the visual system characterized by reduced best corrected visual acuity in one or both eyes (two-line difference between the eyes) with no obvious organic cause.^{15,16} Risk factors for the development of amblyopia include obstruction in the optical pathway that reduces retinal image quality (deprivation amblyopia), refractive error (refractive amblyopia) or ocular misalignment that disrupts binocular fusion (strabismic amblyopia).^{17,18} Amblyopia is the leading cause of monocular visual impairment in both children and adults and a common visual disorder

affecting all age groups.¹⁶ The global prevalence of amblyopia is estimated as 1–5% in children and 2.5–7.14% in adults.^{19–22} The variation in reported prevalence is due to factors such as characteristics of selected cohort of the study, visual acuity criterion, and type of amblyopia. Also, definitions of amblyopia in literature are usually very heterogenous, making prevalence from different studies difficult to compare. Amblyopia can be treated, if detected early and treatment is initiated early enough.²³ However, if left untreated, amblyopia can disrupt vision in diverse ways, affecting contrast sensitivity, visual acuity, and binocular vision.²⁴ In Africa, amblyopia and strabismus have been moderately studied in few regions with varying prevalence reported.^{25–36} Factors such as screening of school-going children to identify refractive errors and strabismus, spectacle coverage, availability of spectacles and/or contact lenses and eyecare personnel in Africa have an impact on incidence of amblyopia and strabismus.^{37–39}

To date, there is no continental representative study on the prevalence of strabismus or amblyopia in Africa. As such, the burden of strabismus and amblyopia in Africa is not known. Given the significance of both strabismus and amblyopia as major causes of visual impairment in both children and adults, it is important to have information of their prevalence on the continental level. Therefore, it is important to provide a pooled prevalence estimate of amblyopia and strabismus as well as associated risk factors in Africa. The aim of the current systematic review and meta-analysis is to sum up the current available literature and provide a current and comprehensive reflection of strabismus and amblyopia epidemiology in Africa using appropriate meta-analytic techniques. In this study, all published data in Africa were evaluated and the overall prevalence estimate of amblyopia and strabismus in Africa was reported.

Methods

Guidelines in the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) statement⁴⁰ were followed.

Literature search strategy

Between 1 June 2021 and 15 July 2021, the following databases were searched for studies reporting prevalence of strabismus and amblyopia in African populations: PubMed, Web of Science, African Journals Online (AJOL), Scopus and Embase. Google scholar was used to access gray literature. No time filters were applied to search. Different variations of search text and medical search headings (MeSH) were used in the literature search, each being an appropriate combination of any of the search words or terms in Table 1. The search terms were first developed in PubMed and was applied in other databases. The PICO of the study was: population (adults and children in Africa), intervention (none), comparison (none) and outcome (prevalence of strabismus and amblyopia). The PICO was used to define the search strategy. The bibliography of eligible studies (from primary literature search) was reviewed for any relevant publication that might have been missed during the initial literature search. Literature search was conducted independently by two authors.

Inclusion and exclusion criteria

School- or population-based studies with prospective cross-sectional design conducted in the “normal” population were included. For this review, normal subjects were defined as individuals without any systemic/genetic/syndromic condition (e.g., Down Syndrome) known to predispose to strabismus or amblyopia. Available full-text articles of studies were used. However, if the full text of a study is not available but the abstract provides

Table 1. Search strategy for PubMed.

1	Strabismus [Text Word] OR Strabismus [MeSH Terms]
2	Exotropia [Text Word] OR Exotropia [MeSH Terms]
3	Esotropia [Text Word] OR Esotropia [MeSH Terms]
4	Squint [Text Word] OR Squint [MeSH Terms]
5	Amblyopia [Text Word] OR Amblyopia [MeSH Terms]
6	Lazy Eye [Text Word] OR Lazy Eye [MeSH Terms]
7	Pediatric [Text Word] OR Pediatric [MeSH Terms]
8	Children [Text Word] OR Children [MeSH Terms]
9	Adolescent [Text Word] OR Adolescent [MeSH Terms]
10	Adult [Text Word] OR Adult [MeSH Terms]
11	Aged [Text Word] OR Aged [MeSH Terms]
12	Prevalence [Text Word] OR Prevalence [MeSH Terms]
13	Frequency [Text Word] OR Frequency [MeSH Terms]
14	Africa [Text Word] OR Africa [MeSH Terms]

enough details for relevant information to be extracted, the study was included. Hospital-based and retrospective studies were excluded from the current review.

Studies screening and appraisal

Studies were initially screened using their titles and abstracts. Full-text articles of studies that passed the initial screening were assessed to ensure all inclusion criteria were met. The following information were extracted from the full-text articles: authors' names, year of publication, sampling period, study location (country), sample size, study design, age range of participants, gender of participants, the diagnostic criteria for strabismus or amblyopia used and the prevalence and number of participants with strabismus or amblyopia. Screening of articles for eligibility was performed by two authors; disagreements about article eligibility were resolved by discussions with a third author.

A 10-item check list produced from the Downs and Black checklist and the Strengthening the Reporting of Observational studies in Epidemiology (STROBE) statement was used to rate and assess the quality of all the full-text articles included in the review. The maximum score a study could receive is 10. Assessment of study quality was conducted by two reviewers; disagreements were resolved by discussions with a third reviewer.

Data analysis

Statistical analysis was performed with R version 4.1.2 (R Core Team (2021). R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. <https://www.R-project.org/>). Individual study proportions and pooled estimates were assessed with a 95% confidence interval. The Freeman-Tukey double arcsine transformation was used before pooling to minimize the effects of studies with extremely high or low prevalence estimates on the overall pooled estimate.^{41–43} Heterogeneity between studies was assessed using the degree of inconsistency (I^2). The I^2 statistic provides an estimate of the percentage of heterogeneity across studies that is truly due to differences

between studies but not chance. Following Higgins et al. recommendation,⁴⁴ $I^2 > 50\%$ was defined as meaningful heterogeneity. The random effect model was used to analyze pooled effects when heterogeneity was meaningful. Subgroup analyses were conducted according to age, gender, study year, and type of amblyopia. To reflect the critical period in childhood where children are considered to develop amblyopia and due to the high variability in the age groupings used by the individual studies, ages were grouped broadly into two categories: ≤ 10 years and > 10 years. Meta-regression analysis was conducted to evaluate the influence of predetermined factors (such as age, sex, type of amblyopia and study year) on prevalence of amblyopia. The Egger's test was used to evaluate the presence of publication bias. For all statistical analysis, a p-value $< .05$ was considered statistically significant.

Results

Four hundred and seven (407) records were identified from database search. Two hundred and seventy-six (276) duplicates were identified and removed. One hundred and thirty-one (131) articles were then screened and eighty-three (83) of them were excluded based on their titles and abstracts, leaving forty-eight (48) articles. Five (5) additional articles were identified from the secondary literature search. A total of fifty-three (53) studies were assessed for eligibility. Twenty-nine (29) studies were excluded for various reasons as shown in **Figure 1**. The final number of studies included in the review was twenty-four (24). **Figure 1** shows the PRISMA flowchart outlining the steps in acquiring the articles for the systematic review and meta-analysis.

Studies included in this review are presented in **Table 2** (strabismus) and **Table 3** (amblyopia). Eight (8) studies (1 population-based and 7 school-based) reported on strabismus and 21 (3 population-based and 18 school-based) reported on amblyopia. The sample sizes of the included studies ranged from 352 to 8715. The lowest and highest age range of participants in the individual studies were ≤ 5 and




Figure 1. PRISMA flowchart outlining the steps in acquiring the articles for the systematic review and meta-analysis. 407 records were identified from database search; 276 duplicates were removed. The final number of studies included in the review was 24.

Table 2. Characteristics of studies included in systematic review and meta-analysis (Strabismus).

Author	Study location	Sampling period	Type of study	Sample size			Age of participants (years)	Method of strabismus assessment	Prevalence (%)			
				Male	Female	Total			strabismus	XT	ET	QOS
Elsahn, 2014 ⁴⁵	Egypt	n.s.	School-based	n.s.	n.s.	6029	6 – 12	n.s.	0.13	0.01	0.12	7
Giorgis and Bejiga, 2001 ⁴⁶	Ethiopia	2000	Population-based	889	1005	1894	≤5	Hirschberg's test	1.53	0.47	1.06	9
Azonobi et al, 2009 ⁴⁷	Nigeria	n.s.	School-based	3766	3522	7288	2 – 17	Hirschberg's test	0.44	0.14	0.30	
Adegbingbe et al, 2005 ³⁴	Nigeria	2003	School-based	802	905	1707	8 – 22	n.s.	1.29	n.s.	n.s.	8
Akpe et al, 2014 ³¹	Nigeria	n.s.	School-based	1024	1115	2139	5 – 19	Hirschberg's test/ Krimsky's test	0.89	0.33	0.56	8
Taha et al, 2015 ⁴⁸	Sudan	2010	School-based	n.s.	n.s.	768	5 – 14	n.s.	2.8	0.6	2.2	9
Wedner et al, 2000 ⁴⁹	Tanzania	1998	School-based	n.s.	n.s.	1386	7 – 19	Hirschberg's test/cover test	0.51			8
Ajaiyeoba et al, 2006 ²⁹	Nigeria	2002	School-based	504	640	1144	4 – 24	n.s.	0.26	0.17	0.09	9

n.s. – not stated XT – exotropia ET – esotropia.

QOS – quality of study.

Table 3. Characteristics of studies included in systematic review and meta-analysis (Amblyopia).

Author	Sampling period	Type of study	Sample size			Age of participants (years)	Prevalence (%)	Type of amblyopia	Diagnostic criteria	QOS
			Male	Female	Total					
Egypt										
Elsahn, 2014 ⁴⁵	n.s.	School-based	n.s.	n.s.	6029	6 – 12	0.13	n.s.	n.s.	7
Rashad et al, 2018 ⁵⁰	n.s.	School-based	193	159	352	8 – 12	1.98	Refractive (100%)	Unilateral: ≥ 2 -line difference in BCVA. Bilateral: BCVA worse than 6/12 in both eyes	8
Ethiopia										
Kedir and Girma, 2010 ⁵¹	2009	Population-based	308	262	570	11 – 15	0.35	Strabismic (100%)	n.s.	9
Giorgis and Bejiga, 2001 ⁴⁶	2000	Population-based	889	1005	1894	≤ 5	0.79	Strabismic (100%)	Unilateral: ≥ 2 -line difference in BCVA	9
Mehari and Yimer, 2013 ⁵²	2010/2011	School-based	2272	1966	4238	7 – 18	0.92	n.s.	n.s.	8
Haile et al, 2017 ⁵³	2015	School-based	582	705	1287	6 – 15	4.66	Refractive(76.6%), strabismic (6.7%), deprivational (8.3%)	Unilateral: ≥ 2 -line difference in BCVA. Bilateral: BCVA worse than 6/9 in both eyes	8
Hailu et al, 2020 ⁵⁴	2019	School-based	370	403	773	7 – 17	0.65	Refractive (100%)	n.s.	9
Ghana										
Ntim-Amponsah et al, 2007 ²⁶	2000/2001	School-based	n.s.	n.s.	975	6 – 22	0.2	Strabismic (100%)	n.s.	9
Kumah et al, 2013 ²⁵	2009	School-based	1143	1311	2454	12 – 15	0.45	n.s.	n.s.	9
Abu et al, 2015 ²⁸	n.s.	School-based	463	566	1029	9 – 22	0.32	n.s.	n.s.	7
Malawi										
Thom et al, 2017 ⁵⁵	2013	School-based	290	304	594	4 – 18	1.69	n.s.	n.s.	9
Nigeria										
Ajaiyeoba et al, 2006 ²⁹	2002	School-based	504	640	1144	4 – 24	0.26	Strabismic (100%)	n.s.	9
Akpe et al, 2015 ³¹	n.s.	School-based	1024	1115	2139	5 – 19	0.23	Refractive only (40%), anisometropic and strabismic (20%), meridional (20%)	Unilateral: ≥ 2 -line difference in BCVA. Bilateral: BCVA worse than 6/9 in both eyes	8
Adegbhingbe et al, 2005 ³⁴	2003	School-based	802	905	1707	8 – 22	0.7	n.s.	n.s.	8
Megbelayin, 2012 ³²	2010	School-based	535	640	1175	9 – 21	0.3	n.s.	Unilateral: ≥ 2 -line difference in BCVA. Bilateral: BCVA worse than 6/9 in both eyes	9
Ekpenyong et al, 2017 ³⁵	n.s.	School-based	993	1117	2110	6 – 17	0.28	n.s.	n.s.	7
Ikuomenisan et al, 2016 ³⁰	2014	School-based	899	803	1702	4 – 16	1.4	Refractive (58.23%), strabismic (20.83%), deprivational (20.83%)	Unilateral: ≥ 2 -line difference in BCVA. Bilateral: BCVA worse than 6/9 in both eyes	10
South Africa										
Naidoo et al, 2003 ⁵⁶	2002	Population-based	n.s.	n.s.	4890	5 – 15	0.29	n.s.	n.s.	9
Sudan										
Alrasheed et al, 2016 ⁵⁷	2014/2015	School-based	827	839	1678	6 – 15	0.36	n.s.	n.s.	8
Tanzania										
Wedner et al, 2000 ⁴⁹	1998	School-based	n.s.	n.s.	1386	7 – 19	0.22	n.s.	n.s.	8
Togo										
Kossi and Argudo, 2019 ⁵⁸	2017/2018	School-based	n.s.	n.s.	8715	11 – 16	0.38	Refractive (100%)	n.s.	8

n.s. – not stated BCVA – best corrected visual acuity QOS – quality of study.

4–24 years. Included studies were published from 1996 to 2019. Egger's test revealed no publication bias for studies on strabismus ($p = .06$) and amblyopia ($p = .08$).

Pooled prevalence estimates of strabismus

Overall prevalence estimate of strabismus in Africa was pooled from eight studies with a total sample size of 22,355. Strabismus prevalence reported by these studies was in the range 0.1% – 2.9% and overall pooled prevalence of strabismus was 0.8% (95% CI: 0.4% – 1.4%; $I^2 = 92.99\%$, $p < .01$). A forest plot for overall pooled strabismus prevalence is presented in Figure 2.

Overall prevalence of esotropia and exotropia were each pooled from six studies. Prevalence estimate for exotropia was 0.2% (95% CI: 0.1% – 0.5%; $I^2 = 83.93\%$, $p < .01$) and prevalence for esotropia was 0.5% (95% CI: 0.1% – 1.2%; $I^2 = 91.81\%$, $p < .01$) (Figure 3).

Pooled prevalence estimates of amblyopia

The overall prevalence estimate of amblyopia in Africa was pooled from 21 studies. Amblyopia prevalence reported by these studies was in the range 0.13% – 4.66%; overall pooled prevalence of amblyopia was 0.6% (95% CI: 0.3% – 0.9%; $I^2 = 90.5\%$, $p < .01$). Forest plot for pooled amblyopia prevalence is presented in Figure 4.

Gender and age group-based prevalence of amblyopia

Prevalence of amblyopia in males and females were each pooled from six studies and their respective

pooled prevalence estimates were 1.8% (95% CI: 0.7% – 3.3%; $I^2 = 86\%$, $p < .01$) and 1.3% (95% CI: 0.4% – 2.6%; $I^2 = 86\%$, $p < .01$) (Figure 5(a)). Meta-regression revealed no significant association between the prevalence of amblyopia and gender ($p = .752$). For analysis, the ages of participants in the included studies were grouped into ≤ 10 years and > 10 years. The prevalence of amblyopia among individuals ≤ 10 years was pooled from three studies; prevalence estimate was 1.7% (95% CI: 0.4% – 3.8%; $I^2 = 84.52\%$, $p < .01$). Estimated prevalence of amblyopia in individuals > 10 years (pooled from 5 studies) was 1.5% (95% CI: 0.2% – 3.8%; $I^2 = 89.53\%$, $p < .01$). Figure 5(b) represents the subgroup group analysis for age. There was no significant association between amblyopia and age ($p = .10$).

Prevalence of amblyopia based on study year

For analysis, studies were grouped based on data collection year – study performed after 2010 (> 2010) and in or before 2010 (≤ 2010). Prevalence estimates of amblyopia in studies published > 2010 (pooled from six studies) was 1.2% (95% CI: 0.3% – 1.2%; $I^2 = 93\%$, $p < .01$) and for studies published ≤ 2010 (pooled from 9 studies) was 0.4% (95% CI: 0.2% – 0.6%; $I^2 = 52\%$, $p = .05$). There was a significant association between prevalence of amblyopia and study year ($p = .006$). Figure 5(c) shows the subgroup group analysis for study year.

Prevalence of amblyopia based on amblyopia type

Subgroup analysis for type of amblyopia was performed when more than two studies reported type




Figure 2. Forest plot of studies reporting on strabismus in Africa. Overall estimate of strabismus prevalence in Africa was calculated using a random effects model. Prevalence of strabismus in Africa was estimated to be 0.8% (95% CI: 0.4% – 1.4%).




Figure 3. Forest plot of studies reporting on esotropia (a) and exotropia (b) in Africa. Estimated prevalence was calculated using a random effects model. Prevalence of esotropia and exotropia were 0.5% (95% CI: 0.1% – 1.2%) and 0.2% (95% CI: 0.1% – 0.5%), respectively.




Figure 4. Forest plot of studies reporting on amblyopia in Africa. Overall estimate of amblyopia prevalence in Africa was calculated using a random effects model. Prevalence of amblyopia in Africa was estimated to be 0.6% (95% CI: 0.3% – 0.9%).




Figure 5. Prevalence of amblyopia in Africa according to gender (a) age, (b) study year, (c) and (d) type of amblyopia. The diamond marks illustrate the pooled prevalence estimates, and the width of each diamond mark represents the confidence interval of the pooled estimate.

of amblyopia. Estimated prevalence of strabismic (pooled from six studies) and refractive amblyopia (pooled from five studies) were 0.4% (95% CI: 0.2% – 0.6%; $I^2 = 27\%$, $p = .23$) and 1.1% (95% CI: 0.2% – 2.5%; $I^2 = 95\%$, $p < .01$), respectively (Figure 5(d)). There was a significant association between prevalence of amblyopia and type of amblyopia ($p = .007$).

Discussion

The current meta-analysis included eight studies (one population-based & seven school-based) on strabismus and 21 studies (3 population-based & 18 school-based) on amblyopia. The overall prevalence of strabismus was estimated to be 0.8% (95% CI: 0.4% – 1.4%); exotropia was 0.2% (95%




Figure 5. (Continued).

CI: 0.1% – 0.5%) and esotropia was 0.5% (95% CI: 0.1% – 1.2%). Prevalence of amblyopia was 0.6% (95% CI: 0.3% – 0.9%); refractive and strabismic amblyopia were 1.1% (95% CI: 0.2% – 2.5%) and 0.4% (95% CI: 0.2% – 0.6%), respectively. Prevalence estimate of amblyopia

in males was 1.8% (95% CI: 0.7% – 3.3%) and in females was 1.3% (95% CI: 0.4% – 2.6%).

The estimated prevalence of strabismus and amblyopia in the current study is lower than prevalence estimated in other regions. Hashemi et al.^{5,19} estimated the global prevalence of strabismus

and amblyopia to be 1.93% and 1.75%, respectively, and also provided estimates for the different global regions according to the World Health Organization regional grouping as follows: American Regional Office (strabismus 2.86%; amblyopia 2.77%); Eastern Mediterranean Regional Office (strabismus 1.96%; amblyopia 1.54%); European Regional Office (strabismus 2.41%; amblyopia 3.67%); Western Pacific Regional Office (strabismus 2.51%; amblyopia 1.19%); and the prevalence of amblyopia in South-East Asia Regional Office was 2.74%. Fu et al.²² also estimated the global prevalence of amblyopia to be 1.44% with estimates in Europe, North America and Asia being 2.90%, 2.41% and 1.09%, respectively. Our current estimate of strabismus and amblyopia in Africa is comparatively lower than reported for the different regions in these global meta-analyses, but relatively higher than previously reported for Africa Regional Office (0.42% and 0.51% for strabismus and amblyopia, respectively) in the meta-analysis by Hashemi et al.^{5,19} The comparatively lower prevalence estimate in the global meta-analysis could be due to the limited number of studies reviewed by the authors and the year of study considering that amblyopia prevalence has tripled since 2010 and was significantly associated with study year in the current study. For instance, in the meta-analysis by Hashemi et al., only three and six studies (all published by 2016) were included in estimating the prevalence of strabismus and amblyopia, respectively, compared to 8 (strabismus) and 21 (amblyopia) studies used in the current study. All three studies on strabismus and five out of the six studies on amblyopia included in the Hashemi et al. study were included in the current meta-analysis. The current study therefore provides a relatively more representative estimate of strabismus and amblyopia prevalence in Africa. A possible reason for the comparatively lower prevalence of strabismus and amblyopia in the current study could be the inclusion of mainly school-based studies. Evidence suggests a relationship between

socioeconomic status and school enrollment; enrollment is lower in groups with lower socioeconomic status compared to those with higher socioeconomic status.^{59,60} Socioeconomic status has been shown to be a significant determinant of amblyopia prevalence; there is negative correlation between socioeconomic status and amblyopia prevalence.^{21,61} Considering that majority of studies included in our analysis were school-based, it is plausible that these studies may have sampled individuals of relatively higher socioeconomic status and lower amblyopia prevalence and hence, the prevalence reported in these studies may reflect the prevalence of amblyopia in school-going children but not the general population. An additional reason for the low prevalence of strabismus in Africa may be the survival rate of preterm infants. The survival rate of preterm babies in developing regions is extremely low. According to the World Health Organization, more than 90% of extremely preterm babies (less than 28 weeks) born in low-income countries die within the first few days of life; yet less than 10% of extremely preterm babies die in high-income settings.⁶² The lower prevalence of strabismus in our study may thus be explained by the low survival rate of preterm babies in Africa given that the prevalence of strabismus tends to be higher in preterm babies and can be as high as 42%.^{63,64}

There is conflicting data on the commonest strabismus type. While some studies have reported that esotropia occurs 3–5 times as often as exotropia in children,⁶⁵ other studies also report exotropia to be more common than esotropia.^{1,9,10} Although not significant, there was a higher prevalence of esotropia compared to exotropia in the current study. This is in contrast with global estimates where exotropia was the common strabismus type.⁵

In agreement with previous studies,^{22,66} the current study reported no gender difference in amblyopia prevalence in Africa. No significant difference in amblyopia prevalence was found between individuals ≤ 10 years and those > 10 years. Contrary to this, the meta-analysis by Hashemi et al.⁵ reported the

prevalence of amblyopia to be higher in adults compared with children. Amblyopia occurs due to disorders in the development of the visual pathway during the critical period of neural development in childhood, but its effect lasts throughout adult lifetime. Amblyopia has a better prognosis when it is detected and treated early, with treatment more effective in children.⁶⁷ Comprehensive eye care services are lacking in most regions of Africa,^{68,69} hence no effective and robust screening programs in many countries to help detect these early signs. This could result in significant undetected/undiagnosed amblyopia on the continent. Challenges within the general and eye health systems of many African countries such as low doctor-to-patient ratio, poor health care seeking behavior, lack of effective health-care policies, and other socioeconomic factors,^{70–72} means that even when the condition is detected, there are limited treatment options, and the cost involved for treatment may be a deterrent to affected individuals. Taken together, these challenges could prevent the early identification and management of strabismus and amblyopia. Accordingly, this highlights the need for and importance of comprehensive screening programs in Africa.

Amblyopia can present in different forms with distinct characteristics. Deprivation amblyopia has the greatest influence on visual acuity and other visual functions.⁷³ Refractive amblyopia on the other hand can be easily treated by correcting refractive errors using spectacle or contact lenses.^{74,75} In the current study, refractive amblyopia was the most common type of amblyopia with a prevalence of 1.1%, with type of amblyopia being significantly associated with amblyopia prevalence in Africa. Although other studies have reported similar findings,^{19,76,77} the peculiar issue of low spectacle coverage^{78,79} in several African countries may be a significant contributory factor for the higher prevalence of refractive amblyopia in Africa.

A major limitation of this review is that studies were from only eight out of the 54 countries in Africa, even though effort was made to include studies from across the continent. Despite including more studies compared to previous reviews, the lack of representativeness from several African countries means the findings from this study should be interpreted with caution and highlights the need for well-conducted epidemiological studies on strabismus and

amblyopia in many parts of Africa. Also, some of the studies included were not originally designed to assess prevalence of amblyopia or strabismus and as such might not have employed measures to prevent bias in reporting the prevalence. Another limitation is the lack of gender and age-specific prevalence of strabismus. There were not enough studies to pool the gender and age-specific prevalence of strabismus in Africa, highlighting the need for more studies, especially population-based studies on strabismus in Africa. Despite these limitations, the estimate from the current study still gives a good impression of the strabismus and amblyopia problem in Africa.

Conclusion

Strabismus and amblyopia are major causes of visual impairment in children and adults, and one of the main causes for public eye health concern. In adulthood, strabismus and amblyopia have an impact on lifestyle, wellbeing, and quality of life. Therefore, it is important to have information of the prevalence of strabismus and amblyopia, specifically in Africa, because the burden of strabismus and amblyopia is poorly studied in Africa continent. Despite the comparatively low prevalence, the current systematic review and meta-analysis highlights the need for more comprehensive epidemiological studies on strabismus and amblyopia in Africa. Information from these studies can help inform policy making and design and implementation of public health intervention programs.


Disclosure statement

No potential conflict of interest was reported by the authors.

Funding

The author(s) reported there is no funding associated with the work featured in this article.

ORCID

Prince Kwaku Akowuah, OD, PhD  <http://orcid.org/0000-0003-3657-6768>

Samuel Adade, OD, PhD  <http://orcid.org/0000-0003-4604-8651>

Ebenezer Owusu, OD  <http://orcid.org/0000-0003-1612-4350>
 Richard Donkor, OD, PhD  <http://orcid.org/0000-0001-5669-5380>
 Stephen Ankamah-Lomotey, OD, MSc  <http://orcid.org/0000-0003-4235-2891>
 Asafo Agyei Frimpong, OD  <http://orcid.org/0000-0002-7246-773X>
 Joseph Adjei-Anang, OD, MSc  <http://orcid.org/0000-0003-4296-5882>
 Emmanuel Kobia-Acquah, OD, MSc  <http://orcid.org/0000-0002-5955-7992>

References

- De Amorim Garcia CA, De Sousa AB, De Melo Mendonça MB, De Andrade LL, Oréface F. Prevalence of strabismus among students in Natal/RN - Brazil. *Arq Bras Oftalmol.* 2004;67:791–794. doi:10.1590/S0004-27492004000500018.
- Zhang XJ, Lau YH, Wang YM, et al. Prevalence of strabismus and its risk factors among school aged children: the Hong Kong Children Eye Study. *Sci Rep.* 2021 Dec 1;11(1). doi:10.1038/s41598-021-93131-w.
- Fieß A, Elflein HM, Urschitz MS, et al. Prevalence of strabismus and its impact on vision-related quality of life: results from the German population-based Gutenberg Health Study. *Ophthalmology.* 2020 Aug 1;127(8):1113–1122. doi:10.1016/j.ophtha.2020.02.026.
- He H, Fu J, Meng Z, Chen W, Li L, Zhao X. Prevalence and associated risk factors for childhood strabismus in Lhasa, Tibet, China: a cross-sectional, school-based study. *BMC Ophthalmol.* 2020 Dec 1;20(1):1–9. doi:10.1186/s12886-020-01732-2.
- Hashemi H, Pakzad R, Heydarian S, et al. Global and regional prevalence of strabismus: a comprehensive systematic review and meta-analysis. *Strabismus.* 2019;27(2):54–65. doi:10.1080/09273972.2019.1604773.
- Lança C, Serra H, Prista J. Strabismus, visual acuity, and uncorrected refractive error in portuguese children aged 6 to 11 years. *Strabismus* [Internet]. 2014;22(3):115–119. <https://pubmed.ncbi.nlm.nih.gov/25019452/> [Accessed 2022 Oct 29].
- Grönlund MA, Andersson S, Aring E, Hård AL, Hellström A. Ophthalmological findings in a sample of Swedish children aged 4-15 years. *Acta Ophthalmol Scand* [Internet]. 2006 Apr;84(2):169–176. <https://pubmed.ncbi.nlm.nih.gov/16637831/> [Accessed 2022 Oct 29].
- Chia A, Dirani M, Chan YH, et al. Prevalence of amblyopia and strabismus in young singaporean chinese children. *Invest Ophthalmol Vis Sci.* 2010;51:3411–3417. doi:10.1167/iovs.09-4461.
- Chen X, Fu Z, Yu J, et al. Prevalence of amblyopia and strabismus in Eastern China: results from screening of preschool children aged 36–72 months. *British J Ophthalmol.* 2016;100:515–519. doi:10.1136/bjophthalmol-2015-306999.
- Group M-ePEDS. Prevalence of amblyopia and strabismus in African American and Hispanic children ages 6 to 72 months. The multi-ethnic pediatric eye disease study. *Ophthalmology.* 2008;115:1229–1236. doi:10.1016/j.ophtha.2007.08.001.
- Archer SM, Musch DC, Wren PA, Guire KE, Del Monte MA. Social and emotional impact of strabismus surgery on quality of life in children. *J AAPOS.* 2005;9:148–151. doi:10.1016/j.jaapos.2004.12.006.
- McBain HB, Au CK, Hancox J, et al. The impact of strabismus on quality of life in adults with and without diplopia: a systematic review. *Surv Ophthalmol.* 2014 Mar 1;59(2):185–191. doi:10.1016/j.survophthal.2013.04.001.
- Chang MY, Velez FG, Demer JL, Isenberg SJ, Coleman AL, Pineles SL. Quality of life in adults with strabismus. *Am J Ophthalmol* [Internet]. 2015 Mar 1;159(3):539–544.e2. <https://pubmed.ncbi.nlm.nih.gov/25498355/> [Accessed 2022 Oct 29].
- Buffenn AN. The impact of strabismus on psychosocial health and quality of life: a systematic review. *Surv Ophthalmol.* 2021 Nov 1;66(6):1051–1064. doi:10.1016/j.survophthal.2021.03.005.
- Levi DM, Knill DC, Bavelier D. Stereopsis and amblyopia: a mini-review. *Vision Res.* 2015;114:17–30. doi:10.1016/j.visres.2015.01.002.
- Holmes JM, Clarke MP. Amblyopia. *Lancet.* 2006;367(9519):1343–1351. doi:10.1016/S0140-6736(06)68581-4.
- Maurer D, McKee SP. Classification and diversity of amblyopia. *Vis Neurosci.* 2018;35:E012. doi:10.1017/S0952523817000190.
- Asper L, Crewther D, Crewther SG. Strabismic amblyopia: part 1: psychophysics. *Clin Exp Optometry.* 2000;83(2):49–58. doi:10.1111/j.1444-0938.2000.tb04892.x.
- Hashemi H, Pakzad R, Yekta A, et al. Global and regional estimates of prevalence of amblyopia: a systematic review and meta-analysis. *Strabismus.* 2018;26(4):168–183. doi:10.1080/09273972.2018.1500618.
- Schmucker C, Grosselfinger R, Riemsma R, et al. Effectiveness of screening preschool children for amblyopia: a systematic review. *BMC Ophthalmol.* 2009;9(1). doi:10.1186/1471-2415-9-3.
- Faghihi M, Hashemi H, Nabovati P, et al. The prevalence of amblyopia and its determinants in a population-based study. *Strabismus.* 2017;25(4):176–183. doi:10.1080/09273972.2017.1391849.
- Fu Z, Hong H, Su Z, Lou B, Pan CW, Liu H. Global prevalence of amblyopia and disease burden projections through 2040: a systematic review and meta-analysis. *British J Ophthalmol.* 2020;104:1164–1170. doi:10.1136/bjophthalmol-2019-314759.
- Attebo K, Mitchell P, Cumming R, Smith W, Jolly N, Sparkes R. Prevalence and cause of amblyopia in an adult population. *Ophthalmology.* 1998;105:154–159. doi:10.1016/S0161-6420(98)91862-0.

24. Webber AL, Wood J. Amblyopia: prevalence, natural history, functional effects and treatment. *Clin Exp Optometry*. 2005;88:365–375. doi:10.1111/j.1444-0938.2005.tb05102.x.
25. Kumah BD, Ebri A, Abdul-Kabir M, et al. Refractive error and visual impairment in private school children in Ghana. *Optometry and Vision Sci*. 2013;90:1456–1461. doi:10.1097/OPX.0000000000000099.
26. Ntim-Amponsah CT, Ofosu-Amaah S. Prevalence of refractive error and other eye diseases in school children in the Greater Accra region of Ghana. *J Pediatric Ophthalmol Strabismus*. 2007;44:294–297.
27. Abdul-Kabir M, Abdul-Sadik A, Ansah DO, et al. Prevalence of anisometropia, strabismus and amblyopia among first year optometry students in Kwame Nkrumah University of Science and Technology, Ghana. *Mathews J Ophthalmol*. 2017;2:018. doi:10.15226/2474-9249/2/2/00126.
28. Abu EK, Yeboah AA, Ocansey S, Kyei S, Abokyi S. Epidemiology of ocular disorders and visual impairment among school pupils in the Cape Coast Metropolis, Ghana. *Br J Vis Impair*. 2015;33:45–53. doi:10.1177/0264619614561690.
29. Ajaiyeoba AI, Isawumi MA, Adeoye AO, Oluleye TS. Prevalence and causes of eye diseases amongst students in south-western Nigeria. *Ann Afr Med*. 2006;5:197–203.
30. Ikuomenisan S, Musa K, Aribaba O, Onakoya A. Prevalence and pattern of amblyopia among primary school pupils in Kosofe town, Lagos state, Nigeria. *Niger Postgrad Med J*. 2016;23:196–201. doi:10.4103/1117-1936.196261.
31. Akpe BA, Dawodu OA, Abadom EG. Prevalence and pattern of strabismus in primary school pupils in Benin city, Nigeria. *Niger J Ophthalmol*. 2014;1:38–43. doi:10.4103/0189-9171.142755.
32. Megbelayin EO. Prevalence of amblyopia among secondary school students in Calabar, south-south Nigeria. *Niger J Med*. 2012;21:407–411.
33. Akpe B, Abadom E, Omoti E. Prevalence of amblyopia in primary school pupils in Benin City, Edo State, Nigeria. *Afr J Med Health Sci*. 2015;14:110–114. doi:10.4103/2384-5589.170174.
34. Screening of adolescents for eye diseases in Nigerian High Schools. *Ghana Med J*. 2005;39:138–142.
35. Ekpenyong BN, Naidoo K, Ahaiwe K, et al. Visual status and prevalence of eye disorders among school-age children in southern Nigeria. *Afr Vis Eye Health*. 2017;76:a377. doi:10.4102/aveh.v76i1.377.
36. Alarape AT, Ulaikere M, Okoye O, et al. Burden and spectrum of amblyopia in a pediatric hospital population southwest Nigeria. *Ann Med Health Sci Res*. 2017;7:60–66.
37. World Health Organisation. Causes of blindness and visual impairment. 2012.
38. Sacharowitz HS. Visual impairment in South Africa: achievements and challenges. *Afr Vis Eye Health*. 2005;64(4). doi:10.4102/aveh.v64i4.239.
39. Muhammad N, Mansur R, Dantani A, Elhassan E, Isiyaku S. Prevalence and causes of blindness and visual impairment in Sokoto State, Nigeria: baseline data for Vision 2020: the right to sight eye care programme. *Middle East Afr J Ophthalmol*. 2011;18(2):123. doi:10.4103/0974-9233.80700.
40. Moher D, Liberati A, Tetzlaff J, Altman DG, Altman D, Antes G, The PRISMA Group. Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *PLoS Med*. 2009;7:e1000097. doi:10.1371/journal.pmed.1000097.
41. Doi SA, Xu C. The Freeman-Tukey double arcsine transformation for the meta-analysis of proportions: recent criticisms were seriously misleading. *J Evid Based Med*. 2021 Dec 1;14(4):259–261. doi:10.1111/jebm.12445.
42. Freeman MF, Tukey JW Transformations related to the angular and the square root. *Ann Math Stat*. 1950 Dec 1;21(4):607–11. doi:10.1214/aoms/1177729756.
43. Akowuah PK, Kobia-Acquah E, Donkor R, Adjei-Anang J, Ankamah-Lomotey S. Keratoconus in Africa: a systematic review and meta-analysis. *Ophthalmic Physiol Optics*. 2021 Jul;41(4):736–747. doi:10.1111/opo.12825.
44. Higgins JPT, Thompson SG, Deeks JJ, Altman DG. Measuring inconsistency in meta-analyses. *Br Med J*. 2003;327:557–560. doi:10.1136/bmj.327.7414.557.
45. Elsahn M. International Vision screening: results from Alexandria, Egypt. *Theor Chem Acc*. 2014;2:137–141. doi:10.1007/s40135-014-0055-3.
46. Giorgis AT, Bejiga A. Prevalence of strabismus among pre-school children community in Butajira Town. *Ethiop J Health Dev*. 2001;15:125–130. doi:10.4314/ejhd.v15i2.9886.
47. Azonobi I, Adido J, Olatunji F, Bello A, Mahmoud A. Risk factors of strabismus in southwestern Nigeria. *Pak J Ophthalmol*. 2009;25:129–132.
48. Taha A, Ibrahim S. Prevalence of manifest horizontal strabismus among basic school children in Khartoum City, Sudan. *Sudan J Ophthalmol*. 2015;7:53–57. doi:10.4103/1858-540X.169437.
49. Wedner SH, Ross DA, Balira R, Kaji L, Foster A. Prevalence of eye diseases in primary school children in a rural area of Tanzania. *British J Ophthalmol*. 2000;84:1291–1297. doi:10.1136/bjo.84.11.1291.
50. Rashad MA, Abd Elaziz KM, Fawzy SM, Abdel Latif AAM, Abdel Latif MAM. Screening of primary school children for amblyopia and amblyogenic factors in Central Cairo. *Egypt J Ophthalmol*. 2018;2018:6. doi:10.1155/2018/8425319.
51. Kedir J, Girma A. Prevalence of refractive error and visual impairment among rural school-age children of Goro District, Gurage Zone, Ethiopia. *Ethiop J Health Sci*. 2010;2:353–358.
52. Mehari ZA, Yimer AW. Prevalence of refractive errors among school children in rural central Ethiopia. *Clin*

- Exp Optometry*. 2013;96:65–69. doi:10.1111/j.1444-0938.2012.00762.x.
53. Haile W, Ayanaw T, Destaye S. Prevalence and types of amblyopia among primary school children in Gondar town, Northwest Ethiopia. *Open Access J Ophthalmol*. 2017;2:000124. doi:10.23880/OAJO-16000124.
 54. Hailu Y, Hiko D, Shaweno T. Prevalence of visual impairment and associated factors among primary school children in Addis Ababa, Central Ethiopia. *Clin Ophthalmol*. 2020;14:767–774. doi:10.2147/OPHTH.S245675.
 55. Thom L, Jogessar S, McGowan SL, Lawless F. The prevalence and causes of decreased visual acuity – a study based on vision screening conducted at Erukweni and Mzuzu Foundation Primary Schools, Malawi. *Clin Optom*. 2017;9:1–10. doi:10.2147/OPTO.S110097.
 56. Naidoo KS, Raghunandan A, Mashige KP, et al. Refractive error and visual impairment in African children in South Africa. *Invest Ophthalmol Vis Sci*. 2003;44:3764–3770. doi:10.1167/iovs.03-0283.
 57. Alrasheed SH, Naidoo KS, Clarke-Farr PC. Prevalence of visual impairment and refractive error in school-aged children in South Darfur State of Sudan. *Afr Vis Eye Health*. 2016;75:a355. doi:10.4102/aveh.v75i1.355.
 58. Kossi B, Amah H. Refractive errors and amblyopia in scholar's population in Togo. *JOJ Ophthalmol*. 2019;7:555714. doi:10.19080/JOJO.2019.07.555714.
 59. Declercq K, Verboven F. Socio-economic status and enrollment in higher education: do costs matter? *Educ Econ* [Internet]. 2015 Sep 3;23(5):532–556. <https://www.tandfonline.com/doi/abs/10.1080/09645292.2015.1047822> [Accessed 2022 Oct 29].
 60. Ara A, Abdul Aziz S. Impact of socio-economic status on the enrollment of girl students in public sector schools of district West Karachi. *Educ Res* [Internet]. 2013;2(3):32–38. www.journals.savap.org.pk/www.savap.org.pk [Accessed 2022 Oct 29]
 61. Nitzan I, Bez M, Megreli J, et al. Socio-demographic disparities in amblyopia prevalence among 1.5 million adolescents. *Eur J Public Health* [Internet]. 2021 Dec 1;31(6):1211–1217. <https://pubmed.ncbi.nlm.nih.gov/34518882/> [Accessed 2022 Oct 29].
 62. Preterm birth. [Internet]. <https://www.who.int/news-room/fact-sheets/detail/preterm-birth> [Accessed 2022 Oct 29].
 63. Bremer DL, Palmer EA, Fellows RR, et al. Strabismus in premature infants in the first year of life. Cryotherapy for Retinopathy of Prematurity Cooperative Group. *Arch Ophthalmol*. [Internet]. 1998;116(3):329–333. <https://pubmed.ncbi.nlm.nih.gov/9514486/> [Accessed 2022 Oct 29].
 64. Fieß A, Kölb-Keerl R, Schuster AK, et al. Prevalence and associated factors of strabismus in former preterm and full-term infants between 4 and 10 years of age. *BMC Ophthalmol* [Internet]. 2017 Dec 2;17(1). /pmc/articles/PMC5712131/. [Accessed 2022 Oct 29].
 65. Donnelly U, Stewart N, Hollinger M. Prevalence and outcomes of childhood visual disorders. *Ophthalmic Epidemiol*. 2005;12:243–250. doi:10.1080/09286580590967772.
 66. Pan C, Chen X, Zhu H, et al. School-based assessment of amblyopia and strabismus among multiethnic children in rural China. *Sci Rep*. 2017;7:13410. doi:10.1038/s41598-017-13926-8.
 67. Tailor V, Bossi M, Greenwood JA, Dahlmann-Noor A. Childhood amblyopia: current management and new trends. *Br Med Bull*. 2016 Sep 1;119(1):75–86. doi:10.1093/bmb/ldw030.
 68. Bechange S, Jolley E, Virendrakumar B, Pente V, Milgate J, Schmidt E. Strengths and weaknesses of eye care services in sub-Saharan Africa: a meta-synthesis of eye health system assessments. *BMC Health Serv Res*. 2020 May 6;20(1):1–8. doi:10.1186/s12913-020-05279-2.
 69. Graham R. Facing the crisis in human resources for eye health in sub-Saharan Africa. *Community Eye Health*. 2017;30:85.
 70. De-Graft Aikins A, Marks DF. Health, disease and healthcare in Africa. *J Health Psychol*. 2007;12:387–402. doi:10.1177/1359105307076228.
 71. Azevedo MJ. The State of Health System(s) in Africa: Challenges and Opportunities. In: *Historical Perspectives on the State of Health and Health Systems in Africa*. Vol. 2. 2017:1–73.
 72. Bonfrer I, Van De Poel E, Grimm M, Van Doorslaer E. Does the distribution of healthcare utilization match needs in Africa? *Health Policy Plan*. 2014;29:921–937. doi:10.1093/heapol/czt074.
 73. Hamm L, Chen Z, Li J, et al. Interocular suppression in children with deprivation amblyopia. *Vision Res*. 2017;133:112–120. doi:10.1016/j.visres.2017.01.004.
 74. Cotter SA. Treatment of anisometropic amblyopia in children with refractive correction. *Ophthalmology*. 2006;113:895–903. doi:10.1016/j.ophtha.2006.01.068.
 75. Steele AL, Bradfield YS, Kushner BJ, France TD, Struck MC, Gangnon RE. Successful treatment of anisometropic amblyopia with spectacles alone. *J AAPOS*. 2006;10:37–43. doi:10.1016/j.jaapos.2005.08.003.
 76. Simons K. Amblyopia characterization, treatment, and prophylaxis. *Surv Ophthalmol*. 2005;50:123–166. doi:10.1016/j.survophthal.2004.12.005.
 77. Jamali P, Fotouhi A, Hashemi H, Younesian M, Jafari A. Refractive errors and amblyopia in children entering school: Shahrood, Iran. *Optometry and Vision Sci*. 2009;86:364–369. doi:10.1097/OPX.0b013e3181993f42.
 78. Chan VF, Mebrahtu G, Ramson P, Wepo M, Naidoo KS. Prevalence of refractive error and spectacle coverage in Zoba Ma'ekel Eritrea: a rapid assessment of refractive error. *Ophthalmic Epidemiol*. 2013 Jun;20(3):131–137. doi:10.3109/09286586.2013.783082.
 79. Mashayo ER, Chan VF, Ramson P, Chinanayi F, Naidoo KS. Prevalence of refractive error, presbyopia and spectacle coverage in Kahama District, Tanzania: a rapid assessment of refractive error. *Clin Exp Optometry*. 2015 Jan 1;98(1):58–64. doi:10.1111/cxo.12207.