

2003

The Tenderisation of Shin Beef Using a Citrus Juice Marinade

Roisin Burke

Technological University Dublin, roisin.burke@tudublin.ie

Frank Monahan

University College Dublin, Ireland, frank.monahan@ucd.ie

Follow this and additional works at: <https://arrow.tudublin.ie/tfschafart>

 Part of the [Agribusiness Commons](#), [Arts and Humanities Commons](#), and the [Food Studies Commons](#)

Recommended Citation

Burke, R.M. and Monahan, F.J. (2003). The tenderisation of shin beef using a citrus juice marinade. *Meat Science*, 63 (2), pp.161-168. doi:10.1016/S0309-1740(02)00062-1

This Article is brought to you for free and open access by the School of Culinary Arts and Food Technology at ARROW@TU Dublin. It has been accepted for inclusion in Articles by an authorized administrator of ARROW@TU Dublin. For more information, please contact arrow.admin@tudublin.ie, aisling.coyne@tudublin.ie, vera.kilshaw@tudublin.ie.



2003-2

The tenderisation of shin beef using a citrus juice marinade

Roisin Burke

Frank Monahan

Follow this and additional works at: <https://arrow.dit.ie/aaconmusart>



Part of the [Food Science Commons](#)

This Article is brought to you for free and open access by the Conservatory of Music and Drama at ARROW@DIT. It has been accepted for inclusion in Articles by an authorized administrator of ARROW@DIT. For more information, please contact yvonne.desmond@dit.ie, arrow.admin@dit.ie, brian.widdis@dit.ie.



The tenderisation of shin beef using a citrus juice marinade

Author links open overlay panel [R.M.Burke](#) · [F.J.Monahan](#)
Show more

[https://doi.org/10.1016/S0309-1740\(02\)00062-1](https://doi.org/10.1016/S0309-1740(02)00062-1) [Get rights and content](#)

Abstract

The effectiveness of organic acids (acetic, citric, lactic) and a citrus juice marinade as tenderising agents in shin beef muscle was investigated. At 0.2 M, citric acid was more effective as a tenderising agent than acetic or lactic acid. Immersion of shin beef strips in citric acid (0–0.05 M) showed that a significant tenderising effect was obtained above a concentration of 0.013 M. When shin beef strips were immersed in the citrus juice marinade (31% orange juice, 31% lemon juice, 38% distilled water) mean pH decreased from 5.7 to 3.1 and mean sample weight increased by ~65%. The mean Warner–Bratzler shear force value decreased from 178 to 44 N cm⁻² following marination while mean sensory analysis scores for tenderness and juiciness increased following marination. A mean total collagen content of 1.4 g/100 g was recorded in shin beef of which 9% was soluble in unmarinated samples and 29% was soluble in marinated samples. The results indicated that the tenderisation of beef samples using a citrus juice marinade could be attributed to marinade uptake by muscle proteins and also to solubilisation of collagen.