

2020

## Effects of Cold Atmospheric Plasma on Mackerel Lipid and Protein Oxidation During Storage

Juan M. Pérez-Andrés  
*Technological University Dublin*

María de Alba  
*Food Safety, Teagasc Food Research Centre, Dublin, Ireland*

Sabine M Harrison  
*University College Dublin*

*See next page for additional authors*

Follow this and additional works at: <https://arrow.tudublin.ie/schfsehart>



Part of the [Food Biotechnology Commons](#), [Food Chemistry Commons](#), [Food Microbiology Commons](#), and the [Food Processing Commons](#)

### Recommended Citation

Perez J.M. et al. (2020) Effects of Cold Atmospheric Plasma on Mackerel Lipid and Protein Oxidation During Storage, *LWT - Food Science and Technology* 118 (2020) 108697. DOI:10.1016/j.lwt.2019.108697

This Article is brought to you for free and open access by the School of Food Science and Environmental Health at ARROW@TU Dublin. It has been accepted for inclusion in Articles by an authorized administrator of ARROW@TU Dublin. For more information, please contact [arrow.admin@tudublin.ie](mailto:arrow.admin@tudublin.ie), [aisling.coyne@tudublin.ie](mailto:aisling.coyne@tudublin.ie).



This work is licensed under a [Creative Commons Attribution-Noncommercial-Share Alike 3.0 License](#)

---

**Authors**

Juan M. Pérez-Andrés, María de Alba, Sabine M Harrison, Nigel P. Brunton, Patrick Cullen, and Brijesh K. Tiwari



## Effects of cold atmospheric plasma on mackerel lipid and protein oxidation during storage

Juan M. Pérez-Andrés<sup>a,b</sup>, María de Alba<sup>c,\*</sup>, Sabine M. Harrison<sup>f</sup>, Nigel P. Brunton<sup>f</sup>, P.J. Cullen<sup>b,d,e</sup>, Brijesh K. Tiwari<sup>a</sup>

<sup>a</sup> Food Chemistry and Technology, Teagasc Food Research Centre, Dublin, 15, Ireland

<sup>b</sup> BioPlasma Research Group, School of Food Science and Environmental Health, Technological University Dublin, Cathal Brugha Street, Dublin, 1, Ireland

<sup>c</sup> Food Safety, Teagasc Food Research Centre, Dublin, Ireland

<sup>d</sup> School of Chemical and Biomolecular Engineering, University of Sydney, Sydney, Australia

<sup>e</sup> Centre for Advanced Food Enjinomics, The University of Sydney, Sydney, 2006, Ireland

<sup>f</sup> UCD School of Agriculture & Food Science, University College Dublin, Dublin, Ireland

### ARTICLE INFO

#### Keywords:

Cold atmospheric plasma  
Mackerel  
Lipid oxidation  
Protein oxidation

### ABSTRACT

The present study investigated the effects of cold atmospheric plasma on the shelf-life stability of lipids and proteins of commercially packaged mackerel fillets. The results showed no significant effects on lipid oxidation between the control samples and those treated at 80 kV for extended treatment times of 5 min using an in-package plasma system. In addition, no significant changes ( $p > 0.05$ ) were found in the fatty acid composition or nutritional quality indices after treatment. Finally, the formation of carbonyls was accelerated by plasma treatment at the end of the storage period for 4 °C and 8 °C. These results suggest that cold atmospheric plasma technologies are potentially suitable as processing technologies for the fish industry.

### 1. Introduction

Fish and seafood products are some of the most common foods in our diet. Mackerel is a popular fish in Europe with more than 800 million tonnes consumed per year (FAO, 2018). It is considered a healthy diet option due to its high content of vitamins, high-quality protein as well as several other essential nutrients such as polyunsaturated fatty acids (Domingo, 2016; Swanson, Block, & Mousa, 2012). In addition, fatty acids present in fish are known to be an important source of essential omega-3 polyunsaturated fatty acids (n-3 PUFA) such as eicosapentaenoic acid (EPA) and docosahexaenoic acid (DHA). These fatty acids are necessary for functions in our metabolism and their intake through the diet is important because humans cannot synthesise them. Numerous studies have highlighted the benefits of EPA and DHA (Shahidi & Ambigaipalan, 2018; Swanson et al., 2012). For example, they are essential for fetal development, and supplementation during pregnancy has also been linked to decreased immune responses in infants including decreased incidence of allergies. The supplementation of both, EPA and DHA, has been reported to lead to optimal pregnancy length (Swanson et al., 2012). Kumar, Mastana, and Lindley (2016) suggested a diet rich in these fatty acids for patients with various chronic inflammatory diseases, like asthma, due to their anti-

inflammatory properties. In addition, they have been reported to play a role in avoiding coronary heart diseases such as atherosclerosis and peripheral arterial disease (Lee, O'keefe, Lavie, & Harris, 2009), as well as Alzheimer's and dementia (Kröger & Laforce, 2016).

Despite all the benefits associated with the intake of fish products, there are challenges facing the industry. Fish quality starts to deteriorate post catch and during storage with spoilage microorganisms the primary factor causing fish spoilage (Gram & Dalgaard, 2002). In this regard, it is paramount to find effective means to ensure the safety and quality of fish products. The food industry is increasingly seeking new processing technologies capable of inactivating microorganisms whilst maintaining food quality. A particular focus has been on non-thermal processing technologies, which are effective at ambient or sub-lethal temperatures (i.e. temperatures below 50 °C) such as high hydrostatic pressure, pulsed electric field, ultrasound processing and more recently cold plasma (Jermann, Koutchma, Margas, Leadley, & Ros-Polski, 2015). These technologies have been applied to inactivate microorganisms such as viruses, spores, yeasts, bacteria and biofilms (Arvanitoyannis, Kotsanopoulos, & Savva, 2017; Daryaei, Yousef, & Balasubramaniam, 2016, pp. 271–294; Saldana, Alvarez, Condon, & Raso, 2014). Numerous studies have examined their potential for the decontamination of fish products, including high pressure treatment of

\* Corresponding author. Teagasc Food Research Centre, Ashtown, Dublin, 15, Ireland.

E-mail address: [Maria.Ortega@teagasc.ie](mailto:Maria.Ortega@teagasc.ie) (M. de Alba).

<https://doi.org/10.1016/j.lwt.2019.108697>

Received 15 March 2019; Received in revised form 16 September 2019; Accepted 25 September 2019

Available online 27 September 2019

0023-6438/ © 2019 Elsevier Ltd. All rights reserved.

mackerel, cod, and salmon (De Alba et al., 2019; Rode & Hovda, 2016), sea bass (Teixeira et al., 2014), herring and haddock (Karim et al., 2011) and smoked cod (Montiel, De Alba, Bravo, Gaya, & Medina, 2012). Ultrasound has been used to reduce the microbiological counts in mackerel, cod, hake and salmon fillets (Pedrós-Garrido et al., 2017).

The principal advantage of such technologies is their operation at low temperatures, thus minimising the negative effects often found with thermal processing (Awuah, Ramaswamy, & Economides, 2007). Nevertheless, they too can also induce chemical/biochemical changes in food constituents, affecting the quality of the food product (Pérez-Andrés, Charoux, Cullen, & Tiwari, 2018).

Cold atmospheric plasma treatment can induce a wide array of active species including ozone, carbon and nitrogen oxides, free radicals, as well as negative and positive ions depending on the different gases used (Lu et al., 2016). Plasma for food applications is still in its infancy but has demonstrated significant potential for food decontamination and associated increases of shelf-life (Bourke, Ziuzina, Boehm, Cullen, & Keener, 2018). A growing literature is addressing the unknowns surrounding the efficacy and mechanisms of action for microbial inactivation (Liao et al., 2017). The effect of plasma on microbial inactivation has been investigated in different food substrates, including orange juice (Xu, Garner, Tao, & Keener, 2017), fresh and dried nut samples (*Juglans regia* L.) (Amini & Ghoranneviss, 2016), ground nuts (Devi, Thirumdas, Sarangapani, Deshmukh, & Annapure, 2017), strawberries (Misra et al., 2014) and blueberries (Lacombe et al., 2017).

Cold atmospheric plasma has also been shown to increase the shelf-life of meat products (Xiang et al., 2018), including chicken breasts (Wang, Zhuang, Hinton, & Zhang, 2016), beef jerky (Yong et al., 2017) and pork loin (Yong et al., 2017). Moreover, fish products, like dried filefish fillets (Park & Ha, 2015) and mackerel (Albertos et al., 2017) have been also tested.

The radical species present in the plasma are responsible for microbial inactivation although they may also lead to chemical reactions affecting the quality of the food product (Kim, Jayasena, Yong, & Jo, 2016, pp. 273–291; Misra, 2016, pp. 253–271; Pérez-Andrés et al., 2018). The challenge for all effective preservation techniques is to retain the sensory and quality attributes of the food while maximising shelf-life and safety profiles. Scale-up of plasma technology is another challenge which has recently been discussed by Cullen et al. (2018). Given the array of radical species found with atmospheric plasma and the wide range of time-scales of their reactions, there are many unknowns surrounding the chemical reactions induced from plasma treatment of foods (Scholtz, Pazlarova, Souskova, Khun, & Julak, 2015).

The goal of this study was to contribute to the understanding of the effects of cold plasma on the lipids and proteins of mackerel during storage and to support scale-up approaches for this technology. For that reason, mackerel fillets were packaged and plasma treated in food packaging trays typically employed by the industry. Key quality markers were monitored for different storage temperatures and time. The impact of cold atmospheric plasma on the stability of mackerel fatty acids, such as EPA, DHA, PUFA, monounsaturated fatty acids (MUFA), saturated fatty acids (SFA), as well as nutritional quality indices and lipid and protein oxidation were investigated.

## 2. Material and methods

### 2.1. Chemicals and reagents

Trichloroacetic acid, HEPES (4-(2-Hydroxyethyl)piperazine-1-ethanesulfonic acid, N-(2-Hydroxyethyl)piperazine-N'-(2-ethanesulfonic acid)), EGTA (Egtazic acid), disodium sulphate, hydrochloric acid, potassium hydroxide, 1,1,3,3-tetraethoxypropane, 2-thiobarbituric acid, methanol, heptane, Supelco-37 FAME standard, tricosanoic acid methyl ester, as well as acetyl chloride, sucrose, mannitol and potassium chloride were purchased from Sigma-Aldrich (Arklow, Co. Wicklow, Ireland). N-pentane as well as sodium chloride was purchased from

Fisher Scientific (Dublin, Ireland). Ultra-pure water ( $18.2 \text{ M}\Omega \text{ cm}^{-1}$ ) was generated in-house using a Millipore water purification system (Millipore, Cork, Ireland). All chemical were GC grade.

### 2.2. Sample preparation

Fresh mackerel was purchased at a local fish monger in Dublin in September 2016. Each mackerel was divided into two fillets and each fillet, with skin attached facing down, was packaged individually (Ilpra Foodpack VG 400 Packaging Machine, Italy) in a black amorphous polyethylene terephthalate (APET/PE) pack ( $195 \times 155 \times 37 \text{ mm}$ ), and sealed under atmospheric air conditions with a low oxygen permeable barrier polyvinyl-chloride film ( $3 \text{ cm}^3/\text{m}^2/24 \text{ h}$  at Standard Temperature and Pressure (STP); Versatile Packaging, Ireland), to mirror common commercial practice.

### 2.3. Plasma treatment

An in-house dielectric barrier discharge atmospheric plasma system was employed, which was described previously (Pankaj, Misra, & Cullen, 2013). Each packaged tray was located between two circular aluminium electrodes (outer diameter = 158 mm) separated by two polypropylene (PP) dielectric layers (2 mm thickness). The distance between the dielectric layers was the height of the tray employed (37 mm). Five minutes of treatment was performed at a discharge voltage of 80 kV Root Mean Square (RMS). The choice of these conditions was based on a recent study in the area carried out by our group (Albertos et al., 2017) which showed this treatment as the most effective one to reduce total aerobic psychrotrophic bacteria and *Pseudomonas* counts in in-packaged plasma treated mackerel fillets with no impact on colour or TBARs values. Both the control and plasma samples were kept at 4 °C and held for 24 h to allow the radical species formed to interact with the food following the method of Ziuzina, Patil, Cullen, Keener, and Bourke (2013). After that, samples were randomly divided into three different batches and stored at three different temperatures: 4 °C, 8 °C and –20 °C. Samples from each temperature were collected after 2, 5 and 7 days and in the case of the –20 °C batch also after 2 weeks. For each day, samples were homogenised using a blender (Robot Coupe-R301 Ultra; Nisbets, Cork, Ireland) and were stored under vacuum at –80 °C until the day of the analysis. All the treatments were carried out in duplicate.

### 2.4. Lipid oxidation

Lipid oxidation was quantified using the TBARs method according to Maraschiello, Sárraga, and Garcia Regueiro (1999), which is based on the detection of malondialdehyde (MDA), a by-product of peroxidation of polyunsaturated fatty acids and esters, associated with off-flavours and off-aromas in meat products (Ghani, Barril, Bedgood, & Prenzler, 2017). Briefly, 1 g of blended mackerel sample was suspended in 20 mL of MilliQ water and homogenised with an Ultraturrax homogeniser (Labortechnik, Staufen, Germany) at 13,500 rpm for 30 s 5 mL of 25% trichloroacetic acid solution was added and the solution which was kept at 4 °C for 15 min. After centrifugation at 3500 rpm for 30 min at 4 °C using a Hettich Rotanta 460R centrifuge (Fisher Scientific Ireland, Dublin, Ireland), the solution was filtered using a Whatman number 51 filter paper (Sparks Lab Supplies Ltd., Dublin, Ireland). A 1.5 mL aliquot of the filtrate was collected and mixed with 1.5 mL 0.6% of 2-thiobarbituric acid with the reaction performed in a water bath at 70 °C for 30 min. A standard curve was prepared using a 1,1,3,3-tetraethoxypropane solution. Absorbance of samples and standards were measured spectrophotometrically on a UV-Vis Spectrophotometer (Shimadzu UV-1700, Columbia, USA) at a wavelength of 532 nm. Results were expressed as milligrams of malondialdehyde per kilogram of mackerel (mg MDA/kg mackerel).

## 2.5. Fatty acid profile

### 2.5.1. Microwave-assisted preparation of fatty acid methyl esters (FAMES)

Microwave assisted FAME preparation was carried out using a MARS 6 Express 40 position Microwave Reaction System (CEM Corporation, Matthews, NC, USA) according to Brunton, Mason, and Collins (2015). Briefly, approximately 0.8 g of chilled blended mackerel was accurately weighed out into a reaction Xpress vessel. 100 µL of internal standard (C23:0 methyl ester, achieving a final concentration following extraction of 0.1 mg/mL) as well as 10 mL of potassium hydroxide (2.5%, w/v) in methanol (MeOH) solution were added to the vessel. The reaction vessels containing a 10 mm stir bar each were closed well and introduced into the MARS 6. The saponification programme consisted of bringing the temperature up to 130 °C over 4 min and holding at this temperature for another 4 min. Then, tubes were cooled down in an ice bath. For esterification, 15 mL of 5% (v/v) acetyl chloride in MeOH solution were added to the vessels. Esterification was carried out by heating the vessels to 120 °C over 4 min and holding at this temperature for 2 min. The reaction tubes were removed and cooled on ice for 5 min. To the cooled vessels, 10 mL of pentane was added and vessels were inverted to achieve extraction. After that, about 20 mL of saturated NaCl solution was added, and the inversion process was repeated. The separation of the organic and aqueous layer was achieved, and an aliquot of the upper pentane layer, where the FAMES were located was transferred into GC vials (1.5 mL) containing about 0.2 g anhydrous sodium sulphate and analysed using gas chromatography.

### 2.5.2. Gas chromatography-flame ionisation detector analysis

Separation was carried out using a Clarus 580 Gas Chromatograph (PerkinElmer, Massachusetts, USA) fitted with a flame ionisation detector (GC-FID). The separation of 37 FAMES was carried out in 35 min employing a CP-Sil 88 capillary column (Agilent, Santa Clara, California, USA) with a length of 100 m × 0.25 mm ID and 0.2 µm film. The inlet temperature was set to 250 °C and the injection volume was 0.5 µL. The carrier gas was hydrogen at a constant flow of 1.25 mL/min, and the split ratio was set at 10:1. The oven was set to 80 °C with an initial temperature ramp of 6.2 °C/min to 220 °C which was held for 3.2 min. A second temperature ramp of 6.3 °C/min to 240 °C followed which was held for 6.5 min. The FID was set at 270 °C. Compounds were identified by comparing their retention times with those of authentic standard of FAMES from the Supelco 37 FAME mix. The content of each fatty acid was calculated using following equation (Eq. (1)).

$$FA \text{ content} = \frac{\text{Peak Area}(FAME)}{\text{Peak Area}(ISTD)} \times \frac{\text{Weight}(ISTD)}{\text{Weight}(Sample)} \times 10 \times 0.96 \quad (1)$$

Where, FA content is the amount of a given fatty acid in the sample (mg/g), 10 is the dilution factor and 0.96 is the conversion factor for the internal standard to account for the internal standard already being presented as a FAME.

Due to the variability of the fat content, fatty acid content was normalised and results were expressed in percentages:

$$FA \text{ content} (\%) = \frac{FA \text{ content} (mg/g)}{\text{Total fat content} (mg/g)} \times 100 \quad (2)$$

A full list of all calculated fatty acids in g/kg fish is provided as supplementary material. (Sup 2).

### 2.5.3. Nutritional quality indices

Nutritional quality indices of plasma treated mackerel were assessed from fatty acids data. Saturation index (SI) is a carcinogenesis marker while atherogenicity (AI) and thrombogenicity (TI) are related to the incidence of coronary heart disease. They were calculated using the following equations (Ulbricht & Southgate, 1991):

$$SI = \frac{[C14:0 + C16:0 + C18:0]}{[\sum MUFA + \sum PUFA]} \quad (3)$$

$$AI = \frac{[C12:0 + 4 \times (C14:0) + C16:0]}{[\sum MUFA + \sum PUFA]} \quad (4)$$

$$TI = \frac{[C14:0 + C16:0 + C18:0]}{\left[ 0.5 \times \left( \sum MUFA + \sum n6 \right) + 3 \times \sum n3 + \frac{\sum n3}{\sum n6} \right]} \quad (5)$$

## 2.6. Protein oxidation

Protein carbonyls are biomarkers of protein oxidation. They are generated through several different mechanisms induced during oxidative stress. The content of carbonyl groups was determined by the dinitrophenylhydrazine (DNPH) method using a Biocell protein carbonyl enzyme Immuno-assay kit (BioCell Corporation Ltd, Auckland, New Zealand) (Buss, Chan, Sluis, Domigan, & Winterbourn, 1997). All required reagents were provided by the test kit. Protein extraction was performed according to the procedure by Devries et al. (2008). Briefly, 1 g of mackerel sample was added to 25 mL protein extraction buffer (70 mmol/L sucrose, 220 mmol/L mannitol, 5 mmol/L Hepes, pH 7.2, 1 mmol/L EGTA, pH 7.2, and 150 mmol/L KCl) and homogenised for 30 s using an Ultraturax homogeniser (Labortechnik, Staufen, Germany) set to 13,500 rpm. Each homogenate was adjusted to pH 7.2 using either 1 mol/L HCl or 1 mol/L NaOH and then centrifuged for 10 min at 3500 rpm at 4 °C using a Hettich Rotina 460R centrifuge (Fisher Scientific Ireland, Dublin, Ireland). The amount of soluble protein in the supernatant was quantified by measuring its absorbance at 280 nm using an ND-100 Nanodrop Spectrophotometer (Nanodrop Technologies, Wilmington, DE, USA). Protein carbonyl content determination was then carried out according to the Biocell protein carbonyl enzyme Immuno-assay kit manufacturer's guidelines and measured using a plate reader (FLUOstar Omega Microplate Reader, BMG Labtech GmbH, Offenburg, Germany). Protein carbonyl content was calculated using a standard curve and results expressed as nmol DNPH/mg protein.

## 2.7. Statistical analysis

Analysis of variance (ANOVA) of the dependent variables was carried out using Minitab 17.1.0 (Minitab Inc, Stage College, Pennsylvania, USA). Statistics were performed using a general linear model (GLM) where the treatment and temperature were considered as fixed and days as random factors. Tukey's multiple comparison was used for post-hoc analysis on the significant factors.


## 3. Results and discussion

### 3.1. Lipid oxidation

One of the most common undesirable reactions found in food is lipid oxidation, which can affect both product taste and odour. Lipid oxidation processes occur via a free radical chain mechanism. Consequently, all radical species forming the plasma bulk could encourage these reactions (Gavahian, Chu, Khaneghah, Barba, & Misra, 2018). One of the most common products of lipid oxidation is malondialdehyde, which can be measured by TBARs analysis.


Plasma treatment was found to have no impact on lipid oxidation, as no significant differences ( $p > 0.05$ ) were found for the TBARs values between untreated and treated samples (Fig. 1). However, the TBARs values significantly increased ( $p < 0.05$ ) during storage for both the control and plasma samples (Fig. 1), which is in line with previous research. For example, Romotowska, Karlsdottir, Gudjonsdottir, Kristinsson, and Arason (2016) investigated lipid deterioration of

A)




**Fig. 1.** TBARs values (mg MDA/kg sample) for control (—■—) and plasma (---▲---) A) at  $-20^{\circ}\text{C}$ , B) at  $4^{\circ}\text{C}$  and C) at  $8^{\circ}\text{C}$ . Different letters (lowercase for control and uppercase for plasma) indicate significant differences on the TBARs values during days of storage at the same temperature ( $p < 0.05$ ). No significant differences between control and plasma treated samples at the same storage day were found ( $p > 0.05$ ). All the differences are measured separately for  $4^{\circ}\text{C}$ ,  $8^{\circ}\text{C}$  and  $-20^{\circ}\text{C}$ .

B)



C)



mackerel during frozen storage at  $-18^{\circ}\text{C}$  and  $-25^{\circ}\text{C}$ , with TBARs formation observed over time. These results agree with the findings of [Albertos et al. \(2017\)](#) for mackerel treated with a dielectric barrier discharge at 70 and 80 kV for treatment times of up to 5 min. These authors observed no significant effect on TBARs values between the treated and untreated samples for any of the applied voltages or

treatment times. No significant differences in TBARs values were also reported in Atlantic herring treated at 70 kV for 5 min compared to the control samples 1 day after treatment but a significant increase of those values was observed during the storage period ([Albertos et al., 2019](#)). When [Lee et al. \(2016\)](#) employed a plasma treatment for chicken breasts using a flexible thin-layer dielectric barrier discharge (FTDBD)

plasma, no significant differences in the TBARs values were detected. Similar results were found after plasma treatment using a dielectric barrier discharge on milk samples during a 5 and 10 min treatment process at 250 W (Kim, Park, Kim, Choe, & Jo, 2015). Contrary to these findings, several articles report that plasma induces lipid oxidation. For example, the TBARs values for plasma treated sushi products (Nigiri and Hosomaki) were found to increase after being treated at 70 kV or 80 kV for 5 min (Kulawik et al., 2018). A higher surface area and a higher concentration of unsaturated fatty acids resulted in higher levels of TBARs in Hosomaki sushi (Dawczynski, Schubert, & Jahreis, 2007; Kulawik et al., 2018). A significant increase in TBARs values has been reported by Albertos et al. (2019) in Atlantic herring treated at 80 kV for 5 min. When dried Alaska pollock shreds, semi dried squid and Gwamegi (semi-dried raw Pacific saury) were treated by corona discharge plasma jet (CDPJ), a significant increase ( $P < 0.05$ ) in TBARs values were detected (Choi, Puligundla, & Mok, 2016, 2017; Puligundla, Choi, & Mok, 2018). According to these authors, that increase during the CDPJ treatment may be attributed to the further oxidation of primary lipid oxidation products by reactive species generated by the jet plasma. Similar results were obtained by Olatunde, Benjakul, and Vongkamjan (2019b) who found an increase in TBARs values in plasma treated Asian sea bass slices with increasing treatment times that could be as a result of the ozone generated. In their study, the samples were packed in heat sealed bags filled with argon/oxygen (90:10), treated with high voltage cold atmospheric plasma (HVCAP) at 80 kV RMS for 2.5, 5.0, 7.5 and 10 min and stored at 4 °C for 1 h as post-treatment time. TBARs values were also significantly higher for plasma treated brown and white rice for all treatment conditions with significant increases found for with increasing treatment times of up to 20 min (Lee et al., 2018). It has been reported that the nature of the food matrix and the way the product is handled prior to plasma exposure can affect the oxidation rate in plasma treated samples (Gavahian et al., 2018). The optimization of the process parameters is also needed in order to reduce the rate of lipid oxidation in plasma treated products.

The limit above which TBARs levels would indicate that secondary oxidation products could be detected by consumers have been reported to be 8 mg MDA/kg of fish tissue (Schormüller, 1968) while Connell (1990) suggested TBARs values of 1–2 mg MDA/kg of fish flesh as the limit beyond which fish will normally develop an objectionable odour. The results obtained in the present work were much higher than these proposed limits. However, it is important to highlight that the aim of the present study was to compare non-treated and plasma treated samples rather than a quantification of MDA during storage, with no significant differences being found. Such values could be explained by the overheating that might be caused by a blender used to homogenize the samples and the penetration of oxygen and other species that could have happen while mincing. However, the results obtained in the present work are in line with those reported in mackerel by Romotowska et al. (2016). According to these authors, an increase of oxidation products may be related to the total lipid content seasonal variation, due to differences in accessibility of the feed source and also due to the geographical location of the catch seemed to exert an influence on the lipid stability. It has also been suggested that higher values of lipid oxidation products detected in mackerel could be generated during storage and transportation of the fillets (Crotova, Mozuraityte, Standal, & Rustad, 2019).

### 3.2. Fatty acids profile

The fatty acid profiles for both the control and plasma treated samples during storage were analysed. The study focused on two fatty acids present in the fish which offer high nutritional values namely eicosapentaenoic acid (EPA, C20:5, n-3) and docosahexaenoic acid (DHA, C22:6, n-3). The effects of cold atmospheric plasma on the fatty acid profiles for the mackerel fillets are summarised in Table 1. In

addition, the complete results of the FAME composition in g/kg, the chromatogram of control and plasma treated samples on day 0 at 4 °C individually, and an overlay of these two chromatograms is provided as supplementary material (Sup 2 and Sup 1).

No significant differences ( $p > 0.05$ ) were found for the fatty acid profiles between the control and treated samples for the same temperature and storage day. These results agree with those reported by Kulawik et al. (2018), who described no significant differences in the content of each fatty acid studied for sushi products. In addition, there was no significant change in the fatty acid composition of beef jerky treated with a radio-frequency atmospheric pressure plasma discharge (Kim, Lee, Choi, & Kim, 2014). However, in another study on mackerel, significant differences between the control and treated samples were observed for the content of palmitic acid (C16:0), oleic acid (C18:1, n-9), eicosapentaenoic acid (C20:5, n-3) and docosahexaenoic acid (C22:6 n-3) after treatment with a DBD operating at a discharge voltage of 70 kV and 80 kV for 1, 3 and 5 min (Albertos et al., 2017). This disagreement with our study could be due to different sample preparation methods employed. In the present work, a whole fillet with the skin attached and facing down was treated. This skin likely provided extra protection against the action of the plasma as only the upper side of the mackerel muscle was exposed and hence allowed to interact with the plasma atmosphere, possibly lessening any effect of the plasma. In addition, in our experiment, the fillet was blended after the treatment and stored. This homogenisation could mask possible localised negative effects induced by plasma.

The effect of cold atmospheric plasma on the fatty acid profile has also been studied on different matrices including soy bean oil (Yepez & Keener, 2016). A significant change of the fatty acid composition was reported after applying a plasma discharge at 90 kV for extended treatment times of 1, 1.5, 2, 4, 6 and 12 h. They reported that  $\alpha$ -linolenic (18:3, n-6) and linoleic (18:2, n-6) acid significantly decreased as a function of treatment time, while oleic (C18:1, n-9), stearic (C18:0), and palmitic (C16:0) acid significantly increased. Kim et al. (2015) found a significant change of butyric (C4:0) and caprylic acid (C8:0) content in milk for plasma treatments of 5 and 10 min with an input power of 250 W. Surowsky, Schluter, and Knorr (2015) reported that the effect of plasma was more evident in liquid samples than solid samples due to their ability to interact with the bulk of the liquid matrix; this could explain why a significant effect of plasma treatment for oil and milk was previously reported. On the other hand, the effect of the temperature on the oxidative stability of fatty acid methyl esters has also been studied. Giua, Blasi, Simonetti, and Cossignani (2013) found that the methyl forms of conjugated linoleic acid (Me-CLA) showed the highest isomer profile modifications during a heating treatment at 180 °C for 15, 30, 45 and 60 min in order to simulate a frying process. In fact, the percentage of cis-9,trans-11 and trans-10,cis-12 CLA isomers decreased during heating time and after 60 min, 13.6% of trans, trans isomers and 4.9% of cis, cis isomers were detected. In another study, Cossignani, Giua, Simonetti, and Blasi (2014) found that methyl octanoate was the main methyl ester compound detected both for Me-CLA and for Me-LA oxidized samples after being treated at 180 °C for 15, 30, 45 and 60 min. Other studies focused on the effect of cold plasma on the fatty acid profile of chocolate milk drink and guava-flavored whey beverage when compared to conventional pasteurization has been published (Coutinho et al., 2019; Silveira et al., 2019). The chocolate milk drink subjected to intermediate cold plasma treatment conditions presented an improved fatty acid profile when compared to the pasteurized product, with a significant reduction in stearic acid and an increase in myristoleic acid, linoleic, and PUFA levels (Coutinho et al., 2019). However, when mild and severe conditions were applied, there was an increase in SFA and a decrease in both MUFA and PUFA. The changes in fatty acids were attributed to the oxygen radicals produced during plasma treatment, including ozone, that react with the unsaturated fatty acids and break down the double bonds, leading to an increase in SFA (Gavahian et al., 2018). No significant differences were

**Table 1**

Fatty acid profile during storage study. No significant differences ( $p > 0.05$ ) were found for any lipid class. The differences are measured separately for 4 °C, 8 °C and –20 °C.

Temperature (°C)	Treatment	Time (days)	SFA (%)	MUFA (%)	PUFA (%)	EPA (%)	DHA (%)		
–20	Control	0	34.3 ± 0.2	30.7 ± 2.7	35.0 ± 2.8	9.9 ± 1.5	16.5 ± 1.3		
		2	37.5 ± 5.9	30.2 ± 3.4	32.3 ± 2.5	8.4 ± 0.7	14.6 ± 1.3		
		5	33.8 ± 1.3	27.8 ± 1.4	38.4 ± 1.3	11.8 ± 0.3	18.6 ± 1.2		
		7	36.7 ± 0.5	30.1 ± 0.2	33.2 ± 0.5	8.1 ± 0.2	16.2 ± 0.3		
		14	35.0 ± 0.3	29.3 ± 4.1	35.6 ± 3.8	9.8 ± 1.1	16.7 ± 2.1		
	Plasma	0	35.1 ± 0.3	27.8 ± 1.4	37.1 ± 1.5	10.2 ± 0.6	18.0 ± 0.7		
		2	36.4 ± 2.4	14.3 ± 8.3	39.3 ± 6.6	11.5 ± 0.7	20.9 ± 5.8		
		5	34.7 ± 1.4	28.9 ± 0.5	36.4 ± 0.9	10.3 ± 0.6	17.7 ± 0.6		
		7	34.2 ± 0.6	27.3 ± 1.1	38.6 ± 0.5	11.7 ± 1.0	17.6 ± 0.6		
		14	34.1 ± 0.9	31.3 ± 0.9	34.7 ± 1.4	9.5 ± 0.2	17.2 ± 1.0		
		4	Control	0	34.3 ± 0.2	30.7 ± 2.7	35.0 ± 2.8	9.9 ± 1.5	16.5 ± 1.3
				2	33.7 ± 0.4	29.4 ± 0.3	36.9 ± 0.5	10.1 ± 0.4	18.0 ± 0.8
				5	34.3 ± 2.0	26.2 ± 4.4	39.5 ± 2.9	11.5 ± 0.9	20.8 ± 3.3
				7	35.1 ± 1.4	31.5 ± 1.4	33.4 ± 1.8	9.0 ± 0.3	15.7 ± 0.4
7	35.1 ± 0.3			27.8 ± 1.4	37.1 ± 1.5	10.2 ± 0.6	18.0 ± 0.7		
Plasma	0		35.1 ± 0.3	27.8 ± 1.4	37.1 ± 1.5	10.2 ± 0.6	18.0 ± 0.7		
	2		34.1 ± 0.5	32.8 ± 0.3	33.1 ± 0.3	9.1 ± 0.1	15.8 ± 0.2		
	5		33.4 ± 2.5	31.9 ± 7.1	34.8 ± 4.8	8.6 ± 1.6	16.0 ± 1.9		
	7		36.9 ± 2.0	25.2 ± 1.8	37.9 ± 1.6	11.9 ± 1.3	18.2 ± 1.0		
	8		Control	0	34.3 ± 0.2	30.7 ± 2.7	35.0 ± 2.8	9.9 ± 1.5	16.5 ± 1.3
				2	36.2 ± 6.4	30.0 ± 5.4	33.8 ± 11.8	9.3 ± 3.5	17.7 ± 7.3
				5	33.7 ± 0.6	30.9 ± 2.5	35.4 ± 3.0	8.9 ± 0.7	17.8 ± 2.5
7				36.1 ± 0.2	30.0 ± 4.1	34.0 ± 4.3	9.0 ± 1.6	15.6 ± 2.3	
7				36.1 ± 0.2	30.0 ± 4.1	34.0 ± 4.3	9.0 ± 1.6	15.6 ± 2.3	
Plasma		0	35.1 ± 0.3	27.8 ± 1.4	37.1 ± 1.5	10.2 ± 0.6	18.0 ± 0.7		
		2	34.7 ± 1.1	30.3 ± 2.3	35.0 ± 3.2	9.63 ± 2.4	15.4 ± 0.9		
		5	33.1 ± 2.6	34.1 ± 2.1	32.8 ± 0.6	9.5 ± 0.7	15.9 ± 0.8		
		7	33.9 ± 0.2	29.7 ± 0.5	36.4 ± 0.3	10.3 ± 0.6	18.3 ± 0.9		

found by [Silveira et al. \(2019\)](#) between the plasma-treated guava-flavored whey beverage and the pasteurized product for butanoic, octanoic, decanoic, dodecanoic, myristic, and stearic acids levels. However, an increase in palmitic acid and/or a decrease in hexanoic acid, oleic acid, and MUFA were observed probably due to the reactivity of the plasma reactive species against the unsaturated fatty acids.

### 3.3. Nutritional indices

Nutritional indices were calculated from the fatty acid profiles of both treated and untreated samples using the equations outlined in section 2.5.3. The results are summarised in [Table 2](#). No significant differences ( $p > 0.05$ ) were found for any index, which is to be expected given no significant differences were found for fatty acid profiles between treatments. To the best of the authors' knowledge, this is the first time that these indices have been calculated for cold plasma treated mackerel. AI relates the risk of atherosclerosis and is based on fatty acids which can increase (C12:0, C14:0 and C16:0) or decrease ( $\Sigma$ MUFA,  $\Sigma$ PUFA) the level of blood cholesterol. TI values relate to the tendency to form clots in the blood vessels, defined as the relationship between the pro-thrombogenic (saturated) and the anti-thrombogenic fatty acids (MUFAs, n-6 PUFAs and n-3 PUFAs) ([Ulbricht & Southgate, 1991](#)). The smaller the AI and TI values, the greater health benefits derived from the product ([Ulbricht & Southgate, 1991](#)). In the present work, no significant differences ( $p > 0.05$ ) were found in the AI values in the control and plasma treated samples along the storage at the temperatures studied. Both treated and untreated samples had an AI value of 0.7 on day 0. The AI values were 0.6 and 0.7 in plasma treated samples along the 14 days at –20 °C, while the control had an AI value which varied from 0.6 to 0.9. At 4 °C, plasma treated samples showed AI values of 0.7 and 0.8 during the 7 days of storage which were in a similar range than the untreated samples. When samples were stored at 8 °C, the AI values in plasma treated and untreated samples varied from

0.6 to 0.8. On the other hand, the TI values obtained on day 0 in plasma treated samples were 0.3 which did not change ( $p > 0.05$ ) along the investigated storage period for any of the temperatures studied. This index showed values between 0.2 and 0.4 in the control samples during the storage. AI values of 0.48 and TI values of 0.24 have been reported for mackerel by [Fernandes et al. \(2014\)](#) who indicated the range of the expected values for AI and TI indices to be up to 1 and 0.5, respectively. According to these authors, AI and TI values detected in the present work would be within the expected range both in plasma treated and untreated mackerel fillets and also along the storage period.

Mackerel would be considered a healthy food because the ratio PUFA/SFA was above 0.45 and the plasma treatment did not decrease the ratio below that recommended value ([Hmsso, 1994](#)). Moreover, it can be observed that this ratio did not decrease during the storage for any of the temperatures studied. SI indicates the relationship between the sum of saturated fatty acids (C14:0, C16:0 and C18:0) (pro-thrombogenic) and unsaturated fatty acids (anti-thrombogenic). It has been reported that myristic acid (C14:0), palmitic acid (C16:0) and stearic acid (C18:0) are associated with an increased risk of coronary heart disease ([Zong et al., 2016](#)). Although to authors' knowledge, there is no numerical values assigned to SI, a food with lower values of these SFA compared to unsaturated fatty acids would be considered a healthier food. In the present work, the SI value in untreated mackerel varied from 0.5 to 0.6 and it was at the same level ( $p > 0.05$ ) in plasma treated samples during the storage period for all the temperatures studied. The n-6/n-3 ratio is considered a good indicator of nutritional value of dietary fat. Diets containing higher amount of n-6 PUFA or high n-6/n-3 ratio have been reported to promote the pathogenesis of cancer, inflammatory and cardiovascular diseases ([Simopoulos, 2002, 2008](#)). According to the nutritional recommendations, the n-6/n-3 ratio should not be higher than 4.0 ([Hmsso, 1994](#)). In our study, the values of this ratio were within this range for plasma treated mackerel, both immediately after the treatment, and along the storage period for all the

**Table 2**

Nutritional indices values during storage study. No significant differences ( $p > 0.05$ ) were found for any nutritional class. The differences are measured separately for 4 °C, 8 °C and -20 °C.

Temperature (°C)	Treatment	Time (days)	Saturation index (SI)	Atherogenic index (AI)	Thrombogenic index (TI)	n-6/n-3	PUFA/SFA	
-20	Control	0	0.5 ± 0.0	0.7 ± 0.0	0.3 ± 0.0	0.1 ± 0.0	1.0 ± 0.1	
		2	0.6 ± 0.2	0.9 ± 0.2	0.4 ± 0.2	0.1 ± 0.0	0.9 ± 0.2	
		5	0.5 ± 0.0	0.6 ± 0.1	0.3 ± 0.0	0.1 ± 0.0	1.1 ± 0.1	
		7	0.5 ± 0.0	0.9 ± 0.0	0.2 ± 0.1	0.1 ± 0.0	0.9 ± 0.0	
		14	0.5 ± 0.0	0.8 ± 0.0	0.3 ± 0.0	0.1 ± 0.0	1.0 ± 0.1	
	Plasma	0	0.5 ± 0.0	0.7 ± 0.0	0.3 ± 0.0	0.1 ± 0.0	1.0 ± 0.0	
		2	0.5 ± 0.1	0.7 ± 0.1	0.3 ± 0.0	0.1 ± 0.0	1.0 ± 0.1	
		5	0.5 ± 0.0	0.7 ± 0.1	0.3 ± 0.0	0.1 ± 0.0	1.0 ± 0.1	
		7	0.5 ± 0.0	0.7 ± 0.0	0.3 ± 0.0	0.1 ± 0.0	1.1 ± 0.0	
		14	0.5 ± 0.0	0.6 ± 0.1	0.3 ± 0.0	0.1 ± 0.0	1.0 ± 0.1	
	Temperature (°C)	Treatment	Time (days)	Saturation index	Atherogenic index	Thrombogenic index	n-6/n-3	PUFA/SFA
	4	Control	0	0.5 ± 0.0	0.7 ± 0.0	0.3 ± 0.0	0.1 ± 0.0	1.0 ± 0.1
			2	0.5 ± 0.0	0.6 ± 0.0	0.3 ± 0.0	0.1 ± 0.0	1.0 ± 0.0
			5	0.5 ± 0.1	0.6 ± 0.1	0.2 ± 0.1	0.1 ± 0.0	1.1 ± 0.1
7			0.5 ± 0.0	0.8 ± 0.1	0.3 ± 0.0	0.1 ± 0.0	0.9 ± 0.1	
7			0.5 ± 0.0	0.8 ± 0.1	0.3 ± 0.0	0.1 ± 0.0	1.0 ± 0.1	
Plasma		0	0.5 ± 0.0	0.7 ± 0.0	0.3 ± 0.0	0.1 ± 0.0	1.0 ± 0.0	
		2	0.5 ± 0.0	0.7 ± 0.0	0.3 ± 0.0	0.1 ± 0.0	0.9 ± 0.0	
		5	0.5 ± 0.1	0.8 ± 0.1	0.3 ± 0.0	0.1 ± 0.0	1.04 ± 0.0	
		5	0.5 ± 0.1	0.8 ± 0.1	0.3 ± 0.0	0.1 ± 0.0	1.04 ± 0.0	
		7	0.5 ± 0.0	0.8 ± 0.1	0.3 ± 0.0	0.1 ± 0.0	1.0 ± 0.0	
Temperature (°C)		Treatment	Time (days)	Saturation index	Atherogenic index	Thrombogenic index	n-6/n-3	PUFA/SFA
8		Control	0	0.5 ± 0.0	0.7 ± 0.0	0.3 ± 0.0	0.1 ± 0.0	1.0 ± 0.1
			2	0.6 ± 0.2	0.7 ± 0.2	0.3 ± 0.0	0.1 ± 0.1	0.9 ± 0.4
			5	0.5 ± 0.0	0.6 ± 0.1	0.3 ± 0.0	0.1 ± 0.0	1.0 ± 0.1
	7		0.5 ± 0.0	0.8 ± 0.0	0.3 ± 0.0	0.1 ± 0.0	0.9 ± 0.1	
	7		0.5 ± 0.0	0.8 ± 0.0	0.3 ± 0.0	0.1 ± 0.0	1.0 ± 0.0	
	Plasma	0	0.5 ± 0.0	0.7 ± 0.0	0.3 ± 0.0	0.1 ± 0.0	1.0 ± 0.0	
		2	0.5 ± 0.1	0.8 ± 0.2	0.3 ± 0.0	0.1 ± 0.0	1.0 ± 0.1	
		5	0.5 ± 0.1	0.6 ± 0.1	0.3 ± 0.0	0.1 ± 0.0	1.0 ± 0.1	
		5	0.5 ± 0.1	0.6 ± 0.1	0.3 ± 0.0	0.1 ± 0.0	1.0 ± 0.1	
		7	0.5 ± 0.0	0.6 ± 0.0	0.3 ± 0.0	0.1 ± 0.0	1.0 ± 0.0	

temperatures studied. Two studies focused on the effect of cold plasma on nutritional indices of chocolate milk drink and guava-flavored whey beverage compared to the pasteurized product has been published (Coutinho et al., 2019; Silveira et al., 2019). According to Coutinho et al. (2019), the chocolate milk drink submitted to mild plasma conditions had similar AI and TI indices ( $p > 0.05$ ) when compared to the pasteurized product. However, when mild and severe plasma treatment conditions were applied, higher AI and TI indices were obtained, highlighting the importance of optimizing the process conditions to minimize negative impacts. In another study, a plasma treated guava-flavored whey beverage showed higher AI and TI values than the pasteurized product (Silveira et al., 2019).

### 3.4. Protein oxidation

Like lipid oxidation, protein oxidation is also initiated by free radicals and can be measured by the loss of sulfhydryl groups (S-H) or by the formation of protein carbonyls which are both formed during this oxidation process. Radical species can lead to the scission of the peptide backbone, the conversion of one amino acid to a different one, the formation of crosslinking as well as oxidative changes of the amino acid side chains (Estévez, 2011). For instance, an increase in the carbonyl content could lead to the formation of crosslinking, resulting in a decrease of the juiciness, tenderness and quality traits of meat products (Estévez, 2011). Hence, it is important to study if plasma treatments encourage this undesirable reaction.

The carbonyl content in mackerel ranged from 0.5 to 1.5 nmol/mg protein for the control samples, to 0.5–2.5 nmol/mg of protein for the plasma treated samples. The results indicated that the plasma treatment significantly accelerated the formation of carbonyl groups ( $p < 0.05$ ) for all storage temperatures studied (Fig. 2); i.e. -20 °C (Fig. 2A), 4 °C (Fig. 2B) and 8 °C (Fig. 2C). In addition, the carbonyl content increased significantly for treated and untreated samples during storage


independent of the temperature. Moreover, in the present study the storage temperature had a significant impact ( $p < 0.05$ ) on carbonyl formation, resulting in greater formation at higher temperatures.

A formation of protein carbonyls in mackerel mince during a storage study at 5 °C has been also reported by Eymard, Baron, and Jacobsen (2009) and at -18 °C by Ozen and Soyer (2018). Eymard et al. (2009) found values much higher than the ones presented in this manuscript (from 2 nmol of carbonyl per mg of protein up to 10 nmol/mg after 92 h). This increase reported by Eymard et al. (2009) could be due to the mincing of the mackerel (before the treatment), making it possibly easier for oxygen to penetrate the matrix, possibly resulting in accelerated oxidation. However, in another study, Babakhani, Farvin, and Jacobsen (2016) reported that the initial content of carbonyl groups in mackerel mince was 1 nmol/mg protein and it increased up to 1.4 nmol/mg protein after 192 h of storage at 5 °C. The reason of this low values can be explained because the mackerel mince was packaged under vacuum. This may have slowed down the oxidation due to the lack of potential gases such as oxygen which can facilitate the oxidation of the food components.


The effect of high voltage cold atmospheric plasma (HVCAP) generated with the mixture of argon and oxygen (90:10) for 5 min at 28 °C with the addition of two different antioxidants such as ascorbic acid and ethanolic coconut husk extract on the quality of sea bass, has been investigated by Olatunde, Benjakul, and Vongkamjan (2019a). In this study, heat-sealed bags containing sea bass slices with and without antioxidants, filled with 90% argon and 10% oxygen using a vacuum-packaging machine, were treated at 16 kV RMS for 5 min using a dielectric barrier discharge system. A significantly higher ( $p < 0.05$ ) carbonyl content of HVCAP treated samples compared to the control ones was reported, regardless the addition of the antioxidants, concluding that HVCAP could lead to the oxidation of proteins.

There is a lack of data in the literature to compare with and to explain in more detail the effects of cold atmospheric plasma on protein

### Protein oxidation -20° C



### Protein oxidation 4° C



### Protein oxidation 8° C




Fig. 2. Carbonyl content values (nmol carbonyl/mg protein) for control (—■—) and plasma (---▲---) A) at -20 °C, B) at 4 °C and C) at 8 °C. Different letters (lowercase for control and uppercase for control) indicate significant differences on the carbonyl content value during days of storage at the same temperature ( $p < 0.05$ ). Stars show significant differences between treatments at the same storage day being \* ( $p < 0.05$ ) and \*\* ( $p < 0.005$ ). All the differences are measured separately for 4 °C, 8 °C and -20 °C.

oxidation in food matrices. To date, most studies have examined the effects of plasma on protein solutions. A whey protein isolate solution (2% (w/v) in 50 mmol/L phosphate buffer, pH 6.8) was treated by cold atmospheric plasma at 70 kV for periods of 1, 5, 10, 15, 30 and 60 min and reported a significant increase in the carbonyl content in plasma treated samples compared to the untreated ones (Segat, Misra, Cullen, & Innocente, 2015). They suggested that the formation of carbonyls could be attributed to the modifications of a number of amino acid side chain groups, especially with NH- or NH<sub>2</sub> or by peptide bond cleavages. On the other hand, K. H. LeeWoo, et al. (2018) and Lee, et al. (2018) found a decrease in the total free sulfhydryl groups (-SH) for peanut protein isolate solutions treated by cold plasma which is related to the oxidation of these groups to form disulphide bonds. Moreover, a decreased in the content of sulfhydryl groups was also reported all the

treatment times (1, 5, 10, 15, 30, and 60 min) after applying a voltage of 70 kV on whey protein isolate solution (Segat et al., 2015). A significant processing time dependant increase ( $p < 0.05$ ) in carbonyl content has been detected in the crude protease extract from squid mantle treated by a dielectric barrier discharge system at 60 kV for different treatment times (Nyaisaba et al., 2019). These authors also found a decrease in the total sulfhydryl group ( $p < 0.05$ ) (up to about 40%) as exposure time extended which was attributed to the formation of disulfide through cross-linking of sulfhydryl groups influenced by reactive species generated by cold plasma (Segat et al., 2015). Similar results were reported by Sharifian, Soltanizadeh, and Abbaszadeh (2019) who found an increase ( $p < 0.05$ ) in the carbonyl content of beef myofibrillar proteins after the DBD plasma treatment which was higher at increased processing times. The carbonyl formation has been

attributed to the modification of certain amino acid side chains with  $-NH_2$  or  $-NH$ , or to the cleavage of peptide bonds (Segat et al., 2015). Sharifian et al. (2019) reported a significant increase ( $p < 0.05$ ) in free sulfhydryl groups in beef myofibrillar proteins after 10 min of atmospheric cold plasma (ACP) treatment compared to the untreated samples and those treated for 5 min. According to these authors, the alteration of the tertiary structure of the myofibrillar proteins caused by 10 min plasma treatment, could have facilitated the hidden sulfhydryl groups to become exposed at the protein surface and consequently, be more vulnerable to the treatment. ACP has been reported to induce the loss of  $-SH$  groups from amino acid cysteine present in the protein structure (Segat et al., 2015).

The data obtained showed that cold atmospheric plasma did not encourage observable undesirable reactions such as lipid oxidation to the bulk of the treated samples. In addition, the stability of the fatty acid composition of mackerel was not affected by the treatment, along with their nutritional quality indices. However, cold atmospheric plasma could accelerate the formation of carbonyls which are related to protein oxidation. Further research is required on the impact of cold plasma technology on the quality of seafood products in order to be implemented by the fish industry.

### Declaration of competing interest

The authors declare no conflict of interest.

### Acknowledgments

The authors would like to thank Dr. Gonzalo Delgado Pando for his statistical assistance. The work on this paper is supported by project that has received funding from the European Union's Horizon 2020 Spreading Excellence and Widening Participation programme under Grant Agreement No. 692276. Authors also wish to acknowledge the financial support received by the JPI project ProHealth (Ref:15/HDHL/1 PROHEALTH) "Innovative processing to preserve positive health effects in pelagic fish products".

### Appendix A. Supplementary data

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.lwt.2019.108697>.

### References

- Albertos, I., Martín-Diana, A., Cullen, P., Tiwari, B., Ojha, S., Bourke, P., ... Rico, D. (2017). Effects of dielectric barrier discharge (DBD) generated plasma on microbial reduction and quality parameters of fresh mackerel (*Scomber scombrus*) fillets. *Innovative Food Science & Emerging Technologies*, 44, 117–122.
- Albertos, I., Martín-Diana, A. B., Cullen, P. J., Tiwari, B. K., Ojha, K. S., Bourke, P., et al. (2019). Shelf-life extension of herring (*Clupea harengus*) using in-package atmospheric plasma technology. [Article] *Innovative Food Science & Emerging Technologies*, 53, 85–91. <https://doi.org/10.1016/j.ifset.2017.09.010>.
- Amini, M., & Ghoranneviss, M. (2016). Effects of cold plasma treatment on antioxidants activity, phenolic contents and shelf life of fresh and dried walnut (*Juglans regia* L.) cultivars during storage. [Article] *LWT-Food Science and Technology*, 73, 178–184. <https://doi.org/10.1016/j.lwt.2016.06.014>.
- Arvanitoyannis, I. S., Kotsanopoulos, K. V., & Savva, A. G. (2017). Use of ultrasounds in the food industry-methods and effects on quality, safety, and organoleptic characteristics of foods: A review. *Critical Reviews in Food Science and Nutrition*, 57(1), 109–128. <https://doi.org/10.1080/10408398.2013.860514>.
- Auwah, G. B., Ramaswamy, H. S., & Economides, A. (2007). Thermal processing and quality: Principles and overview. [Review] *Chemical Engineering and Processing*, 46(6), 584–602. <https://doi.org/10.1016/j.ccep.2006.08.004>.
- Babakhani, A., Farvin, K. H. S., & Jacobsen, C. (2016). Antioxidative effect of seaweed extracts in chilled storage of minced Atlantic mackerel (*Scomber scombrus*): Effect on lipid and protein oxidation. [Article] *Food and Bioprocess Technology*, 9(2), 352–364. <https://doi.org/10.1007/s11947-015-1630-9>.
- Bourke, P., Ziuzina, D., Boehm, D., Cullen, P. J., & Keener, K. (2018). The potential of cold plasma for safe and sustainable food production. [Review] *Trends in Biotechnology*, 36(6), 615–626. <https://doi.org/10.1016/j.tibtech.2017.11.001>.
- Brunton, N. P., Mason, C., & Collins, M. J. (2015). Rapid Microwave assisted preparation of fatty acid methyl esters for the analysis of fatty acid profiles in foods. [Article] *Journal of Analytical Chemistry*, 70(10), 1218–1224. <https://doi.org/10.1134/s1061934815100044>.
- Buss, H., Chan, T. P., Sluis, K. B., Domigan, N. M., & Winterbourn, C. C. (1997). Protein carbonyl measurement by a sensitive ELISA method. *Free Radical Biology and Medicine*, 23(3), 361–366. [https://doi.org/10.1016/S0891-5849\(97\)00104-4](https://doi.org/10.1016/S0891-5849(97)00104-4).
- Choi, S., Puligundla, P., & Mok, C. (2016). Microbial decontamination of dried Alaska pollock shreds using corona discharge plasma jet: Effects on physicochemical and sensory characteristics. *Journal of Food Science*, 81(4), M952–M957.
- Choi, S., Puligundla, P., & Mok, C. (2017). Effect of corona discharge plasma on microbial decontamination of dried squid shreds including physico-chemical and sensory evaluation. *LWT*, 75, 323–328.
- Connell, J. (1990). Methods of assessing and selecting for quality. *Control of Fish Quality*, 2.
- Cossignani, L., Giua, L., Simonetti, M. S., & Blasi, F. (2014). Volatile compounds as indicators of conjugated and unconjugated linoleic acid thermal oxidation. *European Journal of Lipid Science and Technology*, 116(4), 407–412.
- Coutinho, N. M., Silveira, M. R., Fernandes, L. M., Moraes, J., Pimentel, T. C., Freitas, M. Q., ... Cruz, A. G. (2019). Processing chocolate milk drink by low-pressure cold plasma technology. [Article] *Food Chemistry*, 278, 276–283. <https://doi.org/10.1016/j.foodchem.2018.11.061>.
- Cropotova, J., Mozuraityte, R., Standal, I. B., & Rustad, T. (2019). The influence of cooking parameters and chilled storage time on quality of Sous-Vide Atlantic mackerel (*Scomber scombrus*). *Journal of Aquatic Food Product Technology*, 28(5), 505–518.
- Cullen, P. J., Lalor, J., Scally, L., Boehm, D., Milosavljevic, V., Bourke, P., et al. (2018). Translation of plasma technology from the lab to the food industry. [Review] *Plasma Processes and Polymers*, 15(2), 11. <https://doi.org/10.1002/ppap.201700085>.
- Daryaei, H., Yousef, A. E., & Balasubramanian, V. (2016). *Microbiological aspects of high-pressure processing of food: Inactivation of microbial vegetative cells and spores high pressure processing of food*. Springer.
- Dawczynski, C., Schubert, R., & Jahreis, G. (2007). Amino acids, fatty acids, and dietary fibre in edible seaweed products. [Article] *Food Chemistry*, 103(3), 891–899. <https://doi.org/10.1016/j.foodchem.2006.09.041>.
- De Alba, M., Pérez-Andrés, J. M., Harrison, S. M., Brunton, N. P., Burgess, C. M., & Tiwari, B. K. (2019). High pressure processing on microbial inactivation, quality parameters and nutritional quality indices of mackerel fillets. *Innovative Food Science & Emerging Technologies*, 55, 80–87.
- Devi, Y., Thirumdas, R., Sarangapani, C., Deshmukh, R. R., & Annapure, U. S. (2017). Influence of cold plasma on fungal growth and aflatoxins production on groundnuts. *Food Control*, 77, 187–191. <https://doi.org/10.1016/j.foodcont.2017.02.019>.
- Devries, M. C., Hamadeh, M. J., Glover, A. W., Raha, S., Samjoo, I. A., & Tarnopolsky, M. A. (2008). Endurance training without weight loss lowers systemic, but not muscle, oxidative stress with no effect on inflammation in lean and obese women. *Free Radical Biology and Medicine*, 45(4), 503–511. <https://doi.org/10.1016/j.freeradbiomed.2008.04.039>.
- Domingo, J. L. (2016). Nutrients and chemical pollutants in fish and shellfish. Balancing health benefits and risks of regular fish consumption. [Review] *Critical Reviews in Food Science and Nutrition*, 56(6), 979–988. <https://doi.org/10.1080/10408398.2012.742985>.
- Estévez, M. (2011). Protein carbonyls in meat systems: A review. *Meat Science*, 89(3), 259–279.
- Eymard, S., Baron, C. P., & Jacobsen, C. (2009). Oxidation of lipid and protein in horse mackerel (*Trachurus trachurus*) mince and washed minces during processing and storage. [Article] *Food Chemistry*, 114(1), 57–65. <https://doi.org/10.1016/j.foodchem.2008.09.030>.
- FAO (2018). *Herring and mackerel quota cut sharply in the North Atlantic*. from <http://www.fao.org/in-action/globefish/market-reports/resource-detail/en/c/1110767/>.
- Fernandes, C. E., da Silva Vasconcelos, M. A., de Almeida Ribeiro, M., Sarubbo, L. A., Andrade, S. A. C., & de Melo Filho, A. B. (2014). Nutritional and lipid profiles in marine fish species from Brazil. *Food Chemistry*, 160, 67–71.
- Gavahian, M., Chu, Y.-H., Khaneghah, A. M., Barba, F. J., & Misra, N. (2018). A critical analysis of the cold plasma induced lipid oxidation in foods. *Trends in Food Science & Technology*, 77, 32–41.
- Ghani, M. A., Barril, C., Bedgood, D. R., & Prenzler, P. D. (2017). Measurement of anti-oxidant activity with the thiobarbituric acid reactive substances assay. [Review] *Food Chemistry*, 230, 195–207. <https://doi.org/10.1016/j.foodchem.2017.02.127>.
- Giua, L., Blasi, F., Simonetti, M., & Cossignani, L. (2013). Oxidative modifications of conjugated and unconjugated linoleic acid during heating. *Food Chemistry*, 140(4), 680–685.
- Gram, L., & Dalgaard, P. (2002). Fish spoilage bacteria – problems and solutions. *Current Opinion in Biotechnology*, 13(3), 262–266. [https://doi.org/10.1016/S0958-1669\(02\)00309-9](https://doi.org/10.1016/S0958-1669(02)00309-9).
- Hmso, U. (1994). *Nutritional aspects of cardiovascular disease (report on health and social subjects No. 46)* London: HMSO.
- Jermann, C., Koutchma, T., Margas, E., Leadley, C., & Ros-Polski, V. (2015). Mapping trends in novel and emerging food processing technologies around the world. [Article] *Innovative Food Science & Emerging Technologies*, 31, 14–27. <https://doi.org/10.1016/j.ifset.2015.06.007>.
- Karim, N. U., Kennedy, T., Linton, M., Watson, S., Gault, N., & Patterson, M. F. (2011). Effect of high pressure processing on the quality of herring (*Clupea harengus*) and haddock (*Melanogrammus aeglefinus*) stored on ice. [Article] *Food Control*, 22(3–4), 476–484. <https://doi.org/10.1016/j.foodcont.2010.09.030>.
- Kim, Jayasena, D., Yong, H., & Jo, C. (2016). *Quality of cold plasma treated foods of animal origin-cold Plasma in Food and agriculture*. Elsevier.
- Kim, J.-S., Lee, E.-J., Choi, E. H., & Kim, Y.-J. (2014). Inactivation of *Staphylococcus aureus* on the beef jerky by radio-frequency atmospheric pressure plasma discharge

- treatment. *Innovative Food Science & Emerging Technologies*, 22, 124–130. <https://doi.org/10.1016/j.ifset.2013.12.012>.
- Kim, Y. H., Park, S., Kim, K., Choe, W., & Jo, C. (2015). Microbial safety and quality attributes of milk following treatment with atmospheric pressure encapsulated dielectric barrier discharge plasma. *Food Control*, 47, 451–456.
- Kröger, E., & Laforce, R. (2016). Fish consumption, brain mercury, and neuropathology in patients with Alzheimer disease and dementia. *Journal of the American Medical Association*, 315(5), 465–466. <https://doi.org/10.1001/jama.2016.0005>.
- Kulawik, P., Alvarez, C., Cullen, P. J., Aznar-Roca, R., Mullen, A. M., & Tiwari, B. (2018). The effect of non-thermal plasma on the lipid oxidation and microbiological quality of sushi. *Innovative Food Science & Emerging Technologies*, 45, 412–417.
- Kumar, A., Mastana, S. S., & Lindley, M. R. (2016). n-3 Fatty acids and asthma. [Review] *Nutrition Research Reviews*, 29(1), 1–16. <https://doi.org/10.1017/s0954422415000116>.
- Lacombe, A., Niemira, B. A., Gurtler, J. B., Sites, J., Boyd, G., Kingsley, D. H., ... Chen, H. (2017). Nonthermal inactivation of norovirus surrogates on blueberries using atmospheric cold plasma. *Food Microbiology*, 63, 1–5. <https://doi.org/10.1016/j.fm.2016.10.030>.
- Lee, Y. H., Kim, H.-J., Choe, W., Yoo, S. J., Jang, E. J., & Jo, C. (2016). Evaluation of the microbiological safety, quality changes, and genotoxicity of chicken breast treated with flexible thin-layer dielectric barrier discharge plasma. *Food science and biotechnology*, 25(4), 1189–1195.
- Lee, O'keefe, J. H., Lavie, C. J., & Harris, W. S. (2009). Omega-3 fatty acids: Cardiovascular benefits, sources and sustainability. *Nature Reviews Cardiology*, 6(12), 753.
- Lee, Woo, K. S., Yong, H. I., Jo, C., Lee, S. K., Lee, B. W., ... Kim, H.-J. (2018). Assessment of microbial safety and quality changes of brown and white cooked rice treated with atmospheric pressure plasma. *Food science and biotechnology*, 27(3), 661–667.
- Lee, K. H., Woo, K. S., Yong, H. I., Jo, C., Lee, S. K., Lee, B. W., ... Kim, H. J. (2018). Assessment of microbial safety and quality changes of brown and white cooked rice treated with atmospheric pressure plasma. [Article] *Food Science and Biotechnology*, 27(3), 661–667. <https://doi.org/10.1007/s10068-017-0297-6>.
- Liao, X. Y., Liu, D. H., Xiang, Q. S., Ahn, J., Chen, S. G., Ye, X. Q., et al. (2017). Inactivation mechanisms of non-thermal plasma on microbes: A review. [Review] *Food Control*, 75, 83–91. <https://doi.org/10.1016/j.foodcont.2016.12.021>.
- Lu, X., Naidis, G. V., Laroussi, M., Reuter, S., Graves, D. B., & Ostrikov, K. (2016). Reactive species in non-equilibrium atmospheric-pressure plasmas: Generation, transport, and biological effects. *Physics Reports*, 630, 1–84. <https://doi.org/10.1016/j.physrep.2016.03.003>.
- Maraschiello, C., Sárraga, C., & Garcia Regueiro, J. (1999). Glutathione peroxidase activity, TBARS, and  $\alpha$ -tocopherol in meat from chickens fed different diets. *Journal of Agricultural and Food Chemistry*, 47(3), 867–872.
- Misra (2016). *Quality of cold plasma treated plant foods Cold Plasma in Food and Agriculture*. Elsevier.
- Misra, Moiseev, T., Patil, S., Pankaj, S., Bourke, P., Mosnier, J., ... Cullen, P. (2014). Cold plasma in modified atmospheres for post-harvest treatment of strawberries. *Food and Bioprocess Technology*, 7(10), 3045–3054.
- Montiel, R., De Alba, M., Bravo, D., Gaya, P., & Medina, M. (2012). Effect of high pressure treatments on smoked cod quality during refrigerated storage. [Article] *Food Control*, 23(2), 429–436. <https://doi.org/10.1016/j.foodcont.2011.08.011>.
- Nyaisaba, B. M., Miao, W., Hatab, S., Siloam, A., Chen, M., & Deng, S. (2019). Effects of cold atmospheric plasma on squid proteases and gel properties of protein concentrate from squid (*Argentinus illex*) mantle. *Food Chemistry*, 291, 68–76.
- Olatunde, O. O., Benjakul, S., & Vongkamjan, K. (2019a). Combined effects of high voltage cold atmospheric plasma and antioxidants on the qualities and shelf-life of Asian sea bass slices. [Article] *Innovative Food Science & Emerging Technologies*, 54, 113–122. <https://doi.org/10.1016/j.ifset.2019.03.012>.
- Olatunde, O. O., Benjakul, S., & Vongkamjan, K. (2019b). High voltage cold atmospheric plasma: Antibacterial properties and its effect on quality of Asian sea bass slices. [Article] *Innovative Food Science & Emerging Technologies*, 52, 305–312. <https://doi.org/10.1016/j.ifset.2019.01.011>.
- Ozen, B. O., & Soyer, A. (2018). Effect of plant extracts on lipid and protein oxidation of mackerel (*Scomber scombrus*) mince during frozen storage. [Article] *Journal of Food Science and Technology-Mysore*, 55(1), 120–127. <https://doi.org/10.1007/s13197-017-2847-6>.
- Pankaj, S. K., Misra, N. N., & Cullen, P. J. (2013). Kinetics of tomato peroxidase inactivation by atmospheric pressure cold plasma based on dielectric barrier discharge. *Innovative Food Science & Emerging Technologies*, 19, 153–157. <https://doi.org/10.1016/j.ifset.2013.03.001>.
- Park, S. Y., & Ha, S. D. (2015). Application of cold oxygen plasma for the reduction of *Cladosporium cladosporioides* and *Penicillium citrinum* on the surface of dried filefish (*Stephanolepis cirrifer*) fillets. [Article] *International Journal of Food Science and Technology*, 50(4), 966–973. <https://doi.org/10.1111/ijfs.12730>.
- Pedrés-Garrido, S., Condón-Abanto, S., Beltrán, J. A., Lyng, J. G., Brunton, N. P., Bolton, D., et al. (2017). Assessment of high intensity ultrasound for surface decontamination of salmon (*S. salar*), mackerel (*S. scombrus*), cod (*G. morhua*) and hake (*M. merluccius*) fillets, and its impact on fish quality. *Innovative Food Science & Emerging Technologies*, 41, 64–70. <https://doi.org/10.1016/j.ifset.2017.02.006>.
- Pérez-Andrés, J. M., Charoux, C. M., Cullen, P., & Tiwari, B. K. (2018). Chemical modifications of lipids and proteins by nonthermal food processing technologies. *Journal of Agricultural and Food Chemistry*, 66(20), 5041–5054.
- Puligundla, P., Choi, S., & Mok, C. (2018). Microbial decontamination of gwamegi (Semi-dried Pacific Saury) using corona discharge plasma jet, including physicochemical and sensory evaluation. *Journal of Aquatic Food Product Technology*, 27(3), 274–283.
- Rode, T. M., & Hovda, M. B. (2016). High pressure processing extend the shelf life of fresh salmon, cod and mackerel. [Article] *Food Control*, 70, 242–248. <https://doi.org/10.1016/j.foodcont.2016.05.045>.
- Romotowska, P. E., Karlsdottir, M. G., Gudjonsdottir, M., Kristinsson, H. G., & Arason, S. (2016). Influence of feeding state and frozen storage temperature on the lipid stability of Atlantic mackerel (*Scomber scombrus*). [Article] *International Journal of Food Science and Technology*, 51(7), 1711–1720. <https://doi.org/10.1111/ijfs.13146>.
- Saldana, G., Alvarez, I., Condon, S., & Raso, J. (2014). Microbiological aspects related to the feasibility of PEF technology for food pasteurization. *Critical Reviews in Food Science and Nutrition*, 54(11), 1415–1426. <https://doi.org/10.1080/10408398.2011.638995>.
- Scholtz, V., Pazlarova, J., Souskova, H., Khun, J., & Julak, J. (2015). Nonthermal plasma—a tool for decontamination and disinfection. *Biotechnology Advances*, 33(6), 1108–1119.
- Schormüller, J. (1968). *Handbuch der Lebensmittelchemie*. Springer-Verlag.
- Segat, A., Misra, N. N., Cullen, P. J., & Innocente, N. (2015). Atmospheric pressure cold plasma (ACP) treatment of whey protein isolate model solution. [Article] *Innovative Food Science & Emerging Technologies*, 29, 247–254. <https://doi.org/10.1016/j.ifset.2015.03.014>.
- Shahidi, F., & Ambigaipalan, P. (2018). Omega-3 polyunsaturated fatty acids and their health benefits. In M. P. Doyle, & T. R. Klaenhammer (Vol. Eds.), *Annual review of food science and technology: Vol. 9*, (pp. 345–381). Palo Alto: Annual Reviews.
- Sharifian, A., Soltanzadeh, N., & Abbaszadeh, R. (2019). Effects of dielectric barrier discharge plasma on the physicochemical and functional properties of myofibrillar proteins. *Innovative Food Science & Emerging Technologies*, 54, 1–8.
- Silveira, M. R., Coutinho, N. M., Esmerino, E. A., Moraes, J., Fernandes, L. M., Pimentel, T. C., ... Cruz, A. G. (2019). Guava-flavored whey beverage processed by cold plasma technology: Bioactive compounds, fatty acid profile and volatile compounds. [Article] *Food Chemistry*, 279, 120–127. <https://doi.org/10.1016/j.foodchem.2018.11.128>.
- Simopoulos, A. P. (2002). The importance of the ratio of omega-6/omega-3 essential fatty acids. *Biomedicine & Pharmacotherapy*, 56(8), 365–379.
- Simopoulos, A. P. (2008). The importance of the omega-6/omega-3 fatty acid ratio in cardiovascular disease and other chronic diseases. *Experimental Biology and Medicine*, 233(6), 674–688.
- Surowsky, B., Schluter, O., & Knorr, D. (2015). Interactions of non-thermal atmospheric pressure plasma with solid and liquid food systems: A review. [Article; Proceedings Paper] *Food Engineering Reviews*, 7(2), 82–108. <https://doi.org/10.1007/s12393-014-9088-5>.
- Swanson, D., Block, R., & Mousa, S. A. (2012). Omega-3 fatty acids EPA and DHA: Health benefits throughout life. *Adv Nutr*, 3(1), 1–7. <https://doi.org/10.3945/an.111.000893>.
- Teixeira, B., Fidalgo, L., Mendes, R., Costa, G., Cordeiro, C., Marques, A., ... Nunes, M. L. (2014). Effect of high pressure processing in the quality of sea bass (*Dicentrarchus labrax*) fillets: Pressurization rate, pressure level and holding time. [Article] *Innovative Food Science & Emerging Technologies*, 22, 31–39. <https://doi.org/10.1016/j.ifset.2013.12.005>.
- Ulbricht, T., & Southgate, D. (1991). Coronary heart disease: Seven dietary factors. *The Lancet*, 338(8773), 985–992.
- Wang, J., Zhuang, H., Hinton, A., & Zhang, J. (2016). Influence of in-package cold plasma treatment on microbiological shelf life and appearance of fresh chicken breast fillets. *Food Microbiology*, 60, 142–146. <https://doi.org/10.1016/j.fm.2016.07.007>.
- Xiang, Q. S., Liu, X. F., Li, J. G., Ding, T., Zhang, H., Zhang, X. S., et al. (2018). Influences of cold atmospheric plasma on microbial safety, physicochemical and sensorial qualities of meat products. [Review] *Journal of Food Science and Technology-Mysore*, 55(3), 846–857. <https://doi.org/10.1007/s13197-017-3020-y>.
- Xu, L., Garner, A. L., Tao, B., & Keener, K. M. (2017). Microbial inactivation and quality changes in orange juice treated by high voltage atmospheric cold plasma. [Article] *Food and Bioprocess Technology*, 10(10), 1778–1791. <https://doi.org/10.1007/s11947-017-1947-7>.
- Yepez, X. V., & Keener, K. M. (2016). High-voltage Atmospheric Cold Plasma (HVACP) hydrogenation of soybean oil without trans-fatty acids. *Innovative Food Science & Emerging Technologies*, 38, 169–174. <https://doi.org/10.1016/j.ifset.2016.09.001>.
- Yong, H. I., Lee, H., Park, S., Park, J., Choe, W., Jung, S., et al. (2017). Flexible thin-layer plasma inactivation of bacteria and mold survival in beef jerky packaging and its effects on the meat's physicochemical properties. [Article] *Meat Science*, 123, 151–156. <https://doi.org/10.1016/j.meatsci.2016.09.016>.
- Ziuzina, D., Patil, S., Cullen, P. J., Keener, K. M., & Bourke, P. (2013). Atmospheric cold plasma inactivation of *Escherichia coli* in liquid media inside a sealed package. [Article] *Journal of Applied Microbiology*, 114(3), 778–787. <https://doi.org/10.1111/jam.12087>.
- Zong, G., Li, Y., Wanders, A. J., Alsema, M., Zock, P. L., Willett, W. C., ... Sun, Q. (2016). Intake of individual saturated fatty acids and risk of coronary heart disease in US men and women: Two prospective longitudinal cohort studies. *Bmj*, 355, i5796.