

2022

Modulation of Gut Microbiota by Bioactive Compounds for Prevention and Management of Type 2 Diabetes

Basista Rabina Sharma

CSIR-Central Food Technological Research Institute (CFTRI), India

Swarna Jaiswal

Technological University Dublin, swarna.jaiswal@tudublin.ie

P.V. Ravindra

CSIR-Central Food Technological Research Institute (CFTRI), India

Follow this and additional works at: <https://arrow.tudublin.ie/schfsehart>



Part of the [Medicine and Health Sciences Commons](#)

Recommended Citation


Rabina Sharma, B., Jaiswal, S. & Ravindra, P.V. (2020). Modulation of gut microbiota by bioactive compounds for prevention and management of type 2 diabetes. *Biomedicine & Pharmacotherapy*, vol. 152, no. 113148. doi:10.1016/j.biopha.2022.113148

This Article is brought to you for free and open access by the School of Food Science and Environmental Health at ARROW@TU Dublin. It has been accepted for inclusion in Articles by an authorized administrator of ARROW@TU Dublin. For more information, please contact arrow.admin@tudublin.ie, aisling.coyne@tudublin.ie, vera.kilshaw@tudublin.ie.



This work is licensed under a [Creative Commons Attribution-NonCommercial-No Derivative Works 4.0 International License](#).

Funder: Department of Biotechnology (DBT), and Science and Engineering Research Board (SERB), Govt. of India



Review

Modulation of gut microbiota by bioactive compounds for prevention and management of type 2 diabetes

Basista Rabina Sharma^a, Swarna Jaiswal^{b,c}, P.V. Ravindra^{a,*}

^a Department of Biochemistry, CSIR-Central Food Technological Research Institute (CFTRI), KRS Road, Opp. Rail Museum, Mysuru 570020, India

^b School of Food Science and Environmental Health, College of Sciences and Health, Technological University Dublin - City Campus, Central Quad, Grangegorman, Dublin D07 ADY7, Ireland

^c Environmental Sustainability and Health Institute, Technological University Dublin - City Campus, Grangegorman, Dublin D07 H6K8, Ireland



ARTICLE INFO

Keywords:

Type 2 diabetes
Gut microbiota
Eubiosis
Dysbiosis

ABSTRACT

Type 2 diabetes mellitus (T2DM) is a chronic metabolic disorder characterized by hyperglycemia and insulin resistance. Gut microbiota (GM) are specific groups of microbes colonized in the gastrointestinal (GI) tract. They profoundly influence health, disease protection, and associated with metabolic activities, and play a vital role in the production of functional metabolites from dietary substances. Dysbiosis of GM has been linked to the onset of T2DM and can be altered to attain eubiosis by intervention with various nutritional bioactive compounds such as polyphenols, prebiotics, and probiotics. This review presents an overview of the evidence and underlying mechanisms by which bioactive compounds modulate the GM for the prevention and management of T2DM.

1. Introduction

Diabetes is a chronic metabolic disorder characterized by hyperglycemia. The long-term effect of hyperglycemia causes the malfunctioning of several vital organs in the body, causing multiorgan failure and death [1,2]. According to the International Diabetes Federation (IDF), 463 million people were affected with diabetes in 2019, and the number is expected to increase up to 700 million by 2045 [3]. Diabetes is classified broadly into three types based on the underlying causes. Type 1 diabetes is characterized by the destruction and damage of β -cells due to autoimmune response, leading to suppression or absence of production of insulin [4]. Type 2 diabetes (T2DM) is characterized by hyperglycemia and loss of insulin sensitivity in insulin-responsive tissues such as skeletal muscle and adipose tissue [5]. Type 3 diabetes, known as gestational diabetes, is caused temporally during pregnancy or Alzheimer's disease [6]. T2DM is an expanding global public health and economic burden, closely linked to the macro and microvascular complications leading to cardiomyopathy, nephropathy, neuropathy, etc., causing death and significant economic loss [7]. T2DM is caused by interactions between intrinsic (genetic or hereditary), and extrinsic factors such as changes in the lifestyle, a lack of physical activity, and stress factors [5]. A choice of a healthy diet and regular physical activities forms a group of primary and long-term and sustainable, effective measures for the management

of T2DM. Additionally, depending upon the individual's age and physical fitness, lifestyle changes have been shown to complement antidiabetic treatment's efficacy and contribute to lowering the drug dosage [8–10] (Fig. 1).

The variety of gut functions is regulated by the interactions between bioactive compounds of the food and gut microbiota (GM). GM plays an essential role in controlling the fermentation and absorption of dietary nutrients such as SCFAs (Short Chain Fatty Acids) [11,12]. The GM is distributed throughout the GI (gastrointestinal) tract, and their type and activities are linked to various pathophysiological effects, including the development of T2DM [13]. Several metagenomic studies have found that dysbiosis in the GM directly influences the development of T2DM by affecting the gut permeability, inflammation, immune system, and energy metabolism [14–19]. Therefore, modulation of the GM by bioactive compounds has gained substantial momentum in recent years for finding suitable bioactive compounds or their formulation that have prophylactic and/or curative potential on T2DM. Several studies in the recent past described the cross-talk between diet, GM, and the host in the context of metabolic disorders such as diabetes, obesity, and cardiovascular diseases [20–24]. An article by Lazar et al., (2019) [22] highlights the importance of prebiotics and probiotics in modulating the GM and thus indicating their preventive effect on the onset of diabetes and obesity. Other studies have also shown that prebiotics and probiotics

* Corresponding author.

E-mail address: raviravindra1@gmail.com (P.V. Ravindra).




Fig. 1. The figure represents the relationship among the food-derived bioactives, gut microbiota, and risk factors associated with the development of insulin resistance and T2DM. The long-term consumption of the western diet, the overuse of antimicrobials (antibiotics), and contribution from host factors including age, sedentary lifestyle, etc., cause gut dysbiosis, which subsequently causes increased levels of bacterial lipopolysaccharide, -oxidative stress, -the release of pro-inflammatory cytokines, -gut inflammation, and gut permeability. All these changes over a period of time cause the development of insulin resistance and T2DM. On the other hand, bioactive compounds from a healthy diet positively influence the maintenance of gut eubiosis, which causes attenuation in the above-described pathological changes in the gut with concurrent improvement in insulin sensitivity and gut mobility, thus benefiting the prevention and management of T2DM.

have a favorable impact on eubiosis over dysbiosis and thus can be exploited to prevent and treat T2DM [20,22,25,27,28]. Further, previous studies also describe strategies of modulation of GM by antimicrobial agents, bariatric surgery and its implications in managing metabolic disorders [21]. On the other hand, studies also implicate how pharmacological, surgical and nutritional interventions for T2DM impact the GM [23]. This review article provides the update and summarizes the interaction between dietary bioactive compounds, GM, and T2DM. In addition, the article also focuses on three key questions: how different types of diet affect the GM populations, how changing the dietary pattern leads to gut dysbiosis linked to T2DM condition, and how bioactive compounds modulate the GM population help in preventing and managing the T2DM. Further, we have compiled most studies done in experimental models and humans based on the bioactive compounds from different sources modulating the gut population having anti-diabetic properties.

2. T2DM and diet interactions

A popular quote says, "you are what you eat". T2DM is caused by the cumulative effect of lifestyle changes and physical inactivity together with genetic susceptibility [26]. The incidence of T2DM is high in people who consumed a diet rich in high fat and sugar compared to the traditional diet [27]. Further, long-term eating of white rice [28], potatoes [29], fried foods [30], red and processed meat [31], sugar-sweetened beverages, and packaged fruit juices [32] increases the risk of getting T2DM. Therefore, consumption of healthy foods such as whole grains, nuts, yogurt, green leafy vegetables, fruits, coffee, green tea which contain bioactive compounds that lower down risk of T2DM has gained the attention of the public as well as researchers [17,33–37].

Furthermore, consumption of foods rich in $\Omega-6$ polyunsaturated fatty acids [38], dietary fiber [39], magnesium [40], zinc [41], vitamin D [42], selenium [43] lower risk of T2DM. Additionally, Koppe et al. (2005) [44] and other studies [45,46] have observed that moderate alcohol consumption can lower the risk of T2DM. Changing the dietary pattern, including lowering the fat consumption together with lifestyle modifications, can be an effective way for prevention as well as therapy for T2DM [27].

3. Gut microbiome

The gut microbiome refers to the set of genomes of all microorganisms in the gut. The GM refers to a specific microbial population in the gut [46,47]. The interplay between GM and host physiology influences the host's metabolic phenotype, stress response [48]. The equilibrium between the microbial diversity is a prerequisite to maintaining the host homeostasis, including energy metabolism, immunity response, inflammation, and brain-gut axis [19]. The dysbiosis in the GM may contribute to the onset of chronic conditions, including irritable bowel diseases, colorectal cancer, obesity, diabetes, metabolic syndrome, allergy, depression, and anxiety [21,49]. The factors that influence host intestinal microbiota are genetics, lifestyle, diet, and environmental factors [50]. Therefore, balancing the composition of GM is critical to preventing metabolic diseases [51–53].

3.1. Location, composition, and function

The microbiota is present on every surface of the human body, from the skin to genitourinary, GI, and respiratory tracts [51]. The GI tract is sterile at birth but gradually becomes colonized by various microbial

Table 1
Location, composition and function of GM.

GI region	Microbiota	Function of Microbiota	Reference
Stomach	Firmicutes: <i>Lactobacillus</i> , <i>Veillonella</i> , Proteobacteria: <i>Helicobacter</i>	Bacteroidetes: It digest complex polysaccharides, degraded complex O-glycans yielding Volatile Short-chain fatty acids (SCFAs) which is an energy source for host, regulating intestinal epithelial cell growth and differentiation, stimulates immune system	[51,54,57,103,108–114]
Duodenum	Firmicutes: <i>Bacilli</i> , <i>Streptococcaceae</i> , Actinobacteria: <i>Actinomycinaeae</i> , <i>Corynebacteriaceae</i>	Firmicutes: It release butyrate as end product and promote intestinal epithelial health, promote host immune homeostasis	
Jejunum	Firmicutes: <i>Bacilli</i> , <i>Streptococcaceae</i> , Actinobacteria: <i>Actinomycinaeae</i> , <i>Corynebacteriaceae</i>	Actinobacteria: some of them have probiotic effects, protect against pathogens through competitive exclusion, immune modulation, bile salt hydrolase activity, also have the ability to adhere to mucus as well as intestinal epithelium.	
Ileum	Firmicutes: <i>Bacilli</i> , <i>Streptococcaceae</i> , Actinobacteria: <i>Actinomycinaeae</i> , <i>Corynebacteriaceae</i>	Proteobacteria: less health benefit, serve as a potential diagnostic signature of dysbiosis and risk of diseases.	
Colon	Firmicutes: <i>Lachnospiraceae</i> , Bacteroidetes: <i>Bacteroidetes</i>		
Epithelial surface	Firmicutes: <i>Clostridium</i> , <i>Lactobacillus</i> , <i>Enterococcus</i>		
Mucus layer	Firmicutes: <i>Clostridium</i> , <i>Lactobacillus</i> , <i>Enterococcus</i>		
Intestinal lumen	Bacteroidetes: <i>Bacteroides</i> , Actinobacteria: <i>Bifidobacterium</i> , Firmicutes: <i>Streptococcus</i> , <i>Clostridium</i> , <i>Lactobacillus</i> , <i>Enterococcus</i> , <i>Ruminococcus</i> Proteobacteria: <i>Enterobacteriaceae</i>		

species and becomes the GM [13]. There is a notable spatial variation in the composition of GM in the GI tract. GM colonizes heavily in the GI tract, starting with the small concentration in the stomach followed by duodenum, jejunum, ileum, and colon (almost 70% of all microbes) as listed in Table 1. Human gut microbiota is anaerobic, and most belong to the phyla *Firmicutes* (~60%), *Bacteroidetes* (~15%), and *Actinobacteria* (~15%) followed by, *Verrucomicrobia* (~2%), *Proteobacteria* (~1%), and *Methanobacteriales* (~1%) [49,50]. Table 1 depicts the spatial distribution of the GM, along with their functions.

3.2. Effect of diet on GM

The human diet is very complex, having nutrients as well as food-derived bioactive compounds. Hence, dietary patterns are considered the key elements of human health. Changes in the dietary pattern influence GM's composition [56,57]. Alternative dietary patterns or miss management in the dietary component may harm the healthy microorganisms in GM. Several studies have revealed that diet-induced GM dysbiosis is linked to the onset of T2DM and obesity [19,49,58].

The components present in the diet affects GM by various mechanism, as shown in the Table 2. *Bifidobacteria*, *Clostridium*, and *Bacteroidetes* decrease if carbohydrate content in the diet decreases [59,60]. Similarly, increases in the dietary fiber content increase the short-chain fatty acid (SCFA) producing bacteria in the GM [61]. On the other hand, the western diet showed a more significant reduction in GM diversity due to lesser dietary fiber content [62]. Several studies have shown that the Mediterranean diet that is intrinsically rich in polyphenols has been found to lower the risk of T2DM and obesity [63,64]. Table 2 depicts the different types of dietary patterns and how they affect the population of healthy microbes in GM. Diet with high carbohydrates, fiber, protein has shown a positive impact on the person's health, as it acts as an energy source, helps in the production of SCFAs, production of microbial metabolites, and helps in increasing butyrate-producing bacteria. While, the keto diet, western diet, and high-fat diet cause a decrease in gut microbiota including *Bifidobacteria*, *Bacteroidetes* which will lead to the disease condition [59–61].

3.3. Diabetes and GM

Gut microbiota is an environmental factor responsible for controlling energy metabolism, body weight, proinflammatory activity, bile acid metabolism, insulin resistance, and modulating gut hormones [65,66]. The imbalance in the GM causes dysbiosis, which is associated with the onset of T2DM, insulin resistance, and obesity [19,23,24,53,67]. Several studies have revealed significant differences in the two most abundant phyla of GM i.e., *Firmicutes* and *Bacteroidetes*, where the ratio of *Firmicutes* to *Bacteroidetes* increased in people with diabetes as compared to healthy persons [67,68]. On the other hand, *Bifidobacteria*, *Akkermansia*, *Prevotella*, and *Faecalibacterium* have a positive effect on host metabolism and a negative influence on diabetes. Cani et al. (2007) [65] reported the contribution of *Bifidobacteria* towards the improvement of mucosal barrier function and reduction in low-grade chronic inflammation. *Bifidobacteria* also improves glucose tolerance and insulin sensitivity [69]. Hippe et al. (2016) [70] have found decreased levels *Faecalibacterium prausnitzii* in feces contributing to inflammation and poor glucose homeostasis. Similarly, Tap et al. (2015) [61] have studied the effect of *Prevotella* on the host diabetic phenotype, where it increased glucose metabolism by enhancing glycogen storage. Table 3 describes the effect of dysbiosis on diabetes and the contribution of the GM for controlling diabetes. Studies in mice show that alternation in GM can lead to diabetes. Some of the studies show that *Bifidobacteria*, *Bacteroidetes* contribute toward the improvement of mucosal barrier function, reduce inflammation, improve glucose tolerance, reduce oxidative stress which will ultimately help in controlling diabetes. Whereas, increase in *Lachnospiraceae* will contribute toward the development of T2DM, by increasing blood glucose levels and decreasing insulin levels in plasma. And studies in human shows that, *Bifidobacterium*, *Faecalibacterium prausnitzii*, *Akkermansia muciniphila*, *Prevotella* played a pivotal role in controlling diabetes. As shown in Table 3, the decreases in *Bifidobacterium*, *Faecalibacterium*, *Akkermansia*, *Prevotella* is associated with the diabetes condition.

Table 2
Effect of diet on GM.

Type of diet	Effect on microbiota	Mechanism	Reference
Low Carbohydrates	<i>Bifidobacteria</i> ↓ <i>Clostridium</i> ↓ <i>Bacteroidetes</i> ↓	It acts as energy source for microbiota and support microbial growth in the colon. It releases short-chain fatty acids that will help in regulation of gene expression by affecting host energy expenditure and storage.	[59,60]
High fiber	<i>Bacteroidetes</i> ↑ <i>Actinobacteria</i> ↑ <i>Firmicutes</i> to <i>Bacteroidetes</i> ratio↓	It help in production of SCFAs which will further regulate the expression of genes related to glycan and lipid metabolism	[61] [2,3]
High protein diet	<i>Bacteroidetes</i> ↑, <i>Lactobacillus</i> ↑, <i>Bifidobacteria</i> ↑, <i>Firmicutes</i> ↓	Microbiota digest the undigested protein and produced metabolites which are responsible for metabolism, immune system and nervous system.	[115–117]
High-fat diet	<i>Bacteroidetes</i> ↓ <i>Firmicutes</i> ↑ Proteobacteria ↑	It results in impairments in colonic epithelial integrity and barrier function which is associated with the development of Insulin resistance and type 2 diabetes. Increased plasma LPS levels	[118,119]
Keto diet	Dysmicrobism	As microbiota need carbohydrates as energy source, which are diminished in this type of diet	[120]
Western diet	<i>Clostridium innocuum</i> ↑, <i>Eubacterium dolichum</i> ↑, <i>Catenibacterium mitsuokai</i> ↑, <i>Enterococcus</i> spp. ↑ and <i>Bifidobacteria</i> ↓, <i>Bacteroidetes</i> ↓	High fat and digestible saccharides: mostly absorbed in duodenum, leaving very few substrate for colonic bacteria.	[62,118]
High Mediterranean diet	<i>Bacteroidetes</i> ↑ <i>Firmicutes</i> ↑ <i>Actinobacteria</i> ↑	Help in increasing beneficial butyrate-producing bacteria which impart a positive metabolic influence.	[9,63,64]

3.4. Mechanisms of GM on host metabolism

Gut microbiota is responsible for initiating, controlling, and manipulating the host's overall metabolic processes such as energy metabolism, metabolic endotoxemia, maintenance of gut permeability, and host immune system [66]. The interaction of GM and host metabolic changes linked to the onset of T2DM has been studied by various researchers and is presented in Table 4. The metabolites produced by the GM during the fermentation of complex carbohydrates i.e., SCFAs and bile acids, initiate several metabolic pathways that regulate glucose absorption, insulin sensitivity, and inflammation in the host [11,22,72,73]. There are three types of SCFAs i.e., acetate, propionate, and butyrate, produced by bacteria present in the colon [55]. *Bacteroidetes* phylum is known to produce acetate and propionate, while *Firmicutes* phylum produces butyrate by fermenting dietary fibers [74]. The SCFAs help to regulate energy expenditures, glucose homeostasis, and hepatic lipogenesis, which have a substantial impact on T2DM and obesity.

Gao et al. (2009) [75] have found that butyrate reduces insulin sensitivity in high-fat-fed diabetic mice by promoting energy expenditure and inducing mitochondria function. Acetate ingestion in T2DM rats reduces lipogenesis in adipose tissues and the liver. It improves glucose tolerance by inhibiting the transcription factor ChREBP (Carbohydrates-Responsive Element-Binding Protein), which is required for the conversion of glucose to fatty acids in the liver [76]. The diet-derived propionate improves the function of β -cells and promotes insulin secretion through the protein kinase C- dependent pathway [77]. Tolhurst et al. (2012) [78] have found that acetate and propionate influence the L-cells to release GLP-1 (Glucagon-Like Peptide-1) and PYY (Peptide YY hormone), which stimulate insulin secretion, glucose uptake in the muscle and decrease glucagon production in the pancreas. Understanding GM and host metabolism could be the potential strategy for controlling T2DM by modulating the GM of the host.

4. Bioactive compounds: relation with GM and T2DM

Bioactive compounds are phytochemicals such as polyphenols, anthocyanin, flavonoids, carotenoids, alkaloids, and tannins found in various plant parts, including leaf, bark, root, and have functional benefits to maintain normal health. Fruits and vegetables are also a substantial source of several bioactive compounds. Bioactive compounds like phenolics, flavonoids, alkaloids, anthocyanin possess significant antidiabetic properties by increasing α -glycosidase inhibiting activity, improving glucose tolerance by activating the P13K/Akt and inhibiting JNK in signaling, improving serum lipid levels, and reversing the insulin resistance by upregulating the expression levels of insulin receptor and glycolytic enzymes and by down regulating the expression

of insulin receptor substrate-1 and gluconeogenesis enzymes [79]. Foods loaded with bioactive compounds have to pass through various metabolic processes in the body where it finds the space to macerate themselves and release the potential bioactive compounds. Microorganisms are always considered as a sustainable approach to macerate phytochemicals from the foods by fermentation process [80]. As the GI tract contains trillions of GM, the GI tract suits as an appropriate site where the bioactive compounds can be extracted after intake and utilized prominently in several metabolic processes [81]. Several studies have shown a two-way relationship between bioactive compounds and GM [82]. GM plays a crucial role in modulating the production, bioavailability, and biological activities of bioactive compounds, particularly after the intake of food containing high-molecular-weight polyphenols. Furthermore, bioactive compounds might be converted by the colonic GM to metabolites such as SCFAs and bile acids that can affect the intestinal ecology and influence host health by participating in the several metabolic pathways [11,12]. On the other hand, the bioactive compounds supplied to the GM by the various dietary pattern modulates the composition of GM, due to the action of aromatic metabolites [82–85]. In recent years, research focus has shifted towards the modulation of GM by bioactive compounds to combat metabolic diseases such as diabetes and obesity [69,86–90].

Studies have shown that bioactive compounds modulate the composition of microflora either by selective prebiotic effects or antimicrobial effects against the pathogenic bacteria in the gut [82,91–94]. The modulation of colonic microbiota may contribute to control the T2DM, although the mechanisms regarding the antidiabetic action have not been clearly explained. Studies [95–98] have found that allucin modulates the *Firmicutes* to *Bacteroidetes* ratio, consequently maintain glucose homeostasis and insulin sensitivity. Saponin reduces the *Firmicutes* to *Bacteroidetes* ratio, which again beneficial in controlling T2DM [99]. Protocatechuic acid elevated GLP-1 secretion, and serum insulin, also improved insulin resistance by reducing the *Firmicutes* to *Bacteroidetes* ratio [100]. The contradictory results have been found by other studies [101] where the flavonoids from green seaweed increased the *Firmicutes* to *Bacteroidetes* ratio but decreased blood glucose level and improved insulin sensitivity, which might be due to other microflora niches in the ecology of GM [82].

Bifidobacterium spp. enhances the expression of proteins involved in the insulin-signaling pathway and expression of adiponectin hormone, which regulates the blood glucose level and decreases inflammatory adipokines [102]. Cani et al. (2007) [65] reported that *Bifidobacterium* spp. significantly increased the extraction of energy and fat storage, also reduced the expression levels of TNF- α and LPS (lipopolysaccharides) in diabetic models. In addition, *Bifidobacterium* spp. reduced endotoxemia and inflammation and also improved glucose tolerance and insulin

Table 3
GM dysbiosis and interlinkage with diabetes.

Change in GM	Effect on diabetes	References
Studies in Experimental mice		
<i>Firmicutes</i> ↓, <i>Bacteroidetes</i> ↑, <i>Bifidobacteria</i> ↑ <i>Bifidobacteria</i> ↑	<ul style="list-style-type: none"> improved glucose tolerance, help in secretion of GLP-1 from L-cells reduction of low grade inflammation and oxidative stress. 	[121]
<i>Lachnospiraceae</i> ↑	<ul style="list-style-type: none"> contribute toward the improvement of mucosal barrier function by increasing the production of endogenous GLP-2 improved tight junctions decreased plasma LPS concentrations reduction in low grade inflammation oxidative stresses contribute toward the development of type 2 diabetes by increasing fasting blood glucose levels, - decreasing insulin in plasma. 	[65]
Studies in human		
<i>Firmicutes</i> (<i>Clostridia</i>) ↓ <i>Bacteroidetes</i> ↑ <i>Proteobacteria</i> ↑, <i>Bacteroides</i> - versus class <i>Clostridia</i> and <i>C. coccoides</i> ↑ <i>Bifidobacterium</i> spp. ↓ <i>Clostridium leptum</i> ↑, <i>Bacteroides vulgatus</i> ↓	<ul style="list-style-type: none"> Gram negative bacteria, made up of Lipopolysaccharides, which is known as a potent stimulator of inflammation and cause diabetes. <p>Role of <i>Bifidobacterium</i> spp. in host metabolism:</p> <ul style="list-style-type: none"> improved glucose tolerance, mucosal barrier function, glucose-induced insulin and normalized inflammatory tone which lead to diabetes. 	[123,124]
<i>Faecalibacterium prausnitzii</i> ↓	<p>Role of <i>Faecalibacterium prausnitzii</i> in host metabolism:</p> <ul style="list-style-type: none"> Block NF-κB activation Inhibit secretion pro-inflammatory mediators Improve glucose and lipid homeostasis 	[65,69]
<i>Clostridiales</i> sp. SS3/4 ↓, <i>Eubacterium rectale</i> ↓, <i>Faecalibacterium prausnitzii</i> ↓, <i>Roseburia intestinalis</i> ↓, <i>Roseburia inulinivorans</i> ↓, <i>Bacteroides caccae</i> ↑, <i>Clostridium hathewayi</i> ↑, <i>Clostridium ramosum</i> ↑, <i>Clostridium symbiosum</i> ↑, <i>Eggerthella lenta</i> ↑ and <i>E. coli</i> ↑	<p>This led to type 2 diabetes, as this cultures is mainly:</p> <ul style="list-style-type: none"> associated with membrane transport of sugars, transportation of branched-chain amino transport, metabolism of methane, degradation of xenobiotics, metabolism and reduction of sulfate 	[70,109, 125]
<i>Akkermansia muciniphila</i> ↓ <i>Clostridiaceae</i> ↑, <i>Enterococcus casseliflavus</i> ↑, <i>Butyrivibrio</i> ↓, <i>Bifidobacterium</i> ↓, <i>Megasphaera</i> ↓	<p>Role of <i>Akkermansia muciniphila</i> (mucin degrading bacteria) in host metabolism</p> <ul style="list-style-type: none"> stimulates the secretion of glucagon like peptide from the L-cells that suppress glucagen secretion, improve intestinal permeability, ingestion of carbohydrate, improved insulin secretion, Protect against pathogens. <p>SCFAs producing bacteria:</p> <ul style="list-style-type: none"> improved insulin secretion. <p>Role of <i>Bifidobacterium</i> in host phenotype:</p> <ul style="list-style-type: none"> improve of mucosal barrier function improved tight junctions decreased plasma LPS concentrations reduction in low grade inflammation oxidative stresses 	[126]
<i>Clostridium clostridioforme</i> ↑, <i>Lactobacillus gasseri</i> ↑, <i>Streptococcus mutans</i> ↑	<p>Role of <i>Akkermansia muciniphila</i> (mucin degrading bacteria) in host metabolism</p> <ul style="list-style-type: none"> stimulates the secretion of glucagon like peptide from the L-cells that suppress glucagen secretion, improve intestinal permeability, ingestion of carbohydrate, improved insulin secretion, Protect against pathogens. <p>SCFAs producing bacteria:</p> <ul style="list-style-type: none"> improved insulin secretion. <p>Role of <i>Bifidobacterium</i> in host phenotype:</p> <ul style="list-style-type: none"> improve of mucosal barrier function improved tight junctions decreased plasma LPS concentrations reduction in low grade inflammation oxidative stresses lead to diabetes, mainly associated with fasting glucose, HbA1c, insulin plasma triglycerides. 	[127,128]
<i>Prevotella</i> ↑	<p>Increase in <i>Prevotella</i>:</p> <ul style="list-style-type: none"> lead to diabetes, mainly associated with fasting glucose, HbA1c, insulin plasma triglycerides. 	[129]
<i>Firmicutes</i> ↓, <i>Bacteroidetes</i> ↑, <i>Bifidobacteria</i> ↑	<p>Increase in <i>Prevotella</i>:</p> <ul style="list-style-type: none"> lead to diabetes, mainly associated with fasting glucose, HbA1c, insulin plasma triglycerides. 	[61]
	<ul style="list-style-type: none"> improve glucose metabolism by promoting glycogen storage contribute toward the production of butyrate which further help in the secretion of GLP-1 from L-cells that changes the metabolic function 	[130]

↑ shows increase in GM, ↓ shows decrease in GM

secretion by promoting the synthesis and secretion of incretin and GLP-1 [103]. Hence, the increase in the *Bifidobacterium* spp. by certain bioactive compounds helps to control T2DM. Studies have been reported that berberine, polyphenols, phenolic acids such as gallic acid and ellagic acid increase the levels of *Bifidobacterium* spp. which helps to maintain

T2DM (Table 5). Similarly, *Lactobacillus* spp. Improved glucose tolerance and normalized inflammatory tone by reducing LPS and also increased the production of SCFAs and inhibited the pathogenic gut microflora, which might be beneficial for maintaining the T2DM [104, 105]. Table 5 depicts that the various bioactive compounds increase the

Table 4
Effect of GM on host diabetic phenotype.

Function of GM	Mechanism	Effect on diabetes	References
Energy metabolism	SCFA-GPR41/43 Pathway	Gut microbiota ferment indigestible complex carbohydrates and produced monosaccharides, secondary bile acids and short chain fatty acids. SCFAs include: butyrate (activate GPR41), propionate (activate both GPR41 and GPR43), and acetate (activate GPR43). SCFAs act as a signaling molecules and bind to the G-protein coupled receptors GPR43 and GPR41 and played a major role by suppressing insulin signaling and decreased storage of lipid in the adipocytes, increase the oxidation of lipids which will lead to increase in energy expenditure.	[131] [7]
	ChREBP/SREBP-1c Pathway	GM regulate the lipogenesis in liver (production of triglyceride) by controlling two transcription factors i.e., carbohydrate response element binding protein (ChREBP) and sterol response element binding protein 1c (SREBP-1c) which improves glucose absorption and insulin levels.	[67,132]
	AMPK Pathway	GM help to increase the AMPK activity by production of SCFAs. AMPK inhibits lipid and glycogen synthesis and stimulates glucose uptake, glycolysis, fatty acid oxidation.	[75, 133–135]
	FIAF Pathway	GM influence the host metabolism by suppressing the expression of fasting –induced adipose factor (FIAF) which ultimately arrest the Microbiota influence the host by suppressing the expression of fasting –induced adipose factor (FIAF) which will further suppress the lipoprotein lipase (LPL) inhibitor. Hereby increase the activity of LPL will improve the uptake of fatty acids and triacylglycerol accumulation in adipocytes.	[136,137]
Metabolic endotoxemia	Low grade inflammation	High LPS level will form a complex (LPS- binding proteins & the CD14 co-receptor) which is recognized by Toll-like receptor 4, triggering an inflammatory response through Complex signaling pathway by the activation of NF-kB pathway and subsequently expression of proinflammatory cytokines. The activation of NF-kB pathway may also have impaired insulin signaling, which can lead to insulin resistance. LPS also creates a metabolic endotoxemia that cause damage to the mucosal integrity and increase intestinal permeability, also reducing the expression of epithelial tight junction proteins.	[137,138]
Gut permeability	GLP-2 dependent mechanism	Microbiota will help in increasing the production of endogenous GLP-2 and improved the gut barrier functions by a GLP-2 dependent mechanism and hence improves the gut barrier function.	[139,140]
Immune system	Toll-like receptor 5	Toll-like receptor 5 (TLR5) is a component that activated the innate immune system which help to fight against the infection, exhibit hyperphagia, hyperlipidemia, hypertension, increased adiposity and insulin resistance	[141,142]

level of *Lactobacillus* spp. in GM to control the T2DM. Table 5 shows that allicin, flavanols, alkaloids, betacyanins, protocatechuic acid, phlorizin, polyphenols increases the *Akkermansia* spp. in the GM. The antidiabetic action of *Akkermansia* spp. has been described by various researchers. It is a mucin degrading bacteria, helps in preserving the mucus layer thickness, thereby decreasing gut permeability and leakage of LPS [106]. It also improves insulin resistance by inhibiting JNK1 and activating P13K signaling pathways [107]. Xie et al. (2018) [100] also noticed that *Akkermansia* spp. promotes the secretion of GLP-1 from the L-cells to suppress glucagon secretion.

Furthermore, a number of microflora in the GM are modulated by the particular bioactive compounds, which is presented in Table 5. The potential antidiabetic action of GM modulation by bioactive compounds is attributed to increases in the levels of *Bifidobacterium* spp. *Lactobacillus* spp. *Akkermansia* spp., as well as to the reduction in the *Firmicutes* to *Bacteroidetes* ratio. Hence, the particular bioactive compounds having the ability to modulate the gut ecosystem could be beneficial for use as preventive and therapy of T2DM.

The major predictor of human gut microbiota composition is diet. Scientific evidence supports the importance of diet and its composition in modulating gut microbiota. The long-term consumption of the western diet, the overuse of antimicrobials (antibiotics), and host factors including age, sedentary lifestyle, etc., cause gut dysbiosis which, subsequently causes increased levels of bacterial lipopolysaccharide, oxidative stress, the release of pro-inflammatory, gut inflammation, and permeability. All these changes over a period of time cause the development of insulin resistance and T2DM. On the other hand, bioactive compounds from the healthy diet positively influence the maintenance of gut eubiosis, which causes attenuation in the above-described pathological changes in the gut with concurrent improvement in insulin sensitivity and gut mobility, thus benefiting prevention and management of T2DM. In addition to these factors, the influence of specific micronutrients in the diet, frequency in consumption of western diet etc. on GM requires further investigations. Therefore, more studies are required to elucidate the complex interactions between bioactive compounds in the diet, GM, and on the onset and progression as well as prevention and management of T2DM.

5. Conclusion

T2DM is expanding as a most dangerous metabolic disorder after

obesity, which invites various health-related risks. The global health organizations and disease control boards have declared the severity of T2DM and advised to take preventive and curative therapeutic measures to control this metabolic disorder. Several research studies conducted in past decades proved that T2DM is mainly caused by a combination of unhealthy dietary patterns and relative inactivity of insulin. The systemic investigations concerning diet-induced T2DM conclude on the importance of GM and its role to prevent T2DM. Various species of GM like *Firmicutes*, *Bacteroidetes*, *Lactobacillus*, and *Bifidobacterium* play a vital role to control T2DM. Several scientific evidence stated that lifestyle changes and dietary patterns might affect the healthy GM, which causes GM dysbiosis, responsible for the T2DM. Recently researchers have proved that diet-induced GM dysbiosis can be solved by incorporating dietary supplements such as polyphenols, prebiotics, and bioactive compounds. Several systematic studies have recently supported the antioxidative and antimicrobial effects of bioactive compounds on GM, ultimately increasing the ratio of healthy GM. Various in vitro and in vivo studies proved that GM's modulation by bioactive compounds helps prevent T2DM by improving glucose homeostasis, improving insulin sensitivity, and increasing the production of short-chain fatty acids. However, more studies are required to elucidate the intricate relationship between dietary factors including the role of specific micro and macronutrients, combination of bioactive compounds, mechanism of their action on GM, interactions among GM, the role of microbial metabolites, bioaccessibility, and bioavailability of bioactive compounds, the host factors, such as age, gender, stress, etc., and their impact on the onset as well as prevention and management of T2DM.

CRedit authorship contribution statement

Basista Rabina Sharma: Literature search, Writing – original draft. **Swarna Jaiswal:** Addition of literature, editing and language and grammatical corrections. **P.V. Ravindra:** Conceptualization, Supervision, Final draft preparation.

Data availability

Data will be made available on request.

Table 5
Modulation of GM by bioactive compounds for controlling the T2DM.

Bioactive compounds	Sources	Effect on GM	Anti-diabetic mechanisms	References
Anthocyanin	Strawberry	<i>Bifidobacterium</i> ↑, <i>Lactobacillus</i> ↑, <i>Verrucomicrobia</i> ↓,	<ul style="list-style-type: none"> Improved lipid, glucose metabolism, insulin resistance, gut barrier function, reduce low grade inflammation and stimulate the host immune system 	[143]
	Blueberries, Cherries,	<i>Bifidobacterium</i> spp. ↑, <i>Lactobacillus- Enterococcus</i> spp. ↑	<ul style="list-style-type: none"> Decreased blood glucose level Increased the expression of adiponectin Decreased Inflammatory adipokines 	[102,144, 145]
Hesperdin and naringin	Citrus fruit	<i>Bifidobacterium</i> spp. ↑, <i>Lactobacillus</i> spp. ↑, <i>Enterobacteria</i> ↓	<ul style="list-style-type: none"> Improved gut barrier function Improved glucose homeostasis Improved insulin sensitivity Increase the production of SCFAs 	[105]
Flavonoids	<i>E. proliferata</i>	<i>Bacteroidetes</i> ↑, <i>Firmicutes</i> ↓, <i>Lachnospiraceae</i> group and <i>Alisties</i> ↑,	<ul style="list-style-type: none"> Decreased blood glucose levels Improve glucose tolerance (by activating the P13K/Akt and inhibiting JNK in signaling pathway) 	[101] [4,5]
1-Deoxyojirimycin (Alkanoids)	Mulberry Leaves	<i>Lactobacillus</i> ↑, <i>Lachnospiraceae</i> NK4A136 group ↑, <i>Oscillibacter</i> ↑, <i>Alistipes</i> ↑, <i>Lachnospiraceae</i> ↑, <i>Bifidobacterium</i> ↑, <i>Ruminococcaceae</i> UCG-014 ↓, <i>Weissella</i> ↓, <i>Ruminococcus</i> ↓, <i>Prevotellaceae</i> Ga6A1 group ↓, <i>Anaerostipes</i> ↓, <i>Klebsiella</i> ↓, <i>Prevotellaceae</i> UCG-001 ↓ and <i>Bacteroidales</i> S24–7 group ↓	<ul style="list-style-type: none"> Decreased serum glucose and insulin levels Improved serum lipid levels and reversed insulin resistance by upregulating the protein expression of insulin receptor glycolysis enzymes (GK, PK and PFK) and by down regulating the protein expression of insulin receptor substrate-1 and gluconeogenesis enzymes (PCB, PEPCK, FBPase and G-6-Pase). 	[146]
Alkaloids, polyphenols	Mulberry leaves and Oat bran	<i>Firmicutes: Bacteroidetes</i> ↑, <i>Bacteroidales</i> S-247 group ↓, <i>Escherichia Shigella</i> ↓, <i>Alistipes</i> ↑, <i>Ruminiclostridium</i> ↑	<ul style="list-style-type: none"> Improved glucose metabolism by regulating insulin receptor and GK expressions and by down regulating PCB, PEPCK, PDKC expression. 	[147]
Berberine		<i>Bifidobacterium</i> spp. ↑	<ul style="list-style-type: none"> Increased the extraction of energy and fat storage Reduced the expression levels of TNF-α and LPS Reduced endotoxemia and inflammation, Improved glucose tolerance and insulin secretion by promoting the synthesis and secretion of incretin and GLP-1. 	[65,148]
Polyphenols	Cranberry	<i>Akkermansia</i> ↑	Mucin degrading bacteria: <ul style="list-style-type: none"> Preserved the mucus layer thickness 	[106]
	Enteromorpha proliferata	<i>Akkermansia</i> ↑, <i>Alistipes</i> and <i>Turicinbacter</i> ↓	<ul style="list-style-type: none"> Decreasing gut permeability & Leakage of LPS Improved insulin resistance by inhibiting JNK1 and activating P13K signaling pathway 	[107]
	Wild Blueberry	<i>Bifidobacterium</i> ↑, <i>Lactobacillus acidophilus</i> ↑, <i>Eubacteria</i> ↑	<ul style="list-style-type: none"> Decreased blood glucose level Increased the expression of adiponectin Decreased Inflammatory adipokines Improved gut barrier function 	[102,149]
	Green tea, Oolong tea, black tea	<i>Bifidobacterium</i> spp. ↑, <i>Lactobacillus- Enterococcus</i> spp. ↑, <i>Bacteroides</i> ↓, <i>Clostridium Histolyticum</i> ↓	<ul style="list-style-type: none"> Increase the production of SCFAs Inhibit pathogen species Improved glucose tolerance and insulin secretion normalized inflammatory tone. 	[104,150, 151]
	Red wine	<i>Bifidobacterium</i> spp. ↑, <i>Lactobacillus</i> spp. ↑,	<ul style="list-style-type: none"> Improved glucose tolerance and insulin secretion Normalized inflammatory tone Increase the production of SCFAs, Inhibit the adhesion of pathogen. 	[104,152]
Allicin	Garlic	<i>p-Firmicutes: p-Bacteroidetes</i> ↓, <i>f-Lachnospiraceae</i> ↑, <i>g-Akkermansia</i> ↑, <i>g-Lactobacillus</i> ↑, <i>Lachnospiraceae</i> ↑, <i>Ruminococcaceae</i> ↑	<ul style="list-style-type: none"> Maintains glucose homeostasis, Improves insulin sensitivity, Regulate glucose metabolism 	[95–98]
Berberine	European barberry, goldenseal, goldthread, Oregon grape, and tree turmeric	<i>Blautia</i> ↑, <i>Allobaculum</i> ↑, <i>Streptococcus</i> ↓, <i>Prevotella</i> ↓	<ul style="list-style-type: none"> Improves insulin secretion by producing SCFA such as acetate and propionate 	[78, 153–155]
β-glucan	Oats, Barley, Mushroom	<i>g-Bifidobacterium</i> ↑, <i>g-Blautia</i> ↑, <i>p-Actinobacteria</i> ↑	<ul style="list-style-type: none"> Increase the production of SCFAs such as acetate, propionate which influence the L-cells to release the peptide GLP-1 and PYY Stimulate insulin secretion Increase glucose tolerance Increase insulin secretion 	[17,78,156, 157]
Flavonols	Green tea leaves, Red wine, Cocoa, Cranberry	<i>Lactobacillus</i> ↑, <i>Bifidobacterium</i> ↑, <i>p-Bacteroidetes</i> ↑, <i>Akkermansia</i> ↑	<ul style="list-style-type: none"> Increase insulin secretion 	[106,158, 159]
Baicalein	Tuber aestivum, Oroxylym indicum, Scutellaria baicalensis, Scutellaria lateriflora	<i>Bacteroidales</i> ↑, <i>Bacteroides</i> ↑, <i>Prevotellaceae</i> ↑, <i>Alloprevotella</i> ↑, <i>Butyricimonas</i> ↑, <i>Arabacteroide</i> ↑,	<ul style="list-style-type: none"> Increase the production of SCFAs Improved insulin resistance, lipid metabolism, Decreased blood glucose, Decreased circulating LPS and systemic inflammation 	[160,161]
Catechin	Rhododendron groenlandicum	<i>Clostridium perfringens</i> ↓, <i>Clostridium difficile</i> ↓, <i>Lactobacillus- Enterococcus</i> spp. ↑, <i>Bifidobacterium</i> spp. ↑	<ul style="list-style-type: none"> Reduced intestinal endotoxin levels Improve mucosal barrier function Lower serum glucose Increase expression of insulin signaling proteins Improved adipokine 	[65,102, 150,162, 163]

(continued on next page)

Table 5 (continued)

Bioactive compounds	Sources	Effect on GM	Anti-diabetic mechanisms	References
Alkaloids	Rhizomz Coptidis	<i>Sporobacter termitidis</i> ↑, <i>Alcaligenes faecalis</i> ↑, <i>Akkermansia muciniphila</i> ↑, <i>E. coli</i> ↓, <i>Desulfovibrio C21_c20</i> ↓, <i>Parabacteroides</i> ↓	Improved: <ul style="list-style-type: none"> • Insulin sensitivity • Glucose homeostasis • Blood lipids 	[164]
Gallic acid, ellagic acid	Pomegranate by-product	<i>Bifidobacterium</i> spp. ↑, <i>Lactobacillus</i> spp. ↑, <i>Clostridium coccooides</i> ↓: <i>Eubacterium rectale</i> group and <i>C. Histolyticum</i> group	<ul style="list-style-type: none"> • Increase the production of SCFAs • Inhibit pathogenic species • Increased the activity of α-glucosidase • Improved glucose tolerance • Improved insulin secretion. 	[104,165]
Ellagitannins	Pomegranate	<i>Bifidobacterium</i> spp. ↑, <i>Lactobacillus</i> spp. ↑, <i>B. fragilis</i> ↓, <i>Clostridia</i> ↓, <i>Enterobacteriaceae</i> ↓	<ul style="list-style-type: none"> • Inhibits LPS induced inflammation • Improved glucose tolerance & insulin secretion, • Normalized inflammatory tone • Increase the production of SCFAs • Inhibit the adhesion of pathogen. 	[166–168]
Betacyanins	Red pitaya	<i>Akkermansia</i> ↑, <i>Firmicutes/ Bacteroidetes</i> ↓	<ul style="list-style-type: none"> • Improved Hepatic steatosis • Improved insulin resistance • stimulates the secretion of glucagon like peptide from the L-cells that suppress glucagen secretion 	[127,169]
Saponin	Polygonatum kingianum	<i>Firmicutes</i> ↑, <i>Bacteroidetes</i> ↓, <i>Proteobacteria</i> ↓	<ul style="list-style-type: none"> • Decreased blood glucose and LPS, • Improved glucose metabolism • Increase the production of SCFAs 	[99]
Protocatechuic acid	Alpinia oxyphylla	<i>Bacteroidetes</i> to <i>Firmicutes</i> ↑, <i>Akkermansia</i> ↑, <i>Helocobacter</i> ↓	<ul style="list-style-type: none"> • Elevated GLP-1 secretion & serum insulin • Improved insulin resistance & renal function • decreased blood glucose level 	[100]
Triterpenoid	Potentilla discolor Bunge	<i>Firmicutes / Bacteroidetes</i> ↓, <i>Proteobacteria</i> ↑	<ul style="list-style-type: none"> • Improves Insulin resistance, • Protect pancreatic β-cells • decreased LPS, • decreased pro-inflammatory cytokines, Reduced expression of TLR4. • Increased expression of GPR41 & GPR43 	[170,175]
Phlorizin	Apple, Strawberry, pomegranate	<i>Akkermansia muciniphila</i> ↑, <i>Prevotella</i> ↑	Prevent metabolic syndrome, Decreased: <ul style="list-style-type: none"> • weight gain • energy intake • serum LPS • insulin resistance 	[171]
Lycopene	Tomato, Moringa oleifera	<i>Proteobacteria</i> ↓, <i>Bifidobacterium</i> ↑, <i>Lactobacillus</i> ↑	Improved: <ul style="list-style-type: none"> • LPS-induced insulin resistance Mitochondrial dysfunction, • Hepatic inflammation, • Gut barrier function 	[172–174, 176]

↑ shows increase in GM, ↓ shows decrease in GM.

Acknowledgments and funding support

The authors are thankful to the Director, CSIR-CFTRI for providing facilities to carry out this work. The corresponding author acknowledges the financial support from the Department of Biotechnology (DBT), and Science and Engineering Research Board (SERB), Govt. of India.

Conflict of interest statement

We declare no competing interests.

Data availability

Data supporting this systematic review are from previously reported studies and datasets, which have been cited at appropriate places in the text.

References

- [1] A. Nanditha, R.C. Ma, A. Ramachandran, et al., Diabetes in Asia and the Pacific: implications for the global epidemic, *Diabetes Care* 39 (3) (2016) 472–485.
- [2] Y. Zheng, S.H. Ley, F.B. Hu, Global aetiology and epidemiology of type 2 diabetes mellitus and its complications, *Nat. Rev. Endocrinol.* 14 (2) (2018) 88.
- [3] P. Saeeedi, I. Petersohn, P. Salpea, et al., Global and regional diabetes prevalence estimates for 2019 and projections for 2030 and 2045: results from the International Diabetes Federation Diabetes Atlas, *Diabetes Res. Clin. Pract.* 157 (2019), 107843.
- [4] M.A. Atkinson, G.S. Eisenbarth, A.W. Michels, Type 1 diabetes, *Lancet* 383 (9911) (2014) 69–82.
- [5] R.A. DeFronzo, E. Ferrannini, L. Groop, et al., Type 2 diabetes mellitus, *Nat. Rev. Dis. Prim.* 1 (1) (2015) 1–22.
- [6] R. Kandimalla, V. Thirumala, P.H. Reddy, Is Alzheimer's disease a type 3 diabetes? A critical appraisal, *Biochim. Biophys. Acta (BBA) Mol. Basis Dis.* 1863 (5) (2017) 1078–1089.
- [7] I.I. Iid, S. Kumar, S. Shukla, V. Kumar, R. Sharma, Putative antidiabetic herbal food ingredients: nutra/functional properties, bioavailability and effect on metabolic pathways, *Trends Food Sci. Technol.* 97 (2020) 317–340.
- [8] L. Azadbakht, N.R.P. Fard, M. Karimi, et al., Effects of the Dietary Approaches to Stop Hypertension (DASH) eating plan on cardiovascular risks among type 2 diabetic patients: a randomized crossover clinical trial, *Diabetes Care* 34 (1) (2011) 55–57.
- [9] B. Karamanos, A. Thanopoulou, E. Anastasiou, et al., Relation of the Mediterranean diet with the incidence of gestational diabetes, *Eur. J. Clin. Nutr.* 68 (1) (2014) 8–13.
- [10] J. Salas-Salvadó, M. Martínez-González, M. Bullo, E. Ros, The role of diet in the prevention of type 2 diabetes, *Nutr. Metab. Cardiovasc. Dis.* 21 (2011) B32–B48.
- [11] D.J. Morrison, T. Preston, Formation of short chain fatty acids by the gut microbiota and their impact on human metabolism, *Gut Microbes* 7 (3) (2016) 189–200.
- [12] D. Ríos-Covián, P. Ruas-Madiedo, A. Margolles, M. Gueimonde, C.G. De Los Reyes-gavilán, N. Salazar, Intestinal short chain fatty acids and their link with diet and human health, *Front. Microbiol.* 7 (2016) 185.
- [13] G.P. Donaldson, S.M. Lee, S.K. Mazmanian, Gut biogeography of the bacterial microbiota, *Nat. Rev. Microbiol.* 14 (1) (2016) 20–32.
- [14] A. Vacharaksa, B. Finlay, Gut microbiota: metagenomics to study complex ecology, *Curr. Biol.* 20 (13) (2010) R569–R571.
- [15] J.-C. Lagier, F. Armougom, M. Million, et al., Microbial culturomics: paradigm shift in the human gut microbiome study, *Clin. Microbiol. Infect.* 18 (12) (2012) 1185–1193.

- [16] R.S. Mandal, S. Saha, S. Das, Metagenomic surveys of gut microbiota, *Genom. Proteom. Bioinform.* 13 (3) (2015) 148–158.
- [17] F. Wang, M. Miao, H. Xia, L.-G. Yang, S.-K. Wang, G.-J. Sun, Antioxidant activities of aqueous extracts from 12 Chinese edible flowers in vitro and in vivo, *Food Nutr. Res.* (2016).
- [18] V. Tremaroli, F. Bäckhed, Functional interactions between the gut microbiota and host metabolism, *Nature* 489 (7415) (2012) 242–249.
- [19] G. Rogers, D. Keating, R. Young, M. Wong, J. Licinio, S. Wesselingh, From gut dysbiosis to altered brain function and mental illness: mechanisms and pathways, *Mol. Psychiatry* 21 (6) (2016) 738–748.
- [20] R.H. Mallappa, C. Balasubramaniam, M.R. Amarlapudi, S. Kelkar, G.A. Adewumi, S. Kadyan, D. Pradhan, S. Grover, Role of probiotics in the prevention and management of diabetes and obesity. Probiotics in the Prevention and Management of Human Diseases, Academic Press, 2022, pp. 321–336.
- [21] O.O. Erejuwa, S.A. Sulaiman, M.S.A. Wahab, Modulation of gut microbiota in the management of metabolic disorders: the prospects and challenges, *Int. J. Mol. Sci.* 15 (3) (2014) 4158–4188.
- [22] V. Lazar, L.M. Ditu, G.G. Pircalabioru, A. Picu, L. Petcu, N. Cucu, M.C. Chifiriuc, Gut microbiota, host organism, and diet triologue in diabetes and obesity, *Front. Nutr.* 6 (2019) 21.
- [23] N.M. Delzenne, P.D. Cani, A. Everard, A.M. Neyrinck, L.B. Bindels, Gut microorganisms as promising targets for the management of type 2 diabetes, *Diabetologia* 58 (10) (2015) 2206–2217.
- [24] Y. Zhang, H. Zhang, Microbiota associated with type 2 diabetes and its related complications, *Food Sci. Hum. Wellness* 2 (3–4) (2013) 167–172.
- [25] C. Ceapa, H. Wopereis, L. Rezaiki, M. Kleerebezem, J. Knol, R. Oozeer, Influence of fermented milk products, prebiotics and probiotics on microbiota composition and health, *Best Pract. Res. Clin. Gastroenterol.* 27 (1) (2013) 139–155.
- [26] L. Wynnes, Understanding the role of diet in type 2 diabetes prevention, *Br. J. Community Nurs.* 14 (9) (2009) 374–379.
- [27] N.P. Steyn, J. Mann, P.H. Bennett, et al., Diet, nutrition and the prevention of type 2 diabetes, *Public Health Nutr.* 7 (1a) (2004) 147–165.
- [28] E.A. Hu, A. Pan, V. Malik, Q. Sun, White rice consumption and risk of type 2 diabetes: meta-analysis and systematic review, *Bmj* 344 (2012).
- [29] S.C. Larsson, A. Wolk, Potato consumption and risk of cardiovascular disease: 2 prospective cohort studies, *Am. J. Clin. Nutr.* 104 (5) (2016) 1245–1252.
- [30] T.V. Gadiraju, Y. Patel, J.M. Gaziano, L. Djoussé, Fried food consumption and cardiovascular health: a review of current evidence, *Nutrients* 7 (10) (2015) 8424–8430.
- [31] R. Renata Micha, S.K. Wallace, D. Mozaffarian, Red and processed meat consumption and risk of incident coronary heart disease, stroke, and diabetes mellitus, 2010.
- [32] F. Imamura, L. O'Connor, Z. Ye, et al., Consumption of sugar sweetened beverages, artificially sweetened beverages, and fruit juice and incidence of type 2 diabetes: systematic review, meta-analysis, and estimation of population attributable fraction, *Bmj* 351 (2015).
- [33] D. Aune, N. Keum, E. Giovannucci, et al., Whole grain consumption and risk of cardiovascular disease, cancer, and all cause and cause specific mortality: systematic review and dose-response meta-analysis of prospective studies, *bmj* 353 (2016).
- [34] P. Hernández-Alonso, L. Camacho-Barcia, M. Bulló, J. Salas-Salvadó, Nuts and dried fruits: an update of their beneficial effects on type 2 diabetes, *Nutrients* 9 (7) (2017) 673.
- [35] J. Salas-Salvadó, M. Guasch-Ferre, A. Díaz-López, N. Babio, Yogurt and diabetes: overview of recent observational studies, *J. Nutr.* 147 (7) (2017) 1452S–1461S.
- [36] R.M.M. Santos, D.R.A. Lima, Coffee consumption, obesity and type 2 diabetes: a mini-review, *Eur. J. Nutr.* 55 (4) (2016) 1345–1358.
- [37] L. Xing, H. Zhang, R. Qi, R. Tsao, Y. Mine, Recent advances in the understanding of the health benefits and molecular mechanisms associated with green tea polyphenols, *J. Agric. Food Chem.* 67 (4) (2019) 1029–1043.
- [38] E. Patterson, R. Wall, G. Fitzgerald, R. Ross, C. Stanton, Health implications of high dietary omega-6 polyunsaturated fatty acids, *J. Nutr. Metab.* 2012 (2012).
- [39] H. Fujii, M. Iwase, T. Ohkuma, et al., Impact of dietary fiber intake on glycemic control, cardiovascular risk factors and chronic kidney disease in Japanese patients with type 2 diabetes mellitus: the Fukuoka Diabetes Registry, *Nutr. J.* 12 (1) (2013) 1–8.
- [40] M. Barbagallo, L.J. Dominguez, Magnesium and type 2 diabetes, *World J. Diabetes* 6 (10) (2015) 1152.
- [41] J.C. Fernández-Cao, M. Warthon-Medina, V.H. Moran, et al., Zinc intake and status and risk of type 2 diabetes mellitus: a systematic review and meta-analysis, *Nutrients* 11 (5) (2019) 1027.
- [42] J. Mitri, M. Muraru, A. Pittas, Vitamin D and type 2 diabetes: a systematic review, *Eur. J. Clin. Nutr.* 65 (9) (2011) 1005–1015.
- [43] J. Zhou, K. Huang, X.G. Lei, Selenium and diabetes—evidence from animal studies, *Free Radic. Biol. Med.* 65 (2013) 1548–1556.
- [44] L.L. Koppes, J.M. Dekker, H.F. Hendriks, L.M. Bouter, R.J. Heine, Moderate alcohol consumption lowers the risk of type 2 diabetes: a meta-analysis of prospective observational studies, *Diabetes Care* 28 (3) (2005) 719–725.
- [45] Y. Gepner, R. Golan, I. Harman-Boehm, et al., Effects of initiating moderate alcohol intake on cardiometabolic risk in adults with type 2 diabetes: a 2-year randomized, controlled trial, *Ann. Intern. Med.* 163 (8) (2015) 569–579.
- [46] G. Fernandes, J. Tap, T. Bruls, et al., Enterotypes of the human gut microbiome, *Nature* 473 (7346) (2011), 174180.
- [47] T. Yatsunenko, F. Rey, M. Manary, I. Trehan, M. Dominguez-Bello, M. Contreras, & Heath A.C., Human gut microbiome viewed across age and geography nature 486 (7402) (2012) 222.
- [48] M. Kussmann, P. van Bladeren, The extended nutrigenomics—understanding the interplay between the genomes of food, gut microbes, and human host, *Front. Genet.* 2 (2011) 21.
- [49] M.D. Schulz, Ç. Atay, J. Heringer, et al., High-fat-diet-mediated dysbiosis promotes intestinal carcinogenesis independently of obesity, *Nature* 514 (7523) (2014) 508–512.
- [50] A.M. Vaiserman, A.K. Koliada, F. Marotta, Gut microbiota: a player in aging and a target for anti-aging intervention, *Ageing Res. Rev.* 35 (2017) 36–45.
- [51] I. Sekirov, S.L. Russell, L.C.M. Antunes, B.B. Finlay, Gut microbiota in health and disease, *Physiol. Rev.* (2010).
- [52] N.M. Delzenne, A.M. Neyrinck, F. Bäckhed, P.D. Cani, Targeting gut microbiota in obesity: effects of prebiotics and probiotics, *Nat. Rev. Endocrinol.* 7 (11) (2011) 639.
- [53] R. Burcelin, M. Serino, C. Chabo, V. Blasco-Baque, J. Amar, Gut microbiota and diabetes: from pathogenesis to therapeutic perspective, *Acta Diabetol.* 48 (4) (2011) 257–273.
- [54] J. Walter, R. Ley, The human gut microbiome: ecology and recent evolutionary changes, *Annu. Rev. Microbiol.* 65 (2011) 411–429.
- [55] M. Diamant, E. Blaak, W. De Vos, Do nutrient–gut–microbiota interactions play a role in human obesity, insulin resistance and type 2 diabetes? *Obes. Rev.* 12 (4) (2011) 272–281.
- [56] C. Proctor, P. Thiennimitr, N. Chattapakorn, S.C. Chattapakorn, Diet, gut microbiota and cognition, *Metab. Brain Dis.* 32 (1) (2017) 1–17.
- [57] N.-R. Shin, T.W. Whon, J.-W. Bae, Proteobacteria: microbial signature of dysbiosis in gut microbiota, *Trends Biotechnol.* 33 (9) (2015) 496–503.
- [58] S. Carding, K. Verbeke, D.T. Vipond, B.M. Corfe, L.J. Owen, Dysbiosis of the gut microbiota in disease, *Microb. Ecol. Health Dis.* 26 (1) (2015) 26191.
- [59] E. Murphy, P. Cotter, S. Healy, et al., Composition and energy harvesting capacity of the gut microbiota: relationship to diet, obesity and time in mouse models, *Gut* 59 (12) (2010) 1635–1642.
- [60] S.H. Duncan, A. Belenguer, G. Holtrop, A.M. Johnstone, H.J. Flint, G.E. Lobley, Reduced dietary intake of carbohydrates by obese subjects results in decreased concentrations of butyrate and butyrate-producing bacteria in feces, *Appl. Environ. Microbiol.* 73 (4) (2007) 1073–1078.
- [61] J. Tap, J.P. Furet, M. Bensaada, et al., Gut microbiota richness promotes its stability upon increased dietary fibre intake in healthy adults, *Environ. Microbiol.* 17 (12) (2015) 4954–4964.
- [62] P.J. Turnbaugh, V.K. Ridaura, J.J. Faith, F.E. Rey, R. Knight, J.I. Gordon, The effect of diet on the human gut microbiome: a metagenomic analysis in humanized gnotobiotic mice, pp. 6ra14–16ra14, *Sci. Transl. Med.* 1 (6) (2009), pp. 6ra14–16ra14.
- [63] A. Sánchez-Villegas, M.A. Martínez-González, R. Estruch, et al., Mediterranean dietary pattern and depression: the PREDIMED randomized trial, *BMC Med.* 11 (1) (2013) 1–12.
- [64] L. Serra Majem, A. Sánchez Villegas, Mediterranean diet, 2019.
- [65] P.D. Cani, A.M. Neyrinck, F. Fava, et al., Selective increases of bifidobacteria in gut microflora improve high-fat-diet-induced diabetes in mice through a mechanism associated with endotoxaemia, *Diabetologia* 50 (11) (2007) 2374–2383.
- [66] J.-L. Han, H.-L. Lin, Intestinal microbiota and type 2 diabetes: from mechanism insights to therapeutic perspective, *World J. Gastroenterol.* WJG 20 (47) (2014) 17737.
- [67] J. Shen, M.S. Obin, L. Zhao, The gut microbiota, obesity and insulin resistance, *Mol. Asp. Med.* 34 (1) (2013) 39–58.
- [68] P.D. Cani, M. Osto, L. Geurts, A. Everard, Involvement of gut microbiota in the development of low-grade inflammation and type 2 diabetes associated with obesity, *Gut Microbes* 3 (4) (2012) 279–288.
- [69] X. Wu, C. Ma, L. Han, et al., Molecular characterisation of the faecal microbiota in patients with type II diabetes, *Curr. Microbiol.* 61 (1) (2010) 69–78.
- [70] B. Hippe, M. Remely, E. Aumueller, A. Pointner, U. Magnet, A. Haslberger, Faecalibacterium prausnitzii phylogenotypes in type two diabetic, obese, and lean control subjects, *Benef. Microbes* 7 (4) (2016) 511–517.
- [71] F. De Vadder, P. Kovatcheva-Datchary, D. Goncalves, et al., Microbiota-generated metabolites promote metabolic benefits via gut-brain neural circuits, *Cell* 156 (1–2) (2014) 84–96.
- [72] M. Kasubuchi, S. Hasegawa, T. Hiramatsu, A. Ichimura, I. Kimura, Dietary gut microbial metabolites, short-chain fatty acids, and host metabolic regulation, *Nutrients* 7 (2015) 2839–2849.
- [73] G. Den Besten, K. Van Eunen, A.K. Groen, K. Venema, D.-J. Reijngoud, B. M. Bakker, The role of short-chain fatty acids in the interplay between diet, gut microbiota, and host energy metabolism, *J. Lipid Res.* 54 (9) (2013) 2325–2340.
- [74] Z. Gao, J. Yin, J. Zhang, et al., Butyrate improves insulin sensitivity and increases energy expenditure in mice, *Diabetes* 58 (7) (2009) 1509–1517.
- [75] H. Yamashita, K. Fujisawa, E. Ito, et al., Improvement of obesity and glucose tolerance by acetate in Type 2 diabetic Otsuka Long-Evans Tokushima Fatty (OLETF) rats, *Biosci. Biotechnol. Biochem.* 71 (5) (2007) 1236–1243.
- [76] A. Pingitore, E.S. Chambers, T. Hill, et al., The diet-derived short chain fatty acid propionate improves beta-cell function in humans and stimulates insulin secretion from human islets in vitro, *Diabetes Obes. Metab.* 19 (2) (2017) 257–265.
- [77] G. Tolhurst, H. Heffron, Y.S. Lam, et al., Short-chain fatty acids stimulate glucagon-like peptide-1 secretion via the G-protein-coupled receptor FFAR2, *Diabetes* 61 (2) (2012) 364–371.
- [78] Z. Yin, W. Zhang, F. Feng, Y. Zhang, W. Kang, α -Glucosidase inhibitors isolated from medicinal plants, *Food Sci. Hum. Wellness* 3 (3–4) (2014) 136–174.

- [80] B.R. Sharma, V. Kumar, Y. Gat, N. Kumar, A. Parashar, D.J. Pinakin, Microbial maceration: a sustainable approach for phytochemical extraction, *3 Biotech* 8 (9) (2018) 1–13.
- [81] G. Velderrain-Rodríguez, H. Palafox-Carlos, A. Wall-Medrano, et al., Phenolic compounds: their journey after intake, *Food Funct.* 5 (2) (2014) 189–197.
- [82] F. Cardona, C. Andrés-Lacueva, S. Tulipani, F.J. Tinahones, M.I. Queipo-Ortuño, Benefits of polyphenols on gut microbiota and implications in human health, *J. Nutr. Biochem.* 24 (8) (2013) 1415–1422.
- [83] T. Ozdal, D.A. Sela, J. Xiao, D. Boyacioglu, F. Chen, E. Capanoglu, The reciprocal interactions between polyphenols and gut microbiota and effects on bioaccessibility, *Nutrients* 8 (2) (2016) 78.
- [84] M. Duénas, I. Muñoz-González, C. Cueva, et al., A survey of modulation of gut microbiota by dietary polyphenols, *BioMed Res. Int.* 2015 (2015).
- [85] U. Etxeberria, A. Fernández-Quintela, F.I. Milagro, L. Aguirre, J.A. Martínez, M. P. Portillo, Impact of polyphenols and polyphenol-rich dietary sources on gut microbiota composition, *J. Agric. Food Chem.* 61 (40) (2013) 9517–9533.
- [86] M. Mazidi, P. Rezaie, A.P. Kengne, M.G. Mobarhan, G.A. Ferns, Gut microbiome and metabolic syndrome, *Diabetes Metab. Syndr. Clin. Res. Rev.* 10 (2) (2016) S150–S157.
- [87] L.K. Brahe, A. Astrup, L.H. Larsen, Can we prevent obesity-related metabolic diseases by dietary modulation of the gut microbiota? *Adv. Nutr.* 7 (1) (2016) 90–101.
- [88] M.T. Khan, M. Nieuwdorp, F. Bäckhed, Microbial modulation of insulin sensitivity, *Cell Metab.* 20 (5) (2014) 753–760.
- [89] R.A. Luna, J.A. Foster, Gut brain axis: diet microbiota interactions and implications for modulation of anxiety and depression, *Curr. Opin. Biotechnol.* 32 (2015) 35–41.
- [90] H.M. Eid, M.L. Wright, N. Anil Kumar, et al., Significance of microbiota in obesity and metabolic diseases and the modulatory potential by medicinal plant and food ingredients, *Front. Pharmacol.* 8 (2017) 387.
- [91] M. Roberfroid, G.R. Gibson, L. Hoyles, et al., Prebiotic effects: metabolic and health benefits, *Br. J. Nutr.* 104 (S2) (2010) S1–S63.
- [92] D. Vandeputte, G. Falony, S. Vieira-Silva, et al., Prebiotic inulin-type fructans induce specific changes in the human gut microbiota, *Gut* 66 (11) (2017) 1968–1974.
- [93] J.M. Laparra, Y. Sanz, Interactions of gut microbiota with functional food components and nutraceuticals, *Pharmacol. Res.* 61 (3) (2010) 219–225.
- [94] S.G. Parkar, D.E. Stevenson, M.A. Skinner, The potential influence of fruit polyphenols on colonic microflora and human gut health, *Int. J. Food Microbiol.* 124 (3) (2008) 295–298.
- [95] Z. Gao, Q. Li, X. Wu, X. Zhao, L. Zhao, X. Tong, New insights into the mechanisms of chinese herbal products on diabetes: a focus on the “Bacteria-Mucosal Immunity-Inflammation-Diabetes” axis, *J. Immunol. Res.* 2017 (2017).
- [96] B. Zhai, C. Zhang, Y. Sheng, et al., Hypoglycemic and hypolipidemic effect of S-allyl-cysteine sulfoxide (alliin) in DIO mice, *Sci. Rep.* 8 (1) (2018) 1–7.
- [97] K. Chen, K. Xie, Z. Liu, et al., Preventive effects and mechanisms of garlic on dyslipidemia and gut microbiome dysbiosis, *Nutrients* 11 (6) (2019) 1225.
- [98] K. Chen, Y. Nakasone, K. Xie, K. Sakao, D.-X. Hou, Modulation of allicin-free garlic on gut microbiome, *Molecules* 25 (3) (2020) 682.
- [99] H. Yan, J. Lu, Y. Wang, W. Gu, X. Yang, J. Yu, Intake of total saponins and polysaccharides from *Polygonatum kingianum* affects the gut microbiota in diabetic rats, *Phytomedicine* 26 (2017) 45–54.
- [100] Y. Xie, M. Xiao, Y. Ni, et al., *Alpinia oxyphylla* miq. extract prevents diabetes in mice by modulating gut microbiota, *J. Diabetes Res.* 2018 (2018).
- [101] X. Yan, C. Yang, G. Lin, et al., Antidiabetic potential of green seaweed *Enteromorpha prolifera* flavonoids regulating insulin signaling pathway and gut microbiota in type 2 diabetic mice, *J. Food Sci.* 84 (1) (2019) 165–173.
- [102] T.K.C. Le, T. Hosaka, T.T. Nguyen, et al., *Bifidobacterium* species lower serum glucose, increase expressions of insulin signaling proteins, and improve adipokine profile in diabetic mice, *Biomed. Res.* 36 (1) (2015) 63–70.
- [103] C. Picard, J. Fioramonti, A. Francois, T. Robinson, F. Neant, C. Matuchansky, *bifidobacteria* as probiotic agents—physiological effects and clinical benefits, *Aliment. Pharmacol. Ther.* 22 (6) (2005) 495–512.
- [104] S. Yun, H. Park, J. Kang, Effect of *Lactobacillus gasseri* BNR17 on blood glucose levels and body weight in a mouse model of type 2 diabetes, *J. Appl. Microbiol.* 107 (5) (2009) 1681–1686.
- [105] A.C.D. Lima, C. Cecatti, M.P. Fidélis, et al., Effect of daily consumption of orange juice on the levels of blood glucose, lipids, and gut microbiota metabolites: controlled clinical trials, *J. Med. Food* 22 (2) (2019) 202–210.
- [106] F.F. Anhé, D. Roy, G. Pilon, et al., A polyphenol-rich cranberry extract protects from diet-induced obesity, insulin resistance and intestinal inflammation in association with increased *Akkermansia* spp. population in the gut microbiota of mice, *Gut* 64 (6) (2015) 872–883.
- [107] G. Lin, X. Liu, X. Yan, et al., Role of green macroalgae *Enteromorpha prolifera* polyphenols in the modulation of gene expression and intestinal microflora profiles in type 2 diabetic mice, *Int. J. Mol. Sci.* 20 (1) (2019) 25.
- [108] D.N. Frank, A.L.S. Amand, R.A. Feldman, E.C. Boedeker, N. Harpaz, N.R. Pace, Molecular-phylogenetic characterization of microbial community imbalances in human inflammatory bowel diseases, *Proc. Natl. Acad. Sci. USA* 104 (34) (2007) 13780–13785.
- [109] F. Navab-Moghadam, M. Sedighi, M.E. Khamesh, et al., The association of type II diabetes with gut microbiota composition, *Microb. Pathog.* 110 (2017) 630–636.
- [110] G.M. Nava, H.J. Friedrichsen, T.S. Stappenbeck, Spatial organization of intestinal microbiota in the mouse ascending colon, *ISME J.* 5 (4) (2011) 627–638.
- [111] F. Thomas, J.-H. Hehemann, E. Rebuffet, M. Czjzek, G. Michel, Environmental and gut bacteroidetes: the food connection, *Front. Microbiol.* 2 (2011) 93.
- [112] C.G. Buffie, E.G. Pamer, Microbiota-mediated colonization resistance against intestinal pathogens, *Nat. Rev. Immunol.* 13 (11) (2013) 790–801.
- [113] L.R. Lopetuso, F. Scaldaferri, V. Petito, A. Gasbarrini, Commensal Clostridia: leading players in the maintenance of gut homeostasis, *Gut Pathog.* 5 (1) (2013) 1–8.
- [114] P.M. Smith, M.R. Howitt, N. Panikov, et al., The microbial metabolites, short-chain fatty acids, regulate colonic Treg cell homeostasis, *Science* 341 (6145) (2013) 569–573.
- [115] L.A. David, C.F. Maurice, R.N. Carmody, et al., Diet rapidly and reproducibly alters the human gut microbiome, *Nature* 505 (7484) (2014) 559–563.
- [116] W.R. Russell, S.W. Gratz, S.H. Duncan, et al., High-protein, reduced-carbohydrate weight-loss diets promote metabolite profiles likely to be detrimental to colonic health, *Am. J. Clin. Nutr.* 93 (5) (2011) 1062–1072.
- [117] L. Madsen, L.S. Myrmet, E. Fjære, B. Liasset, K. Kristiansen, Links between dietary protein sources, the gut microbiota, and obesity, *Front. Physiol.* 8 (2017) 1047.
- [118] M.A. Hildebrandt, C. Hoffmann, S.A. Sherrill-Mix, et al., High-fat diet determines the composition of the murine gut microbiome independently of obesity, *e1712*, *Gastroenterology* 137 (5) (2009) 1716–1724. e1712.
- [119] J.R. Araújo, J. Tomas, C. Brenner, P.J. Sansonetti, Impact of high-fat diet on the intestinal microbiota and small intestinal physiology before and after the onset of obesity, *Biochimie* 141 (2017) 97–106.
- [120] S. Bibbò, G. Ianiro, V. Giorgio, et al., The role of diet on gut microbiota composition, *Eur. Rev. Med. Pharmacol. Sci.* 20 (22) (2016) 4742–4749.
- [121] A. Everard, V. Lazarevic, M. Derrien, et al., Responses of gut microbiota and glucose and lipid metabolism to prebiotics in genetic obese and diet-induced leptin-resistant mice, *Diabetes* 60 (11) (2011) 2775–2786.
- [122] K. Kameyama, K. Itoh, Intestinal colonization by a Lachnospiraceae bacterium contributes to the development of diabetes in obese mice, *Microbes Environ.* (2014) ME14054.
- [123] G. Allcock, M. Allegra, R. Flower, M. Perretti, Neutrophil accumulation induced by bacterial lipopolysaccharide: effects of dexamethasone and annexin 1, *Clin. Exp. Immunol.* 123 (1) (2001) 62–67.
- [124] N. Larsen, F.K. Vogensen, F.W. Van Den Berg, et al., Gut microbiota in human adults with type 2 diabetes differs from non-diabetic adults, *PLoS One* 5 (2) (2010), e9085.
- [125] J.-P. Furet, L.-C. Kong, J. Tap, et al., Differential adaptation of human gut microbiota to bariatric surgery-induced weight loss: links with metabolic and low-grade inflammation markers, *Diabetes* 59 (12) (2010) 3049–3057.
- [126] C. Zhang, H. Diao, F. Lu, X. Bie, Y. Wang, Z. Lu, Degradation of triphenylmethane dyes using a temperature and pH stable spore laccase from a novel strain of *Bacillus vallismortis*, *Bioresour. Technol.* 126 (2012) 80–86.
- [127] A. Everard, C. Belzer, L. Geurts, et al., Cross-talk between *Akkermansia muciniphila* and intestinal epithelium controls diet-induced obesity, *Proc. Natl. Acad. Sci. USA* 110 (22) (2013) 9066–9071.
- [128] J. De La Cuesta-Zuluaga, N.T. Mueller, V. Corrales-Agudelo, et al., Metformin is associated with higher relative abundance of mucin-degrading *Akkermansia muciniphila* and several short-chain fatty acid-producing microbiota in the gut, *Diabetes Care* 40 (1) (2017) 54–62.
- [129] M. Murri, I. Leiva, J.M. Gomez-Zumaquero, et al., Gut microbiota in children with type 1 diabetes differs from that in healthy children: a case-control study, *BMC Med.* 11 (1) (2013) 1–12.
- [130] H. Yadav, J.-H. Lee, J. Lloyd, P. Walter, S.G. Rane, Beneficial metabolic effects of a probiotic via butyrate-induced GLP-1 hormone secretion, *J. Biol. Chem.* 288 (35) (2013) 25088–25097.
- [131] C.N. Heiss, L.E. Olofsson, Gut microbiota-dependent modulation of energy metabolism, *J. Innate Immun.* 10 (3) (2018) 163–171.
- [132] R. Dentin, J.-P. Pégurier, F. Benhamed, et al., Hepatic glucokinase is required for the synergistic action of ChREBP and SREBP-1c on glycolytic and lipogenic gene expression, *J. Biol. Chem.* 279 (19) (2004) 20314–20326.
- [133] S. Sakakibara, T. Yamauchi, Y. Oshima, Y. Tsukamoto, T. Kadowaki, Acetic acid activates hepatic AMPK and reduces hyperglycemia in diabetic KK-A (y) mice, *Biochem. Biophys. Res. Commun.* 344 (2) (2006) 597–604.
- [134] S.M. Cabarcas, E.M. Hurt, W.L. Farrar, Defining the molecular nexus of cancer, type 2 diabetes and cardiovascular disease, *Curr. Mol. Med.* 10 (8) (2010) 741–755.
- [135] K. Inoki, J. Kim, K.-L. Guan, AMPK and mTOR in cellular energy homeostasis and drug targets, *Annu. Rev. Pharmacol. Toxicol.* 52 (2012) 381–400.
- [136] F. Bäckhed, H. Ding, T. Wang, et al., The gut microbiota as an environmental factor that regulates fat storage, *Proc. Natl. Acad. Sci. USA* 101 (44) (2004) 15718–15723.
- [137] A. Vriee, F. Holleman, E. Zoetendal, W. De Vos, J. Hoekstra, M. Nieuwdorp, The environment within: how gut microbiota may influence metabolism and body composition, *Diabetologia* 53 (4) (2010) 606–613.
- [138] P.D. Cani, S. Possemiers, T. Van de Wiele, et al., Changes in gut microbiota control inflammation in obese mice through a mechanism involving GLP-2-driven improvement of gut permeability, *Gut* 58 (8) (2009) 1091–1103.
- [139] P. Cani, N. Delzenne, Involvement of the gut micro biota in the development of low grade inflammation as associated with obesity: focus on this neglected partner, *Acta Gastroenterol. Belg.* 73 (2010) 267–269.
- [140] A.P.B. Moreira, T.F.S. Teixeira, A.B. Ferreira, M.D.C.G. Peluzio, R.D.C.G. Alfenas, Influence of a high-fat diet on gut microbiota, intestinal permeability and metabolic endotoxaemia, *Br. J. Nutr.* 108 (5) (2012) 801–809.
- [141] M. Vijay-Kumar, J.D. Aitken, F.A. Carvalho, et al., Metabolic syndrome and altered gut microbiota in mice lacking Toll-like receptor 5, *Science* 328 (5975) (2010) 228–231.

- [142] K.M. Maslowski, C.R. Mackay, Diet, gut microbiota and immune responses, *Nat. Immunol.* 12 (1) (2011) 5–9.
- [143] C. Petersen, U.D. Wankhade, D. Bharat, et al., Dietary supplementation with strawberry induces marked changes in the composition and functional potential of the gut microbiome in diabetic mice, *J. Nutr. Biochem.* 66 (2019) 63–69.
- [144] M. Hidalgo, M.J. Oruna-Concha, S. Kolida, et al., Metabolism of anthocyanins by human gut microflora and their influence on gut bacterial growth, *J. Agric. Food Chem.* 60 (15) (2012) 3882–3890.
- [145] M. Horbowicz, R. Kosson, A. Grzesiuk, H. Debski, Anthocyanins of fruits and vegetables—their occurrence, analysis and role in human nutrition, *Veg. Crops Res. Bull.* 68 (2008) 5.
- [146] T.-G. Hu, P. Wen, W.-Z. Shen, et al., Effect of 1-deoxynojirimycin isolated from mulberry leaves on glucose metabolism and gut microbiota in a streptozotocin-induced diabetic mouse model, *J. Nat. Prod.* 82 (8) (2019) 2189–2200.
- [147] T.-G. Hu, P. Wen, J. Liu, et al., Combination of mulberry leaf and oat bran possessed greater hypoglycemic effect on diabetic mice than mulberry leaf or oat bran alone, *J. Funct. Foods* 61 (2019), 103503.
- [148] L. Chen, W. Lu, Y. Li, Berberine ameliorates type 2 diabetes via modulation of *Bifidobacterium* species, tumor necrosis factor- α , and lipopolysaccharide, *Int. J. Clin. Exp. Med.* 9 (6) (2016).
- [149] S. Vendrame, S. Guglielmetti, P. Riso, S. Arioli, D. Klimis-Zacas, M. Porrini, Six-week consumption of a wild blueberry powder drink increases bifidobacteria in the human gut, *J. Agric. Food Chem.* 59 (24) (2011) 12815–12820.
- [150] J.C. Lee, H. Koerten, P. Van den Broek, et al., Adherence of *Acinetobacter baumannii* strains to human bronchial epithelial cells, *Res. Microbiol.* 157 (4) (2006) 360–366.
- [151] H. Sun, Y. Chen, M. Cheng, X. Zhang, X. Zheng, Z. Zhang, The modulatory effect of polyphenols from green tea, oolong tea and black tea on human intestinal microbiota in vitro, *J. Food Sci. Technol.* 55 (1) (2018) 399–407.
- [152] P. Dolara, C. Luceri, C. De Filippo, et al., Red wine polyphenols influence carcinogenesis, intestinal microflora, oxidative damage and gene expression profiles of colonic mucosa in F344 rats, *Mutat. Res. Fundam. Mol. Mech. Mutagen.* 591 (1–2) (2005) 237–246.
- [153] J. Han, H. Lin, W. Huang, Modulating gut microbiota as an anti-diabetic mechanism of berberine, *Med. Sci. Monit. Int. Med. J. Exp. Clin. Res.* 17 (7) (2011) RA164.
- [154] X. Zhang, Y. Zhao, J. Xu, et al., Modulation of gut microbiota by berberine and metformin during the treatment of high-fat diet-induced obesity in rats, *Sci. Rep.* 5 (1) (2015) 1–10.
- [155] X. Zhang, Y. Zhao, M. Zhang, et al., Structural changes of gut microbiota during berberine-mediated prevention of obesity and insulin resistance in high-fat diet-fed rats, *PLoS One* 7 (8) (2012), e42529.
- [156] J. Miyamoto, K. Watanabe, S. Taira, et al., Barley β -glucan improves metabolic condition via short-chain fatty acids produced by gut microbial fermentation in high fat diet fed mice, *PLoS One* 13 (4) (2018), e0196579.
- [157] Y. Zhu, L. Dong, L. Huang, et al., Effects of oat β -glucan, oat resistant starch, and the whole oat flour on insulin resistance, inflammation, and gut microbiota in high-fat-diet-induced type 2 diabetic rats, *J. Funct. Foods* 69 (2020), 103939.
- [158] X. Tzounis, A. Rodriguez-Mateos, J. Vulevic, G.R. Gibson, C. Kwik-Urbe, J. P. Spencer, Prebiotic evaluation of cocoa-derived flavanols in healthy humans by using a randomized, controlled, double-blind, crossover intervention study, *Am. J. Clin. Nutr.* 93 (1) (2011) 62–72.
- [159] M.I. Queipo-Ortuño, M. Boto-Ordóñez, M. Murri, et al., Influence of red wine polyphenols and ethanol on the gut microbiota ecology and biochemical biomarkers, *Am. J. Clin. Nutr.* 95 (6) (2012) 1323–1334.
- [160] J. Zhao, L. Nian, L. Kwok, T. Sun, Reduction in fecal microbiota diversity and short-chain fatty acid producers in methicillin-resistant *Staphylococcus aureus* infected individuals as revealed by PacBio single molecule, real-time sequencing technology, *Eur. J. Clin. Microbiol. Infect. Dis.* 36 (8) (2017) 1463–1472.
- [161] B. Zhang, W. Sun, N. Yu, et al., Anti-diabetic effect of baicalein is associated with the modulation of gut microbiota in streptozotocin and high-fat-diet induced diabetic rats, *J. Funct. Foods* 46 (2018) 256–267.
- [162] X. Tzounis, J. Vulevic, G.G. Kuhnle, et al., Flavanol monomer-induced changes to the human faecal microflora, *Br. J. Nutr.* 99 (4) (2008) 782–792.
- [163] H.M. Eid, M. Ouchfoun, A. Saleem, et al., A combination of (+)-catechin and (–)-epicatechin underlies the in vitro adipogenic action of Labrador tea (*Rhododendron groenlandicum*), an antidiabetic medicinal plant of the Eastern James Bay Cree pharmacopeia, *J. Ethnopharmacol.* 178 (2016) 251–257.
- [164] K. He, Y. Hu, H. Ma, et al., Rhizoma *Coptidis* alkaloids alleviate hyperlipidemia in B6 mice by modulating gut microbiota and bile acid pathways, *Biochim. Biophys. Acta (BBA) Mol. Basis Dis.* 1862 (9) (2016) 1696–1709.
- [165] J.P. Piwowski, A.K. Kiss, S. Granica, T. Moeslinger, Urolithins, gut microbiota-derived metabolites of ellagitannins, inhibit LPS-induced inflammation in RAW 264.7 murine macrophages, *Mol. Nutr. Food Res.* 59 (11) (2015) 2168–2177.
- [166] H. Song, Q. Chu, F. Yan, Y. Yang, W. Han, X. Zheng, Red pitaya betacyanins protects from diet-induced obesity, liver steatosis and insulin resistance in association with modulation of gut microbiota in mice, *J. Gastroenterol. Hepatol.* 31 (8) (2016) 1462–1469.
- [167] L. Han, T. Li, M. Du, R. Chang, B. Zhan, X. Mao, Beneficial effects of potentilla discolor bunge water extract on inflammatory cytokines release and gut microbiota in high-fat diet and streptozotocin-induced type 2 diabetic mice, *Nutrients* 11 (3) (2019) 670.
- [168] M. Van Hul, L. Geurts, H. Plovier, et al., Reduced obesity, diabetes, and steatosis upon cinnamon and grape pomace are associated with changes in gut microbiota and markers of gut barrier, *Am. J. Physiol. Endocrinol. Metab.* 314 (4) (2018) E334–E352.
- [169] R. Gupta, M. Mathur, V.K. Bajaj, et al., Evaluation of antidiabetic and antioxidant activity of *Moringa oleifera* in experimental diabetes, *J. Diabetes* 4 (2) (2012) 164–171.
- [170] J. Wang, Q. Zou, Y. Suo, et al., Lycopene ameliorates systemic inflammation-induced synaptic dysfunction by improving insulin resistance and mitochondrial dysfunction in the liver–brain axis, *Food Funct.* 10 (4) (2019) 2125–2137.
- [171] B. Zhao, J. Wu, J. Li, et al., Lycopene alleviates DSS-induced colitis and behavioral disorders via mediating microbes–gut–brain axis balance, *J. Agric. Food Chem.* 68 (13) (2020) 3963–3975.
- [172] A. Arora, T. Behl, A. Sehgal, S. Singh, N. Sharma, S. Bhatia, E. Sobarzo-Sanchez, S. Bungau, Unravelling the involvement of gut microbiota in type 2 diabetes mellitus, *Life Sci.* 273 (15) (2021), 119311.
- [173] B.Y. Li, X.Y. Xu, R.Y. Gan, Q.C. Sun, J.M. Meng, A. Shang, Q.Q. Mao, H.B. Li, Targeting gut microbiota for the prevention and management of diabetes mellitus by dietary natural products, *Foods* 8 (10) (2019) 440.

Further reading

- [1] J.L. Sonnenburg, F. Bäckhed, Diet–microbiota interactions as moderators of human metabolism, *Nature* 535 (7610) (2016) 56–64.
- [2] Q. Nie, H. Chen, J. Hu, S. Fan, S. Nie, Dietary compounds and traditional Chinese medicine ameliorate type 2 diabetes by modulating gut microbiota, *Crit. Rev. Food Sci. Nutr.* 59 (6) (2019) 848–863.